

REDUCTION OF ONE-LOOP AMPLITUDES
AT THE INTEGRAND LEVEL —
NLO QCD CALCULATIONS* **

GIOVANNI OSSOLA, COSTAS G. PAPADOPOULOS

NCSR Demokritos, 15310, Athens, Greece

ROBERTO PITTAU

Departamento de Fisica Teorica y del Cosmos y CAFPE
Universidad de Granada, 18071 Granada, Spain

(Received May 6, 2008)

The recently proposed method (OPP) to extract the coefficients of the scalar one-loop integrals to any multi-particle (sub)-amplitude is described. Within this method no analytical information on the structure of the amplitude is needed, allowing for a purely numerical, but still algebraic, implementation of the algorithm. The algorithm can be used to automatically perform one-loop calculation both in QCD and in the EW Theory. As an application, we give QCD one-loop results for the process $pp \rightarrow ZZZ$ at the LHC.

PACS numbers: 11.10.-z, 12.38.Bx, 12.15.Lk, 13.85.Hd

1. Introduction

In two recent papers [1], we proposed a reduction method (OPP) for arbitrary one-loop sub-amplitudes at *the integrand level* [2]. The method is based on idea of expressing the integrand of the one-loop amplitude in terms of the propagators that depends on the integration momentum. The solution of this equation can proceed in an hierarchical way, by exploiting numerically the set of kinematical equations for the integration momentum, corresponding to the so-called quadruple, triple and double cuts used in the unitarity-cut method [3–5]. The method requires a minimal information

* Presented by C.G. Papadopoulos at the Cracow Epiphany Conference on LHC Physics, Cracow, Poland, 4–6 January 2008.

** Work supported by HEPTOOLS, MRTN-CT-2006-035505 and ALGOTOOLS, MTKD-CT-2004-014319.

about the form of the one-loop (sub-)amplitude and therefore it is well suited for a numerical implementation. The method works for any set of internal and/or external masses, so that one is able to study the full electroweak model, without being limited to massless theories.

2. The OPP method

The starting point of the OPP reduction method is the general expression for the *integrand* of a generic m -point one-loop (sub-)amplitude

$$A(\bar{q}) = \frac{N(q)}{\bar{D}_0 \bar{D}_1 \cdots \bar{D}_{m-1}}, \quad \bar{D}_i = (\bar{q} + p_i)^2 - m_i^2, \quad p_0 \neq 0. \quad (1)$$

In the previous equation, we use a bar to denote objects living in $n = 4 + \epsilon$ dimensions, and $\bar{q}^2 = q^2 + \tilde{q}^2$, where \tilde{q}^2 is ϵ -dimensional and $(\tilde{q} \cdot q) = 0$. $N(q)$ is the 4-dimensional part of the numerator function of the amplitude. If needed, the ϵ -dimensional part of the numerator should be treated separately, as will be explained later. $N(q)$ depends on the 4-dimensional denominators $D_i = (q + p_i)^2 - m_i^2$ as follows

$$\begin{aligned} N(q) = & \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i \\ & + \sum_{i_0 < i_1 < i_2}^{m-1} [c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2)] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\ & + \sum_{i_0 < i_1}^{m-1} [b(i_0 i_1) + \tilde{b}(q; i_0 i_1)] \prod_{i \neq i_0, i_1}^{m-1} D_i \\ & + \sum_{i_0}^{m-1} [a(i_0) + \tilde{a}(q; i_0)] \prod_{i \neq i_0}^{m-1} D_i \\ & + \tilde{P}(q) \prod_i^{m-1} D_i. \end{aligned} \quad (2)$$

Inserted back in Eq. (1), this expression simply states the multi-pole nature of any m -point one-loop amplitude, that, clearly, contains a pole for any propagator in the loop, thus one has terms ranging from 1 to m poles. The coefficients of the poles can be further split into two pieces. A piece that still depends on q (the terms $\tilde{d}, \tilde{c}, \tilde{b}, \tilde{a}$), that vanishes upon integration, and a piece that does not depend on q (the terms d, c, b, a). Such a separation is

always possible and the latter set of coefficients is immediately interpretable as the ensemble of the coefficients of all possible 4, 3, 2, 1-point one-loop functions contributing to the amplitude.

Once Eq. (2) is established, the task of computing the one-loop amplitude is then reduced to the algebraical problem of fitting the coefficients d, c, b, a by evaluating the function $N(q)$ a sufficient number of times, at different values of q , and then inverting the system. It can be achieved quite efficiently by singling out particular choices of q such that, systematically, 4, 3, 2 or 1 among all possible denominators D_i vanishes. Then the system of equations is solved iteratively¹. First one determines all possible 4-point functions, then the 3-point functions and so on. For example, calling q_0^\pm the two solutions (in general complex) for which

$$D_0 = D_1 = D_2 = D_3 = 0, \quad (3)$$

(there are 2 solutions because of the quadratic nature of the propagators) and since the functional form of $\tilde{d}(q; 0123)$ is known, one directly finds the coefficient of the box diagram containing the above 4 denominators through the two simple equations

$$N(q_0^\pm) = \left[d(0123) + \tilde{d}(q_0^\pm; 0123) \right] \prod_{i \neq 0,1,2,3} D_i(q_0^\pm). \quad (4)$$

This algorithm also works in the case of complex denominators, namely with complex masses. Notice that the described procedure can be performed *at the amplitude level*. One does not need to repeat the work for all Feynman diagrams, provided their sum is known: we just suppose to be able to compute $N(q)$ numerically.

As is well known, even starting from a perfectly finite tensor integral, the tensor reduction may eventually lead to integrals that need to be regularized (we use dimensional regularization). Such tensors are finite, but tensor reduction iteratively leads to rank m m -point tensors with $1 \leq m \leq 5$, that are ultraviolet divergent when $m \leq 4$. For this reason, we introduced, in Eq. (1), the d -dimensional denominators \bar{D}_i , that differs by an amount \tilde{q}^2 from their 4-dimensional counterparts

$$\bar{D}_i = D_i + \tilde{q}^2. \quad (5)$$

The result of this is a mismatch in the cancellation of the d -dimensional denominators of Eq. (1) with the 4-dimensional ones of Eq. (2). The rational part of the amplitude, called R_1 [7], comes from such a lack of cancellation.

¹ An interesting method to optimize the solution of the system has been very recently presented in [6].

A different source of Rational Terms, called R_2 , can also be generated from the ϵ -dimensional part of $N(q)$ (that is missing in Eq. (1)). For the time being, it should be added by hand by looking at the analytical structure of the Feynman Diagrams or via a dedicated set of Feynman Rules. Examples on how to compute R_2 are reported in [7] and [8]. The Rational Terms R_1 are generated by the following extra integrals, introduced in [1]

$$\begin{aligned} \int d^n \bar{q} \frac{\tilde{q}^2}{\bar{D}_i \bar{D}_j} &= -\frac{i\pi^2}{2} \left[m_i^2 + m_j^2 - \frac{(p_i - p_j)^2}{3} \right] + \mathcal{O}(\epsilon), \\ \int d^n \bar{q} \frac{\tilde{q}^2}{\bar{D}_i \bar{D}_j \bar{D}_k} &= -\frac{i\pi^2}{2} + \mathcal{O}(\epsilon), \quad \int d^n \bar{q} \frac{\tilde{q}^4}{\bar{D}_i \bar{D}_j \bar{D}_k \bar{D}_l} = -\frac{i\pi^2}{6} + \mathcal{O}(\epsilon). \end{aligned} \quad (6)$$

The coefficients of the above integrals can be computed by looking at the implicit mass dependence (namely reconstructing the \tilde{q}^2 dependence) in the coefficients d, c, b of the one-loop functions, once \tilde{q}^2 is reintroduced through the mass shift $m_i^2 \rightarrow m_i^2 - \tilde{q}^2$. One gets

$$b(ij; \tilde{q}^2) = b(ij) + \tilde{q}^2 b^{(2)}(ij), \quad c(ijk; \tilde{q}^2) = c(ijk) + \tilde{q}^2 c^{(2)}(ijk). \quad (7)$$

Furthermore, by defining

$$\mathcal{D}^{(m)}(q, \tilde{q}^2) \equiv \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[d(i_0 i_1 i_2 i_3; \tilde{q}^2) + \tilde{d}(q; i_0 i_1 i_2 i_3; \tilde{q}^2) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} \bar{D}_i, \quad (8)$$

the following expansion holds

$$\mathcal{D}^{(m)}(q, \tilde{q}^2) = \sum_{j=2}^m \tilde{q}^{(2j-4)} d^{(2j-4)}(q), \quad (9)$$

where the last coefficient is independent on q , $d^{(2m-4)}(q) = d^{(2m-4)}$. In practice, once the 4-dimensional coefficients have been determined, one can redo the fits for different values of \tilde{q}^2 , in order to determine $b^{(2)}(ij)$, $c^{(2)}(ijk)$ and $d^{(2m-4)}$. Such three quantities are the coefficients of the three extra scalar integrals listed in Eq. (6), respectively. Therefore, the OPP method allows an easy and purely numerical computation of the Rational Terms of type R_1 .

3. $pp \rightarrow ZZZ$ at NLO

The calculation is composed of two parts: the evaluation of virtual corrections, namely one-loop contributions obtained by adding a virtual particle to the tree-order diagrams, and corrections from the real emission of one additional massless particle from initial and final states, which is necessary in

order to control and cancel infrared singularities. The virtual corrections are computed using the **OPP** reduction [1]. In particular, we make use of **CutTools** [9]. Concerning the contributions coming from real emission we used the dipole subtraction method [10] to isolate the soft and collinear divergences and checked the results using the phase space slicing method [11] with soft and collinear cutoffs, as outlined in [12].

These results have been already obtained, following a very different approach, by Lazopoulos *et al.* in Ref. [13]. We also presented some preliminary results in [14]. A more complete study, that also includes the case of W^+W^-Z [15], W^+ZZ , and $W^+W^-W^+$ production, has been recently presented in [16].

Let us begin with the evaluation of the virtual QCD corrections to the process $q\bar{q} \rightarrow ZZZ$. We consider the process

$$q(p_1) + \bar{q}(p_2) \longrightarrow Z(p_3) + Z(p_4) + Z(p_5). \quad (10)$$




Fig. 1. Diagrams contributing to virtual QCD corrections to $q\bar{q} \rightarrow ZZZ$.

At the tree-level, there are six contributions to this process, obtained by permuting the final legs in all possible ways. One-loop corrections are obtained by adding a virtual gluon to the tree-level structures. The calculation involves the reduction of 48 diagrams.

We perform a reduction to scalar integrals using the **OPP** reduction method [1]. The coefficients determined in this manner should be multiplied by the corresponding scalar integrals. Since, in the process that we are studying, no q -dependent massive propagator appears, we will only need massless scalar integrals. They are computed using the package **OneLOop** written by van Hameren [17].

We turn now to the calculation of Rational Terms. As explained in Section 2, part of this contribution, that we call R_1 , is automatically included by the reduction algorithm. The second term R_2 , coming from the explicitly ϵ -dimensional part of the amplitude, has been added after having been computed separately; it turns out that only three- and two-point functions contribute and the result is proportional to the tree-order amplitude.

We checked that our results, agree with the results obtained by the authors of Ref. [13].

In what concerns the real emission, we only have to deal with initial state singularities, where we distinguish $q\bar{q}$ and qg initial states. For the qg initial state, no soft singularity is present because the corresponding tree-level contribution vanishes. We recall that the structure of the NLO partonic cross sections is as follows:

$$\begin{aligned}\sigma_{q\bar{q}}^{\text{NLO}} &= \int_{VVV} \left[d\sigma_{q\bar{q}}^B + d\sigma_{q\bar{q}}^V + d\sigma_{q\bar{q}}^C + \int_g d\sigma_{q\bar{q}}^A \right] + \int_{VVVg} \left[d\sigma_{q\bar{q}}^R - d\sigma_{q\bar{q}}^A \right], \\ \sigma_{qg}^{\text{NLO}} &= \int_{VVV} \left[+d\sigma_{qg}^C \int_g d\sigma_{qg}^A \right] + \int_{VVVg} \left[d\sigma_{qg}^R - d\sigma_{qg}^A \right],\end{aligned}\quad (11)$$

where $d\sigma^B, d\sigma^V, d\sigma^C, d\sigma^R, d\sigma^A$ are respectively the Born cross section, the virtual, virtual counterterm, real and real-subtraction cross sections. For the $q\bar{q}$ initial state two dipoles are needed as subtraction terms. If p_6 is the momentum which can become soft or collinear, the dipole term for gluon emission off the quark is given by

$$\begin{aligned}\mathcal{D}^{q_1g_6,\bar{q}_2} &= \frac{8\pi\alpha_s C_F}{2\tilde{x} p_1 \cdot p_6} \left(\frac{1+\tilde{x}^2}{1-\tilde{x}} \right) |\mathcal{M}_{q\bar{q}}^B(\{\tilde{p}\})|^2, \\ \tilde{x} &= \frac{p_1 \cdot p_2 - p_2 \cdot p_6 - p_1 \cdot p_6}{p_1 \cdot p_2},\end{aligned}\quad (12)$$

where the $\{\tilde{p}\}$ are redefined momenta, $\{\tilde{p}_j\} = \{\tilde{p}_{16}, \tilde{p}_2, \tilde{p}_3, \tilde{p}_4, \tilde{p}_5\}$, which are again on-shell and go to $\{p_1, \dots, p_5\}$ in the singular limit, *e.g.* $\tilde{p}_{16} = \tilde{x} p_1$. The regularised real emission part then reads

$$d\sigma_{q\bar{q}}^R - d\sigma_{q\bar{q}}^A = \frac{1}{6} \frac{1}{N} \frac{1}{2s_{12}} \left[C_F |\mathcal{M}_{q\bar{q}}^R(\{p_j\})|^2 - \mathcal{D}^{q_1g_6,\bar{q}_2} - \mathcal{D}^{\bar{q}_2g_6,q_1} \right] d\Phi_{VVVg},$$

where the factor $1/6$ accounts for the three identical bosons in the final state. More details can be found in [10, 16].

The hadronic differential cross section with hadron momenta P_1 and P_2 is the sum over all partonic initial states convoluted with the parton distribution functions

$$d\sigma(P_1, P_2) = \sum_{ab} \int dz_1 dz_2 f_a(z_1, \mu_F) f_b(z_2, \mu_F) d\sigma_{ab}(z_1 P_1, z_2 P_2), \quad (13)$$

where the sum runs over the partonic configurations $q\bar{q}$, $\bar{q}q$, gq , qg , $g\bar{q}$, $\bar{q}g$.

As an explicit example we present the numerical results for the case $u\bar{u} \rightarrow ZZZ$ for $\sqrt{s} = 14$ TeV and using CTEQ6L1 [18]. Tree-order cross section has been evaluated using the **HELAC** event generator [19]. The same generator has been appropriately developed and used for the calculation of the real corrections as outlined above. In the following table the results in fb are presented for the tree-order cross section σ_0 , the ratio of the virtual to the tree-level cross section, and the real contribution, combining 5- and 6-point contributions, as described above, for all channels, *i.e.*, $u\bar{u}$, ug , $g\bar{u}$, for different values of the factorization(renormalization) scale ($\mu = \mu_F = \mu_R$).

Scale	σ_0	σ_V/σ_0	σ_R	σ_{NLO}
$\mu = M_Z$	1.481(5)	0.536(1)	0.238(2)	2.512(2)
$\mu = 2M_Z$	1.487(5)	0.481(1)	0.232(2)	2.434(2)
$\mu = 3M_Z$	1.477(5)	0.452(1)	0.232(2)	2.376(2)
$\mu = 4M_Z$	1.479(5)	0.436(1)	0.232(2)	2.355(2)
$\mu = 5M_Z$	1.479(5)	0.424(1)	0.237(2)	2.343(2)




Fig. 2. Scale dependence of the total cross section.

As is evident from these results, the K -factor is quite sizeable (1.58–1.69), whereas the dependence on the scale μ (see also Fig. 2) is for both cases quite weak, due mainly to the electroweak character of the process.

4. Conclusions

In conclusion we have presented, a reduction method at the integrand level which is changing the way we are looking at the NLO calculations: a full numerical but still algebraic method has been born.

The efficiency of the OPP is quite good. The main future improvement is the efficiency with which the one-loop amplitude, at the integrand level, is computed.

Finally, taking into account the speed, precision and easiness of the OPP method, a universal NLO calculator/event-generator seems feasible.

REFERENCES

- [1] G. Ossola, C.G. Papadopoulos, R. Pittau, *Nucl. Phys.* **B763**, 147 (2007) [arXiv:hep-ph/0609007]; *J. High Energy Phys.* **0707**, 085 (2007) [arXiv:0704.1271[hep-ph]].
- [2] F. del Aguila, R. Pittau, *J. High Energy Phys.* **0407**, 017 (2004) [arXiv:hep-ph/0404120].
- [3] Z. Bern, L.J. Dixon, D.C. Dunbar, D.A. Kosower, *Nucl. Phys.* **B435**, 59 (1995) [arXiv:hep-ph/9409265].
- [4] R. Britto, F. Cachazo, B. Feng, *Nucl. Phys.* **B725**, 275 (2005) [arXiv:hep-th/0412103].
- [5] W.B. Kilgore, arXiv:0711.5015[hep-ph]; R.K. Ellis, W.T. Giele, Z. Kunszt, *J. High Energy Phys.* **0803**, 003 (2008) [arXiv:0708.2398[hep-ph]]; R. Britto, B. Feng, P. Mastrolia, arXiv:0803.1989[hep-ph]; W.T. Giele, Z. Kunszt, K. Melnikov, *J. High Energy Phys.* **0804**, 049 (2008) [arXiv:0801.2237[hep-ph]]; M. Moretti, F. Piccinini, A.D. Polosa, arXiv:0802.4171[hep-ph].
- [6] P. Mastrolia, G. Ossola, C.G. Papadopoulos, R. Pittau, arXiv:0803.3964[hep-ph].
- [7] G. Ossola, C.G. Papadopoulos, R. Pittau, *J. High Energy Phys.* **0805**, 004 (2008) [arXiv:0802.1876[hep-ph]].
- [8] R. Pittau, *Comput. Phys. Commun.* **104**, 23 (1997) [arXiv:hep-ph/9607309]; **111**, 48 (1998) [arXiv:hep-ph/9712418].
- [9] G. Ossola, C.G. Papadopoulos, R. Pittau, *J. High Energy Phys.* **0803**, 042 (2008) [arXiv:0711.3596[hep-ph]].
- [10] S. Catani, M.H. Seymour, *Nucl. Phys.* **B485**, 291 (1997); Erratum ibid. *Nucl. Phys.* **B510**, 503 (1998) [arXiv:hep-ph/9605323].
- [11] W.T. Giele, E.W. N. Glover, D.A. Kosower, *Nucl. Phys.* **B403**, 633 (1993) [arXiv:hep-ph/9302225].

- [12] B.W. Harris, J.F. Owens, *Phys. Rev.* **D65**, 094032 (2002) [[arXiv:hep-ph/0102128](#)].
- [13] A. Lazopoulos, K. Melnikov, F. Petriello, *Phys. Rev.* **D76**, 014001 (2007) [[arXiv:hep-ph/0703273](#)].
- [14] Z. Bern *et al.*, [arXiv:0803.0494\[hep-ph\]](#).
- [15] V. Hankele, D. Zeppenfeld, *Phys. Lett.* **B661**, 1038 (2008) [[arXiv:0712.3544\[hep-ph\]](#)].
- [16] T. Binoth, G. Ossola, C.G. Papadopoulos, R. Pittau, [arXiv:0804.0350\[hep-ph\]](#).
- [17] A. van Hameren, J. Vollinga, S. Weinzierl, *Eur. Phys. J.* **C41**, 361 (2005) [[arXiv:hep-ph/0502165](#)].
- [18] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P. Nadolsky, W.K. Tung, *J. High Energy Phys.* **0207**, 012 (2002) [[arXiv:hep-ph/0201195](#)].
- [19] A. Kanaki, C.G. Papadopoulos, *Comput. Phys. Commun.* **132**, 306 (2000) [[arXiv:hep-ph/0002082](#)]; A. Kanaki, C.G. Papadopoulos, [arXiv:hep-ph/0012004](#); A. Cafarella, C.G. Papadopoulos, M. Worek, [arXiv:0710.2427\[hep-ph\]](#).