

Jacques Distler, Skip Garibaldi

# There is No “Theory of Everything” Inside $E_8$

Received: 30 July 2009 / Accepted: 26 October 2009  
© Springer-Verlag 2010

**Abstract** We analyze certain subgroups of real and complex forms of the Lie group  $E_8$ , and deduce that any “Theory of Everything” obtained by embedding the gauge groups of gravity and the Standard Model into a real or complex form of  $E_8$  lacks certain representation-theoretic properties required by physical reality. The arguments themselves amount to representation theory of Lie algebras in the spirit of Dynkin’s classic papers and are written for mathematicians.

## 1 Introduction

Recently, the preprint (1) by Garrett Lisi has generated a lot of popular interest. It boldly claims to be a sketch of a “Theory of Everything”, based on the idea of combining the local Lorentz group and the gauge group of the Standard Model in a real form of  $E_8$  (necessarily not the compact form, because it contains a group isogenous to  $SL(2, \mathbb{C})$ ). The purpose of this paper is to explain some reasons why an entire class of such models—which include the model in (1)—cannot work, using mostly mathematics with relatively little input from physics.

The mathematical set up is as follows. Fix a real Lie group  $E$ . We are interested in subgroups  $SL(2, \mathbb{C})$  and  $G$  of  $E$  so that:

$G$  is connected, compact, and centralizes  $SL(2, \mathbb{C})$ . (ToE1)

We complexify and then decompose  $\text{Lie}(E) \otimes \mathbb{C}$  as a direct sum of representations of  $SL(2, \mathbb{C})$  and  $G$ . We identify  $SL(2, \mathbb{C}) \otimes_{\mathbb{R}} \mathbb{C}$  with  $SL_{2, \mathbb{C}} \times SL_{2, \mathbb{C}}$  and write

$$\text{Lie}(E) = \bigoplus_{m, n \geq 1} m \otimes n \otimes V_{m, n}, \quad (1.1)$$

---

Theory Group, Department of Physics, and Texas Cosmology Center, University of Texas, Austin, TX 78712, USA. distler@golem.ph.utexas.edu · Department of Mathematics & Computer Science, 400 Dowman Dr., Emory University, Atlanta, GA 30322, USA. skip@member.ams.org; skip@mathcs.emory.edu

where  $m$  and  $n$  denote the irreducible representation of  $\mathrm{SL}_{2,\mathbb{C}}$  of that dimension and  $V_{m,n}$  is a complex representation of  $G \otimes_{\mathbb{R}} \mathbb{C}$ . (Physicists would usually write  $2$  and  $\bar{2}$  instead of  $2 \otimes 1$  and  $1 \otimes 2$ .) Of course,

$$\overline{m \otimes n \otimes V_{m,n}} \simeq n \otimes m \otimes \overline{V_{m,n}},$$

and since the action of  $\mathrm{SL}(2, \mathbb{C}) \cdot G$  on  $\mathrm{Lie}(E)$  is defined over  $\mathbb{R}$ , we deduce that  $\overline{V_{m,n}} \simeq V_{n,m}$ . We further demand that

$$V_{m,n} = 0 \quad \text{if } m+n > 4, \text{ and} \quad (\text{ToE2})$$

$$V_{2,1} \text{ is a complex representation of } G. \quad (\text{ToE3})$$

We recall the definition of complex representation and explain the physical motivation for these hypotheses in the next section. Roughly speaking, (ToE1) is a trivial requirement based on trying to construct a Theory of Everything along the lines suggested by Lisi, (ToE2) is the requirement that the model not contain any “exotic” higher-spin particles, and (ToE3) is the statement that the gauge theory (with gauge group  $G$ ) is *chiral*, as required by the Standard Model. In fact, physics requires slightly stronger hypotheses on  $V_{m,n}$ , for  $m+n=4$ . We will not impose the stronger version of (ToE2).

**Definition 11** *A candidate ToE subgroup of a real Lie group  $E$  is a subgroup generated by a copy of  $\mathrm{SL}(2, \mathbb{C})$  and a subgroup  $G$  such that (ToE1) and (ToE2) hold. A ToE subgroup is a candidate ToE subgroup for which (ToE3) also holds.*

Our main result is:

**Theorem 12** *There are no ToE subgroups in (the transfer of) the complex  $E_8$  nor in any real form of  $E_8$ .*

## Notation

Unadorned Lie algebras and Lie groups mean ones over the real numbers. We use a subscript  $\mathbb{C}$  to denote complex Lie groups—e.g.,  $\mathrm{SL}_{2,\mathbb{C}}$  is the (complex) group of 2-by-2 complex matrices with determinant 1. We can view a  $d$ -dimensional complex Lie group  $G_{\mathbb{C}}$  as a  $2d$ -dimensional real Lie group, which we denote by  $R(G_{\mathbb{C}})$ . (Algebraists call this operation the “transfer” or “Weil restriction of scalars”; geometers, and many physicists, call this operation “realification.”) We use the popular notation of  $\mathrm{SL}(2, \mathbb{C})$  for the transfer  $R(\mathrm{SL}_{2,\mathbb{C}})$  of  $\mathrm{SL}_{2,\mathbb{C}}$ ; it is a double covering of the “restricted Lorentz group”, i.e., of the identity component  $\mathrm{SO}(3, 1)_0$  of  $\mathrm{SO}(3, 1)$ .

## Strategy and main results

Our strategy for proving Theorem 12 will be as follows. We will first catalogue, up to conjugation, all possible embeddings of  $SL(2, \mathbb{C})$  satisfying the hypotheses of (ToE2). The list is remarkably short. Specifically, for every candidate ToE subgroup of  $E$ , the group  $G$  is contained in the maximal compact, connected subgroup  $G_{\max}$  of the centralizer of  $SL(2, \mathbb{C})$  in  $E$ . The proof of Theorem 12 shows that the only possibilities are:

$E$	$G_{\max}$	$V_{2,1}$	
$E_{8(-24)}$	$Spin(11)$	32	
$E_{8(8)}$	$Spin(5) \times Spin(7)$	$(4, 8)$	
$E_{8(-24)}$	$Spin(9) \times Spin(3)$	$(16, 2)$	(1.2)
$R(E_{8, \mathbb{C}})$	$E_7$	56	
$R(E_{8, \mathbb{C}})$	$Spin(12)$	$32 \oplus 32'$	
$R(E_{8, \mathbb{C}})$	$Spin(13)$	64	

We then note that the representation  $V_{2,1}$  of  $G_{\max}$  (and hence, of any  $G \subseteq G_{\max}$ ) has a self-conjugate structure. In other words, (ToE3) fails.

## 2 Physics Background

One of the central features of modern particle physics is that the world is described by a *chiral gauge theory*.

**Definition 21** *Let  $M$  be a four-dimensional pseudo-Riemannian manifold, of signature  $(3, 1)$ , which we will take to be oriented, time-oriented and spin. Let  $G$  be a compact Lie group. The data of a **gauge theory on  $M$  with gauge group  $G$**  consists of a connection,  $A$ , on a principal  $G$ -bundle,  $P \rightarrow M$ , and some “matter fields” transforming as sections of vector bundle(s) associated to unitary representations of  $G$ .*

Of particular interest are the *fermions* of the theory. The orthonormal frame bundle of  $M$  is a principal  $SO(3, 1)_0$  bundle. A choice of spin structure defines a lift to a principal  $Spin(3, 1)_0 = SL(2, \mathbb{C})$  bundle. Let  $S_{\pm} \rightarrow M$  be the irreducible spinor bundles, associated, via the defining two-dimensional representation and its complex conjugate, to this  $SL(2, \mathbb{C})$  principal bundle.

The *fermions of our gauge theory* are denoted

$$\psi \in \Gamma(S_+ \otimes V), \quad \bar{\psi} \in \Gamma(S_- \otimes \bar{V}),$$

where  $V \rightarrow M$  is a vector bundle associated to a (typically reducible) representation  $R$  of  $G$ .

**Definition 22** *Consider,  $V$ , a unitary representation of  $G$  over  $\mathbb{C}$ —i.e., a homomorphism  $G \rightarrow U(V)$ —and an antilinear map  $J: V \rightarrow V$  that commutes with the action of  $G$ . The map  $J$  is called a **real structure** on  $V$  if  $J^2 = 1$ ; physicists call a representation possessing a real structure **real**. The map  $J$  is called a **quaternionic structure** on  $V$  if  $J^2 = -1$ ; physicists call a representation possessing a quaternionic structure **pseudoreal**.*

Subsuming these two subcases, we will say that  $V$  **has a self-conjugate structure** if there exists an antilinear map  $J : V \rightarrow V$  commuting with the action of  $G$  and satisfying  $J^4 = 1$ . Physicists call a representation  $V$  that does not possess a self-conjugate structure **complex**.

*Remark 23* We sketch how to translate the above definition into the language of algebraic groups and Galois descent as in (2) and (3, X.2). Let  $G$  be an algebraic group over  $\mathbb{R}$  and fix a representation  $\rho : G \otimes \mathbb{C} \rightarrow \mathrm{GL}(V)$  for some complex vector space  $V$ . Let  $J$  be an antilinear map  $V \rightarrow V$  that satisfies

$$\rho(g) = J^{-1} \rho(\bar{g}) J \quad \text{for } g \in G(\mathbb{C}). \quad (2.1)$$

We define real, quaternionic, etc., by copying the second and third sentences verbatim from Definition 22.

(In the special case where  $G$  is compact, there is necessarily a positive-definite invariant hermitian form on  $V$  and  $\rho$  arises by complexifying some map  $G \rightarrow \mathrm{U}(V)$ ; this puts us back in the situation of Def. 22. In the special case where  $G$  is connected, the hypothesis from Def. 22 that  $J$  commutes with  $G(\mathbb{R})$ —which is obviously implied by Eq. (2.1)—is actually equivalent to Eq. (2.1). Indeed, both sides of Eq. (2.1) are morphisms of varieties over  $\mathbb{C}$ , so if they agree on  $G(\mathbb{R})$ —which is Zariski-dense by (2, 18.2(ii))—then they are equal on  $G(\mathbb{C})$ .)

If  $V$  has a real structure  $J$ , then the  $\mathbb{R}$ -subspace  $V'$  of elements of  $V$  fixed by  $J$  is a real vector space and  $V$  is canonically identified with  $V' \otimes \mathbb{C}$  so that  $J(v' \otimes z) = v' \otimes \bar{z}$  for  $v' \in V'$  and  $z \in \mathbb{C}$ ; this is Galois descent. Because  $\rho$  commutes with complex conjugation (which acts in the obvious manner on  $G(\mathbb{C})$  and via  $J$  on  $V$ ), it is the complexification of a homomorphism  $\rho' : G \rightarrow \mathrm{GL}(V')$  defined over  $\mathbb{R}$  by (2, AG.14.3). Conversely, if there is a representation  $(V', \rho')$  whose complexification is  $(V, \rho)$ , then taking  $J$  to be complex conjugation on  $V = V' \otimes \mathbb{C}$  defines a real structure on  $(V, \rho)$ .

If  $V$  has a quaternionic structure  $J$ , then we define a real structure  $\hat{J}$  on  $\hat{V} := V \oplus V$  via  $\hat{J}(v_1, v_2) := (Jv_2, -Jv_1)$ .

Finally, suppose that  $G$  is reductive and  $V$  is irreducible (as a representation over  $\mathbb{C}$ , of course). Then by (4, 7), there is a unique irreducible real representation  $W$  whose complexification  $W \otimes \mathbb{C}$  contains  $V$  as a summand. By Schur,  $\mathrm{End}_G(W)$  is a division algebra, and we have three possibilities:

- $\mathrm{End}_G(W) = \mathbb{R}$ ,  $W \otimes \mathbb{C} \simeq V$ , and  $V$  has a real structure.
- $\mathrm{End}_G(W) = \mathbb{H}$ ,  $W \otimes \mathbb{C} \simeq V \oplus V$ , and  $V$  has a quaternionic structure.
- $\mathrm{End}_G(W) = \mathbb{C}$ ,  $W \otimes \mathbb{C} \simeq V \oplus \bar{V}$  where  $V \not\simeq \bar{V}$ , and  $V$  is complex.

We have stated this remark for  $G$  a group over  $\mathbb{R}$ , but all of it generalizes easily to the case where  $G$  is reductive over a field  $F$  and is split by a quadratic extensions  $K$  of  $F$ .

**Definition 24** A gauge theory, with gauge group  $G$ , is said to be **chiral** if the representation  $R$  by which the fermions (21) are defined is complex in the above sense. By contrast, a gauge theory is said to be **nonchiral** if the representation  $R$  in 21 has a self-conjugate structure.

Note that whether a gauge theory is chiral depends crucially on the choice of  $G$ . A gauge theory might be chiral for gauge group  $G$ , but *nonchiral* for a subgroup

$H \subset G$ . That is, there can be a self-conjugate structure on  $R$  compatible with  $H$ , even though no such structure exists that is compatible with the full group  $G$ .

Conversely, suppose that a gauge theory is nonchiral for the gauge group  $G$ . It is also necessarily nonchiral for any gauge group  $H \subset G$ .

## GUTs

The Standard Model is a chiral gauge theory with gauge group

$$G_{\text{SM}} := (\text{SU}(3) \times \text{SU}(2) \times \text{U}(1))/(\mathbb{Z}/6\mathbb{Z}).$$

Various grand unified theories (GUTs) proceed by embedding  $G_{\text{SM}}$  in some (usually simple) group,  $G_{\text{GUT}}$ . Popular choices for  $G_{\text{GUT}}$  are  $\text{SU}(5)$  (5),  $\text{Spin}(10)$ ,  $E_6$ , and the Pati-Salam group,  $(\text{Spin}(6) \times \text{Spin}(4))/(\mathbb{Z}/2\mathbb{Z})$  (6).

It is easiest to explain what the fermion representation of  $G_{\text{SM}}$  is after embedding  $G_{\text{SM}}$  in  $G_{\text{GUT}} := \text{SU}(5)$ . Let  $W$  be the five-dimensional defining representation of  $\text{SU}(5)$ . The representation  $R$  from 21 is the direct sum of three copies of

$$R_0 = \wedge^2 W \oplus \overline{W}.$$

Each such copy is called a “generation” and is 15-dimensional. One identifies each of the 15 weights of  $R_0$  with left-handed fermions: 6 quarks (two in a doublet, each in three colors), two leptons (e.g., the electron and its neutrino), 6 antiquarks, and a positron. With three generations,  $R$  is 45-dimensional.

**Definition 25** *As a generalization, physicists sometimes consider the  $n$ -generation Standard Model, which is defined in similar fashion, but with  $R = R_0^{\oplus n}$ . The  $n$ -generation Standard Model is a chiral gauge theory, for any positive  $n$ . Particle physics, in the real world, is described by “the” Standard Model, which is the case  $n = 3$ .*

For the other choices of GUT group, the analogue of a generation ( $R_0$ ) is higher-dimensional, containing additional fermions that are not seen at low energies. When decomposed under  $G_{\text{SM}} \subset G_{\text{GUT}}$ , the representation decomposes as  $R_0 + R'$ , where  $R'$  is a real representation of  $G_{\text{SM}}$ . In  $\text{Spin}(10)$ , a generation is the 16-dimensional half-spinor representation. In  $E_6$ , it is a 27-dimensional representation, and for the Pati-Salam group it is the  $(4, 1, 2) \oplus (\overline{4}, 2, 1)$  representation. In each case, these representations are complex representations (in the above sense) of  $G_{\text{GUT}}$ , and the complex-conjugate representation is called an “anti-generation.”

## 3 Lisi’s Proposal from [1]

In the previous section, we have described a chiral gauge theory in a fixed (pseudo) Riemannian structure on  $M$ . Lisi’s proposal (1) is to try to combine the spin connection on  $M$  and the gauge connection on  $P$  into a single dynamical framework. This motivates Definition 11 of a ToE subgroup.

More precisely, following (1), we fix subgroups  $\text{SL}(2, \mathbb{C})$  and  $G$ —say, with  $G = G_{\text{SM}}$ —satisfying (ToE1) in some real Lie group  $E$ . The action of the central

element  $-1 \in \mathrm{SL}(2, \mathbb{C})$  provides a  $\mathbb{Z}/2\mathbb{Z}$ -grading on the Lie algebra of  $E$ . This  $\mathbb{Z}/2\mathbb{Z}$ -grading allows one to define a sort of superconnection associated to  $E$  (precisely what sort of superconnection is explained in a blog post by the first author (7)). In the proposal of (1), we are supposed to identify each of the generators of  $\mathrm{Lie}(E)$  as either a boson or a fermion. (See Table 9 in (1) for an identification of the 240 roots.)

The Spin-Statistics Theorem (8) says that fermions transform as spinorial representations of  $\mathrm{Spin}(3, 1)$ ; bosons transform as “tensorial” representations (representations which lift to the double cover,  $\mathrm{SO}(3, 1)$ ). To be consistent with the Spin-Statistics Theorem, we must, therefore, require that the fermions belong to the  $-1$ -eigenspace of the aforementioned  $\mathbb{Z}/2\mathbb{Z}$  action, and the bosons to the  $+1$ -eigenspace.

In fact, to agree with 21, we should require that the  $-1$ -eigenspace (when tensored with  $\mathbb{C}$ ) decomposes as a direct sum of two-dimensional representations (over  $\mathbb{C}$ ) of  $\mathrm{SL}(2, \mathbb{C})$ , corresponding to “left-handed” and “right-handed” fermions, in the sense of 21.

### 3.1 Interpretations of $V_{m,n}$ and (ToE2)

In the notation of (1.1), the  $V_{m,n}$ , with  $m+n$  odd, correspond to fermions; those with  $m+n$  even correspond to bosons. In Lisi’s setup, the bosons are 1-forms on  $M$ , with values in a vector bundle associated to the aforementioned  $\mathrm{Spin}(3, 1)_0$  principal bundle via the  $m \otimes n$  representation (with  $m+n$  even). The  $V_{m,n}$  with  $m+n=4$  are special; they correspond to the gravitational degrees of freedom in Lisi’s theory.  $(3 \otimes 1) \oplus (1 \otimes 3)$  is the adjoint representation of  $\mathrm{SL}(2, \mathbb{C})$ ; these correspond to the spin connection. The 1-form with values in the  $2 \otimes 2$  representation is the vierbein<sup>1</sup>.

It is a substantial result from physics (see Sects. 13.1, 25.4 of (9)) that a unitary interacting theory is incompatible with massless particles in higher representations ( $m+n \geq 6$ ). Our hypothesis (ToE2) reflects this and also forbids gravitinos ( $m+n=5$ ). In 10, we will revisit the possibility of admitting gravitinos.

### 3.2 Explanation of (ToE3)

Our hypothesis (ToE3) says that the candidate “Theory of Everything” one obtains from subgroups  $\mathrm{SL}(2, \mathbb{C})$  and  $G$  as in (ToE1) must be *chiral* in the sense of Definition 24.<sup>2</sup>

<sup>1</sup> In making this identification, we have tacitly assumed that  $V_{2,2}$  is one-dimensional. This is, in fact, required for a unitary interacting theory. We will not, however, impose this additional constraint. Suffice to say that it is not satisfied by any of the *candidate ToE subgroups* (per Definition 11) of  $E_8$ .

<sup>2</sup> Of course, there are many other features of the Standard Model that a candidate Theory of Everything must reproduce. We have chosen to focus on the requirement that the theory be chiral for two reasons. First, it is “physically robust”: Whatever intricacies a quantum field theory may possess at high energies, if it is *non-chiral*, there is no known mechanism by which it could reduce to a *chiral* theory at low energies (and there are strong arguments (10) that no such mechanism exists). Second, chirality is easily translated into a mathematical criterion—

In private communication, Lisi has indicated that he objects to our condition (ToE3), because he no longer wishes to identify all 248 generators of  $\text{Lie}(E)$  as particles (either bosons or fermions). In his new—and unpublished—formulation, only a subset are to be identified as particles. In particular,  $V_{2,1}$  is typically a reducible representation of  $G$  and, in his new formulation, only a subrepresentation corresponds to particles (fermions). This is not the approach followed in (1), where all 248 generators are identified as particles and where, moreover, 20-odd of these are claimed to be new as-yet undiscovered particles—a prediction of his theory. As recently as April 2009, Lisi reiterated this prediction in an essay published in the *Financial Times*, (11).

Our paper assumes that the approach of (1) is to be followed, and that all 248 generators are to be identified as particles, hence (ToE3). In any case, even if one identifies only a subset of the generators as particles, all the fermions must come from the  $(-1)$ -eigenspace, which is too small to accommodate 3 generations, as we now show.

### 3.3 No-go based on dimensions

The fermions of Lisi’s theory correspond to weight vectors in  $V_{m,n}$ , with  $m+n$  odd. In particular, the weight vectors in  $V_{2,1}$  and  $V_{1,2}$  correspond (as in 21) to left- and right-handed fermions, respectively. Since there are  $3 \times 15 = 45$  known fermions of each chirality,  $V_{2,1}$  must be at least 45-dimensional, and similarly for  $V_{1,2}$ . Thus, the  $-1$ -eigenspace of the central element of  $\text{SL}(2, \mathbb{C})$ , which contains  $(2 \otimes 1 \otimes V_{2,1}) \oplus (1 \otimes 2 \otimes V_{1,2})$ , must have dimension at least  $2 \times 2 \times 45 = 180$ .

When  $E$  is a real form of  $E_8$ , the  $-1$ -eigenspace has dimension 112 or 128 (this is implicit in Elie Cartan’s classification of real forms of  $E_8$  as in (12, p. 518, Table V)),<sup>3</sup> so no identification of the fermions as distinct weight vectors in  $\text{Lie}(E)$  (as in Table 9 in (1)) can be compatible with the Spin-Statistics Theorem and the existence of three generations.

These dimensional considerations do not, however, rule out the possibility of accommodating a 1- or 2-generation Standard Model (per Definition 25) in a real form of  $E_8$ . That requires more powerful considerations, which are the subject of our main theorem. We now turn to the proof of that theorem.

## 4 $\mathfrak{sl}_2$ Subalgebras and the Dynkin Index

### The Dynkin index

In (15, 2), Dynkin defined the *index* of an inclusion  $f: \mathfrak{g}_1 \hookrightarrow \mathfrak{g}_2$  of simple complex Lie algebras as follows. Fix a Chevalley basis of the two algebras, so that the Cartan subalgebra  $\mathfrak{h}_1$  of  $\mathfrak{g}_1$  is contained in the Cartan subalgebra  $\mathfrak{h}_2$  of  $\mathfrak{g}_2$ . The

---

our (ToE3). This allows us to study a purely representation-theoretic question and side-step the difficulties of making sense of Lisi’s proposal as a dynamical quantum field theory.

<sup>3</sup> Alternatively, Serre’s marvelous bound on the trace from (13, Th. 3) or (14, Th. 1) implies that for every element  $x$  of order 2 in a reductive complex Lie group  $G$ , the  $-1$ -eigenspace of  $\text{Ad}(x)$  has dimension  $\leq (\dim G + \text{rank } G)/2$ . In particular, when  $G$  is a real form of  $E_8$ , the  $-1$ -eigenspace has dimension  $\leq 128$ .

Chevalley basis identifies  $\mathfrak{h}_i$  with the complexification  $Q_i^\vee \otimes \mathbb{C}$  of the coroot lattice  $Q_i^\vee$  of  $\mathfrak{g}_i$ , and the inclusion  $f$  gives an inclusion  $Q_1^\vee \otimes \mathbb{C} \hookrightarrow Q_2^\vee \otimes \mathbb{C}$ . Fix the Weyl-invariant inner product  $(\cdot, \cdot)_i$  on  $Q_i^\vee$  so that  $(\alpha^\vee, \alpha^\vee)_i = 2$  for short coroots  $\alpha^\vee$ . Then the *Dynkin index* of the inclusion is the ratio  $(f(\alpha^\vee), f(\alpha^\vee))_2 / (\alpha^\vee, \alpha^\vee)_1$  where  $\alpha^\vee$  is a short coroot of  $\mathfrak{g}_1$ . For example, the irreducible representation  $\mathfrak{sl}_2 \rightarrow \mathfrak{sl}_n$  has index  $\binom{n+1}{3}$  by (15, Eq. (2.32)).

### $\mathfrak{sl}_2$ subalgebras

We now consider the case  $\mathfrak{g}_1 = \mathfrak{sl}_2$  and write simply  $\mathfrak{g}$  and  $Q^\vee$  for  $\mathfrak{g}_2$  and  $Q_2^\vee$ . The coroot lattice of  $\mathfrak{sl}_2$  is  $\mathbb{Z}$  and the image of 1 under the map  $\mathbb{Z} \hookrightarrow Q^\vee$  is an element  $h \in \mathfrak{h}$  called the *defining vector* of the inclusion. In 8 of his paper (or see (16, VIII.11)), Dynkin proved that, after conjugating by an element of the automorphism group of  $\mathfrak{g}$ , one can assume that the defining vector  $h$  satisfies the strong restrictions:

$$h = \sum_{\delta \in \Delta} p_\delta \delta^\vee \quad \text{for } p_\delta \text{ real and non-negative [15, Lemma 8.3],}$$

where  $\Delta$  denotes the set of simple roots of  $\mathfrak{g}$  and further that

$$\delta(h) \in \{0, 1, 2\} \quad \text{for all } \delta \in \Delta. \quad (4.1)$$

But note that for each simple root  $\delta$ , the fundamental irreducible representation of  $\mathfrak{g}$  with highest weight dual to  $\delta^\vee$  restricts to a representation of  $\mathfrak{sl}_2$  with  $p_\delta$  as a weight, hence  $p_\delta$  is an integer.

As a consequence of these generalities and specifically (15, Lemma 8.2), one can identify an  $\mathfrak{sl}_2$  subalgebra of  $\mathfrak{g}$  up to conjugacy by writing the Dynkin diagram of  $\mathfrak{g}$  and putting the number  $\delta(h)$  from Display (4.1) at each vertex; this is the *marked Dynkin diagram* of the  $\mathfrak{sl}_2$  subalgebra.

Here is an alternative formula for computing the index of an  $\mathfrak{sl}_2$  subalgebra from its marked Dynkin diagram. Write  $\kappa_{\mathfrak{g}}$  and  $m^\vee$  for the Killing form and dual Coxeter number of  $\mathfrak{g}$ . We have:

$$(\text{Dynkin index}) = \frac{1}{2}(h, h) = \frac{1}{4m^\vee} \kappa_{\mathfrak{g}}(h, h) = \frac{1}{2m^\vee} \sum_{\text{positive roots } \alpha \text{ of } \mathfrak{g}} \alpha(h)^2, \quad (4.2)$$

where the second equality is by, e.g., (17, 5), and the third is by the definition of  $\kappa_{\mathfrak{g}}$ . One can calculate the number  $\alpha(h)$  by writing  $\alpha$  as a sum of positive roots and applying the marked Dynkin diagram for  $h$ .

**Lemma 41** *For every simple complex Lie algebra  $\mathfrak{g}$ , there is a unique copy of  $\mathfrak{sl}_2$  in  $\mathfrak{g}$  of index 1, up to conjugacy.*

This is (equivalent to) Theorem 2.4 in (15). We give a different proof for the convenience of the reader.

*Proof* The index of an  $\mathfrak{sl}_2$ -subalgebra is  $(h, h)/2$ , where the defining vector  $h$  belongs to the coroot lattice  $Q^\vee$ . If  $\mathfrak{g}$  is not of type B, then the coroot lattice is not of type C, and the claim amounts to the statement that the vectors of minimal length in the coroot lattice are actually coroots. This follows from the constructions of the root lattices in (18, 12.1).

Otherwise  $\mathfrak{g}$  has type B and is  $\mathfrak{so}_n$  for some odd  $n \geq 5$ . The conjugacy class of an  $\mathfrak{sl}_2$ -subalgebra is determined by the restriction of the natural  $n$ -dimensional representation; they are parameterized by partitions of  $n$  (i.e.,  $\sum n_i = n$ ) so that the even  $n_i$  occur with even multiplicity and some  $n_i > 1$ , see (19, 5.1.2) or (20, 6.2.2). The index of the composition  $\mathfrak{sl}_2 \rightarrow \mathfrak{so}_n \rightarrow \mathfrak{sl}_n$  is then  $\sum \binom{n_i+1}{3}$ ; we must classify those partitions such that this sum equals the Dynkin index of  $\mathfrak{so}_n \rightarrow \mathfrak{sl}_n$ , which is 2. The unique such partition is  $2 + 2 + 1 + \cdots + 1 > 0$ .  $\square$

In the bijection between conjugacy classes of  $\mathfrak{sl}_2$  subalgebras and orbits of nilpotent elements in  $\mathfrak{g}$  from (19, 3.2.10), the unique orbit of index 1  $\mathfrak{sl}_2$ ’s corresponds to the minimal nilpotent orbit described in (19, 4.3.3).

If  $\mathfrak{g}$  has type C,  $F_4$ , or  $G_2$ , then the argument in the proof of the lemma shows that there is up to conjugacy a unique copy of  $\mathfrak{sl}_2$  in  $\mathfrak{g}$  with index 2, 2, or 3 respectively. For  $\mathfrak{g}$  of type  $B_n$  with  $n \geq 4$ , there are two conjugacy classes of  $\mathfrak{sl}_2$ -subalgebras of index 2. This amounts to the fact that there are vectors in the  $C_n$  root lattice that are not roots but have the same length as a root—specifically, sums of two strongly orthogonal short roots, cf. Exercise 5 in 12 of (18).

## 5 Copies of $\mathfrak{sl}_{2,\mathbb{C}}$ in the Complex $E_8$

We now prove some facts about copies of  $\mathfrak{sl}_{2,\mathbb{C}}$  in the complex Lie algebra  $\mathfrak{e}_8$  of type  $E_8$ . Of course, the 69 conjugacy classes of such are known—see (15, pp. 182–185) or (21, pp. 430–433)—but we do not need this information.

Fix a pinning for  $\mathfrak{e}_8$ ; this includes a Cartan subalgebra  $\mathfrak{h}$ , a set of simple roots  $\Delta := \{\alpha_i \mid 1 \leq i \leq 8\}$  (numbered

$$\begin{array}{cccccccc} 1 & 3 & 4 & 5 & 6 & 7 & 8 \\ & & 2 & & & & \end{array} \quad (5.1)$$

as in (22)), and fundamental weights  $\omega_i$  dual to  $\alpha_i$ . As all roots of the  $E_8$  root system have the same length, we can and do identify the root system with its coroot system (also called the “inverse” or “dual” root system).

*Example 51* Taking any root of  $E_8$ , one can generate a copy of  $\mathfrak{sl}_{2,\mathbb{C}}$  in  $\mathfrak{e}_8$  with index 1. Doing this with the highest root gives an  $\mathfrak{sl}_{2,\mathbb{C}}$  with marked Dynkin diagram

$$\text{index 1:} \quad \begin{array}{cccccccc} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ & & 0 & & & & \end{array}.$$

Every index 1 copy of  $\mathfrak{sl}_2$  in  $\mathfrak{e}_8$  is conjugate to this one by Lemma 41.

*Example 52* One can find a copy of  $\mathfrak{sl}_{2,\mathbb{C}} \times \mathfrak{sl}_{2,\mathbb{C}}$  in  $\mathfrak{e}_8$  by taking the first copy to be generated by the highest root of  $E_8$  and the second copy to be generated by the

highest root of the obvious  $E_7$  subsystem. If you embed  $\mathfrak{sl}_{2,\mathbb{C}}$  diagonally in this algebra, you find a copy of  $\mathfrak{sl}_{2,\mathbb{C}}$  with index 2 and marked Dynkin diagram

$$\text{index 2: } \begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & 0 & & & & \end{array} .$$

**Proposition 53** *The following collections of copies of  $\mathfrak{sl}_{2,\mathbb{C}}$  in  $\mathfrak{e}_8$  are the same:*

- (1) *copies such that  $\pm 1$  are weights of  $\mathfrak{e}_8$  (as a representation of  $\mathfrak{sl}_{2,\mathbb{C}}$ ) and no other odd weights occur,*
- (2) *copies such that every weight of  $\mathfrak{e}_8$  is in  $\{0, \pm 1, \pm 2\}$ ,*
- (3) *copies such that the inclusion  $\mathfrak{sl}_{2,\mathbb{C}} \subset \mathfrak{e}_8$  has Dynkin index 1 or 2,*
- (4) *copies of  $\mathfrak{sl}_{2,\mathbb{C}}$  conjugate to one of those defined in Examples 51 or 52.*

*Proof* One easily checks that (4) is contained in (1)–(3); we prove the opposite inclusion.

For (3), we identify  $\mathfrak{h}$  with the complexification  $Q \otimes \mathbb{C}$  of the (co)root lattice  $Q$ , hence  $h$  with  $\sum \alpha_i(h) \omega_i$ . By Eq. (4.2), the index of  $h$  satisfies:

$$\frac{1}{60} \sum_{\alpha} \alpha(h)^2 = \frac{1}{60} \sum_{\alpha} \left( \sum_i \alpha_i(h) \langle \omega_i, \alpha \rangle \right)^2 \geq \sum_i \left( \alpha_i(h)^2 \sum_{\alpha} \frac{\langle \omega_i, \alpha \rangle^2}{60} \right),$$

where the sums vary over the positive roots. We calculate for each fundamental weight  $\omega_i$  the number  $\sum_{\alpha} \langle \omega_i, \alpha \rangle^2 / 60$ :

$$\begin{array}{ccccccc} 2 & 7 & 15 & 10 & 6 & 3 & 1 \\ & & 4 & & & & \end{array} . \quad (5.2)$$

As the numbers  $\alpha_i(h)$  are all 0, 1, or 2, the numbers in Display (5.2) show that  $h$  for an  $\mathfrak{sl}_{2,\mathbb{C}}$  with Dynkin index 1 or 2 must be  $\omega_1$  (index 2) or  $\omega_8$  (index 1).

For (2), the highest root  $\tilde{\alpha}$  of  $E_8$  is  $\tilde{\alpha} = \sum_i c_i \alpha_i$ , where  $c_1 = c_8 = 2$  and the other  $c_i$ 's are all at least 3. As  $\tilde{\alpha}(h)$  is a weight of  $\mathfrak{e}_8$  relative to a given copy of  $\mathfrak{sl}_{2,\mathbb{C}}$ , we deduce that an  $\mathfrak{sl}_{2,\mathbb{C}}$  as in (2) must have  $h = \omega_1$  or  $\omega_8$ , as claimed.

Suppose now that we are given an  $h$  for a copy of  $\mathfrak{sl}_{2,\mathbb{C}}$  as in (1). As  $\pm 1$  occur as weights, there is at least one 1 in the marked Dynkin diagram.

But note that there cannot be three or more 1's in the marked Dynkin diagram for  $h$ . Indeed, for every connected subset  $S$  of vertices of the Dynkin diagram of  $E_8$ ,  $\sum_{i \in S} \alpha_i$  is a root (22, VI.1.6, Cor. 3b). If the number of 1's in the marked diagram of  $h$  is at least three, then one can pick  $S$  so that it meets exactly three of the  $\alpha_i$ 's with  $\alpha_i(h) = 1$ , in which case  $\sum_{i \in S} \alpha_i(h)$  is odd and at least 3, violating the hypothesis of (1).

For the sake of contradiction, suppose that there are two 1's in the marked diagram for  $h$ , say, corresponding to simple roots  $\alpha_i$  and  $\alpha_j$  with  $i < j$ . For each  $i, j$ , one can find a root  $\beta$  in the list of roots of  $E_8$  of large height in (22, Plate VII) such that the coefficients of  $\alpha_i$  and  $\alpha_j$  in  $\beta$  have opposite parity and sum at least 3. (Merely taking  $\beta$  to be the highest root suffices for many  $(i, j)$ .) This contradicts (1), so there is a unique 1 in the marked diagram for  $h$ , i.e.,  $\alpha_i(h) = 1$  for a unique  $i$ .

If  $\alpha_i(h) = 1$  for some  $i \neq 1, 8$ , then we find a contradiction because there is a root  $\alpha$  of  $E_8$  with  $\alpha_i$ -coordinate 3. Therefore  $\alpha_i(h) = 1$  only for  $i = 1$  or 8 and not for both. By the fact used two paragraphs above,  $\beta := \sum_i \alpha_i$  is a root of  $E_8$ , so  $\beta(h) = \sum \alpha_i(h)$  is odd and must be 1. It follows that  $h = \omega_1$  or  $\omega_8$ .  $\square$

*Observation 54 (Centralizer for index 1)* The  $\mathfrak{sl}_{2,\mathbb{C}}$  of index 1 in  $\mathfrak{e}_8$  has centralizer the obvious regular subalgebra  $\mathfrak{e}_7$  of type  $E_7$ . (A subalgebra is *regular* if it is generated by the root subalgebras corresponding to a closed sub-root-system (15, no. 16).) Indeed, it is clear that  $\mathfrak{e}_7$  centralizes this  $\mathfrak{sl}_{2,\mathbb{C}}$ . Conversely, the centralizer of  $\mathfrak{sl}_{2,\mathbb{C}}$  is contained in the centralizer of  $h = \omega_8$ —i.e.,  $\mathfrak{e}_7 \oplus \mathbb{C}h$ —but does not contain  $h$ .

*Technique 55 (Decomposing  $\mathfrak{e}_8$ )* Suppose we are given a copy of  $\mathfrak{sl}_{2,\mathbb{C}}$  in  $\mathfrak{e}_8$  specified by a defining vector  $h$ . By applying the 240 roots of  $\mathfrak{e}_8$  to  $h$  (and throwing in also 0 with multiplicity 8), we obtain the weights of  $\mathfrak{e}_8$  as a representation of  $\mathfrak{sl}_{2,\mathbb{C}}$  and therefore also the decomposition of  $\mathfrak{e}_8$  into irreducible representations of  $\mathfrak{sl}_{2,\mathbb{C}}$  as in, e.g., (18, 7.2).

Extending this, suppose we are given a copy of  $\mathfrak{sl}_{2,\mathbb{C}} \times \mathfrak{sl}_{2,\mathbb{C}}$  in  $\mathfrak{e}_8$ , where the two summands are specified by defining vectors in  $\mathfrak{h}$ . (Here we want the defining vectors to span the Cartan subalgebras in the images of the two  $\mathfrak{sl}_{2,\mathbb{C}}$ ’s. In particular, they need not be normalized in the sense of Display (4.1).) Computing as in the previous paragraph, we can decompose  $\mathfrak{e}_8$  as a direct sum of irreducible representations  $m \otimes n$  of  $\mathfrak{sl}_{2,\mathbb{C}} \times \mathfrak{sl}_{2,\mathbb{C}}$ . It is easy to write code from scratch to make a computer algebra system perform this computation. We remark that applying this recipe in the situation from the Introduction gives the dimension of  $V_{m,n}$  as the multiplicity of  $m \otimes n$ .

## 6 Index 2 Copies of $\mathfrak{sl}_{2,\mathbb{C}}$ in the Complex $E_8$

**Lemma 61** *The centralizer of the index 2  $\mathfrak{sl}_{2,\mathbb{C}}$  in  $\mathfrak{e}_8$  from Example 52 is a copy of  $\mathfrak{so}_{13}$  contained in the regular subalgebra  $\mathfrak{so}_{14}$  of  $\mathfrak{e}_8$ .*

*Proof* The centralizer of the  $\mathfrak{sl}_{2,\mathbb{C}}$  of index 2 in  $\mathfrak{e}_8$  is contained in the centralizer of the defining vector  $h$ ; this centralizer is reductive with semisimple part the regular subalgebra  $\mathfrak{so}_{14}$  of type  $D_7$ . The centralizer of  $\mathfrak{sl}_{2,\mathbb{C}}$  contains the centralizer of the  $\mathfrak{sl}_{2,\mathbb{C}} \times \mathfrak{sl}_{2,\mathbb{C}}$  from Example 52, which is the regular subalgebra  $\mathfrak{so}_{12}$  of type  $D_6$ , as can be seen by the recipe from (15, pp. 147, 148). Computing as in 55, we see that the centralizer of  $\mathfrak{sl}_{2,\mathbb{C}}$  has dimension 78 (as is implicitly claimed in the statement of the lemma), so it lies properly between the regular  $\mathfrak{so}_{12}$  and the regular  $\mathfrak{so}_{14}$ .

For concreteness, let us suppose that the structure constants for  $\mathfrak{e}_8$  are as in (23). Define a copy of  $\mathfrak{sl}_{2,\mathbb{C}}$  by sending  $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$  to the sum of the elements in the Chevalley basis of  $\mathfrak{e}_8$  spanning the root subalgebras corresponding to  $-\alpha_8$  and the highest root in the obvious  $D_7$  subdiagram. This copy of  $\mathfrak{sl}_{2,\mathbb{C}}$  has defining vector  $\alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + 2\alpha_7$ . One checks using the structure constants that this  $\mathfrak{sl}_{2,\mathbb{C}}$  centralizes the index 2  $\mathfrak{sl}_{2,\mathbb{C}}$  we started with, and that together with  $\mathfrak{so}_{12}$  it generates a copy of  $\mathfrak{so}_{13}$ . In particular, the coroot lattice of this  $\mathfrak{so}_{13}$  has basis  $\beta_1^\vee, \dots, \beta_6^\vee$ , embedded in the (co)root lattice of  $\mathfrak{e}_8$  as in the table:

$$\begin{array}{c|cccccc} \mathfrak{so}_{13} & \beta_1^\vee & \beta_2^\vee & \beta_3^\vee & \beta_4^\vee & \beta_5^\vee & \beta_6^\vee \\ \mathfrak{e}_8 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & -\alpha_2 - \alpha_3 - 2\alpha_4 - 2\alpha_5 - 2\alpha_6 - 2\alpha_7 \end{array}. \quad (6.1)$$

We remark that the numbering of the coroots  $\beta_1^\vee, \dots, \beta_6^\vee$  corresponds to a numbering of the simple roots of  $\mathfrak{so}_{13}$  as in the diagram

Dimension count shows that this  $\mathfrak{so}_{13}$  is the centralizer.  $\square$

The claim of the lemma is already in (24, p. 125). We gave the details of a proof because it specifies an inclusion of  $\mathfrak{so}_{13}$  in  $\mathfrak{e}_8$  and a comparison of the pinnings of the two algebras as in Table (6.1).

The index 2  $\mathfrak{sl}_2$  and the copy of  $\mathfrak{so}_{13}$  give an  $\mathfrak{sl}_2 \times \mathfrak{so}_{13}$  subalgebra of  $\mathfrak{e}_8$ . We now decompose  $\mathfrak{e}_8$  into irreducible representations of  $\mathfrak{sl}_2 \times \mathfrak{so}_{13}$ . We can do this from first principles by restricting the roots of  $\mathfrak{e}_8$  to the Cartan subalgebras of  $\mathfrak{sl}_2$  (using the marked Dynkin diagram from Example 52) and  $\mathfrak{so}_{13}$  (using Table (6.1)). Alternatively, we can read the decomposition off the tables in (25) as follows. As in the proof of Lemma 61,  $\mathfrak{sl}_2$  is contained in the regular subalgebra  $\mathfrak{sl}_2 \times \mathfrak{sl}_2 \times \mathfrak{so}_{12}$  of  $\mathfrak{e}_8$ , and the tables on pages 301 and 305 of *ibid.* show that  $\mathfrak{e}_8$  decomposes as a sum of

$$\text{the adjoint representation, } 2 \otimes 1 \otimes S_+, \quad 1 \otimes 2 \otimes S_-, \quad \text{and } 2 \otimes 2 \otimes V, \quad (6.2)$$

where  $S_{\pm}$  denotes the half-spin representations of  $\mathfrak{so}_{12}$  and  $V$  is the vector representation. We can restrict the representations of  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$  to the diagonal  $\mathfrak{sl}_2$  subalgebra to obtain a decomposition of  $\mathfrak{e}_8$  into representations of  $\mathfrak{sl}_2 \times \mathfrak{so}_{12}$ . Consulting the tables in *ibid.* for restricting representations from type  $B_6$  to  $D_6$  allows us to deduce the decomposition

$$1 \otimes \mathfrak{so}_{13, \mathbb{C}} \oplus 2 \otimes (\text{spin}) \oplus 3 \otimes 1 \oplus 3 \otimes (\text{vector}) \quad (6.3)$$

of  $\mathfrak{e}_8$  as a representation of  $\mathfrak{sl}_2 \times \mathfrak{so}_{13}$ . From this it is obvious that  $\mathfrak{so}_{13, \mathbb{C}}$  is the Lie algebra of a copy of  $\text{Spin}_{13}$  in  $E_8$ .

The main result of this section is the following:

**Lemma 62** *Up to conjugacy, there is a unique copy of  $\text{SL}_{2, \mathbb{C}} \times \text{SL}_{2, \mathbb{C}}$  in  $E_{8, \mathbb{C}}$  so that each inclusion of  $\text{SL}_{2, \mathbb{C}}$  in  $E_{8, \mathbb{C}}$  has index 2. The centralizer of this  $\text{SL}_{2, \mathbb{C}} \times \text{SL}_{2, \mathbb{C}}$  has identity component  $\text{Sp}_{4, \mathbb{C}} \times \text{Sp}_{4, \mathbb{C}}$ .*

*Proof* As in the proof of Lemma 41 (or by the method used to prove Prop. 53), there are two index 2 copies of  $\mathfrak{sl}_2$  in  $\mathfrak{so}_{13}$ , corresponding to the partitions

$$(a) \quad 3 + 1 + 1 + \cdots + 1 \quad \text{and} \quad (b) \quad 2 + 2 + 2 + 2 + 1 + 1 + \cdots + 1$$

of 13. The recipe in (19, 5.3) gives defining vectors for these  $\mathfrak{sl}_2$ 's, which we can rewrite in terms of the  $E_8$  simple roots using Table (6.1):

$$\begin{aligned} (a) \quad & 2\beta_1^\vee + 2\beta_2^\vee + 2\beta_3^\vee + 2\beta_4^\vee + 2\beta_5^\vee + \beta_6^\vee = -\alpha_2 + \alpha_3, \\ (b) \quad & \beta_1^\vee + 2\beta_2^\vee + 3\beta_3^\vee + 4\beta_4^\vee + 4\beta_5^\vee + 2\beta_6^\vee = -2\alpha_2 - \alpha_3 - 2\alpha_4 - \alpha_5. \end{aligned} \quad (6.4)$$

We can pair each of (a) and (b) with the copy of  $\mathfrak{sl}_2$  from Example 52 to get an  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$  subalgebra of  $\mathfrak{e}_8$  where both  $\mathfrak{sl}_2$ 's have index 2. Clearly, these represent the only two  $E_8$ -conjugacy classes of such subalgebras. With Display (6.4) in hand, we can calculate the multiplicities of the irreducible representations of  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$  in  $\mathfrak{e}_8$  as in 55.

In case (a), every irreducible summand  $m \otimes n$  has  $m + n$  even. Therefore, this copy of  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$  is the Lie algebra of a subgroup of  $E_8$  isomorphic to  $(\text{SL}_2 \times \text{SL}_2)/(-1, -1)$ .

(An alternative way to see this is to note that the simple roots with odd coefficients are the same in Display (6.4a) and the defining vector in Example 52.)

In case (b), we have the following table of multiplicities for  $m \otimes n$ :

$$\begin{array}{c|ccc} & 1 & 2 & 3 & m \\ \hline 1 & 20 & 20 & 6 & \\ n & 2 & 20 & 16 & 4 \\ & 3 & 6 & 4 & 0 \end{array} \quad (6.5)$$

In particular, it is the Lie algebra of a copy of  $SL_2 \times SL_2$  in  $E_8$ . The centralizer of (b) in  $Spin_{13}$  has been calculated in (26, IV.2.25), and the identity component is  $Sp_4 \times Sp_4$ , as claimed.  $\square$

We can decompose  $\mathfrak{e}_8$  into a direct sum of irreducible representations of the  $\mathfrak{sl}_2 \times \mathfrak{sl}_2 \times \mathfrak{sp}_4 \times \mathfrak{sp}_4$  subalgebra from Lemma 62 by combining the decomposition of  $\mathfrak{e}_8$  into irreducible representations of  $\mathfrak{sl}_2 \times \mathfrak{so}_{13}$  from Decomposition (6.3) with the tables in (25). Specifically, we restrict representations from  $\mathfrak{so}_{13}$  to an  $\mathfrak{sp}_4 \times \mathfrak{so}_8$  subalgebra and then from  $\mathfrak{so}_8$  to  $\mathfrak{sp}_4 \times \mathfrak{sl}_2$ , where this  $\mathfrak{sl}_2$  also has index 2. Recall that  $\mathfrak{sp}_4$  has two fundamental irreducible representations: one that is 4-dimensional symplectic and another that is 5-dimensional orthogonal; we denote them by their dimensions. With this notation and 1.1, we find:

$$V_{2,1} \simeq 5 \otimes 4, \quad V_{1,2} \simeq 4 \otimes 5, \quad V_{2,3} \simeq 1 \otimes 4, \quad V_{3,2} \simeq 4 \otimes 1, \quad \text{and} \quad V_{2,2} \simeq 4 \otimes 4. \quad (6.6)$$

## 7 Copies of $SL(2, \mathbb{C})$ in a Real Form of $E_8$

Suppose now that we have a copy of  $SL(2, \mathbb{C})$  inside a real Lie group  $E$  of type  $E_8$ . Over the complex numbers, we decompose  $\text{Lie}(E) \otimes \mathbb{C}$  into a direct sum of irreducible representations of  $SL(2, \mathbb{C}) \otimes \mathbb{C} \simeq SL_{2, \mathbb{C}} \times SL_{2, \mathbb{C}}$ ; each irreducible representation can be written as  $m \otimes n$  where  $m$  and  $n$  denote the dimension of an irreducible representation of the first or second  $SL_{2, \mathbb{C}}$  respectively. The goal of this section is to prove:

**Proposition 71** *Maintain the notation of the previous paragraph. If  $\text{Lie}(E) \otimes \mathbb{C}$  contains no irreducible summands  $m \otimes n$  with  $m + n > 4$ , then the identity component  $Z$  of the centralizer of  $SL(2, \mathbb{C})$  in  $E$  is a subgroup isomorphic to*

- (1)  $Spin(7, 5)$  if  $E$  is split; or
- (2)  $Spin(3, 9)$  or  $Spin(11, 1)$  if the Killing form of  $\text{Lie}(E)$  has signature  $-24$ .

*In either case,  $\text{Lie}(Z) \otimes \mathbb{C}$  is the regular  $\mathfrak{so}_{12}$  subalgebra of  $\text{Lie}(E) \otimes \mathbb{C}$ .*

*Proof* Complexifying the inclusion of  $SL(2, \mathbb{C})$  in  $E$  and going to Lie algebras gives an inclusion of  $\mathfrak{sl}_{2, \mathbb{C}} \times \mathfrak{sl}_{2, \mathbb{C}}$  in the complex Lie algebra  $\mathfrak{e}_8$ . The hypothesis on the irreducible summands  $m \otimes n$  implies that each of the two  $\mathfrak{sl}_{2, \mathbb{C}}$ 's has index 1 or 2 by Proposition 53. As complex conjugation interchanges the two components, they must have the same index.

Suppose first that both  $\mathfrak{sl}_2$ 's have index 2. When we decompose  $\mathfrak{e}_8$  as in 1.1, we find the representation  $2 \otimes 3$  with positive multiplicity 4 by Table (6.5), which violates our hypothesis on the  $SL(2, \mathbb{C})$  subgroup of  $E$ .

Therefore both  $\mathfrak{sl}_2$ 's have index 1. Lemma 41 (twice) gives that this  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$  is conjugate to the one generated by the highest root of  $E_8$  from Example 51 (so the second  $\mathfrak{sl}_2$  belongs to the centralizer of type  $E_7$ ) and by the highest root of the  $E_7$  subsystem and makes up the first two summands of an  $\mathfrak{sl}_2 \times \mathfrak{sl}_2 \times \mathfrak{so}_{12}$  subalgebra, the same one used to find decomposition (6.2). That is,  $\mathfrak{so}_{12}$  centralizes  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$ . Conversely, the centralizer of the defining vectors of the two copies of  $\mathfrak{sl}_2$  has semisimple part  $\mathfrak{so}_{12}$ ; it follows that  $\text{Lie}(Z) \otimes \mathbb{C}$  is isomorphic to  $\mathfrak{so}_{12}$ .

From this and decomposition (6.2), we see that  $Z$  is a real form of  $\text{Spin}_{12}$ . As  $\text{Lie}(E)$  is a real representation of  $Z$ , we deduce that  $V$  is also a real representation of  $Z$  but  $S_+$  and  $S_-$  are not; they are interchanged by the Galois action. The first observation shows that  $Z$  is  $\text{Spin}(12-a, a)$  for some  $0 \leq a \leq 6$ . The second shows that  $a$  must be 1, 3, or 5, as claimed in the statement of the proposition.

It remains to prove the correspondence between  $a$  and the real forms of  $E_8$ . For  $a = 5$ , this is clear: the subgroup generated by  $\text{SL}(2, \mathbb{C})$  and  $\text{Spin}(7, 5)$  has real rank 6, so it can only be contained in the split real form.

Now suppose that  $a = 3$  or 1 and that  $\text{SL}(2, \mathbb{C})$  is in the split  $E_8$ ; we will show that the Killing form of  $E$  has signature  $-24$ . Over  $\mathbb{C}$ ,  $\text{SL}(2, \mathbb{C})$  is conjugate to the copy of  $\text{SL}_{2, \mathbb{C}} \times \text{SL}_{2, \mathbb{C}}$  in  $E_{8, \mathbb{C}}$  generated by the highest root of  $E_8$  and the highest root of the natural subsystem of type  $E_7$ . Writing out these two roots in terms of the  $E_8$  simple roots, we see that  $\alpha_3$  and  $\alpha_5$  are the only simple roots whose coefficients have different parities. It follows that the element  $-1 \in \text{SL}(2, \mathbb{C})$ —equivalently,  $(-1, -1) \in \text{SL}_2 \times \text{SL}_2$ —is  $h_{\alpha_2}(-1)h_{\alpha_3}(-1)$  in the notation of (27), where  $h_{\alpha_i} : \mathbb{C}^\times \rightarrow E \otimes \mathbb{C}$  is the cocharacter corresponding to the coroot  $\alpha_i^\vee$ . Now,  $\alpha_2$  and  $\alpha_3$  are the only simple roots with odd coefficients in the fundamental weight  $\omega_1$ , so the subgroup of  $E \otimes \mathbb{C}$  fixed by conjugation by this  $-1$  is generated by root subgroups corresponding to roots  $\alpha$  such that  $\langle \omega_1, \alpha \rangle$  is even. These roots form the natural  $D_8$  subsystem of  $E_8$ , and in this way we see  $\text{SL}(2, \mathbb{C}) \cdot \text{Spin}(12-a, a)$  as a semisimple subgroup of maximal rank in a copy of a half-spin group  $H$  in 16 dimensions—the identity component of the centralizer of  $-1$ .

We claim that  $H$  is isogenous to  $\text{SO}(12, 4)$ . As  $H$  is a half-spin group with a half-spin representation defined over  $\mathbb{R}$ , it is isogenous to  $\text{SO}(16-b, b)$  for  $b = 0, 4$ , or 8 or it is quaternionic; these possibilities have Killing forms of signature  $-120, -24, 8$ , or  $-8$  respectively, as can be looked up in (28), for example. The adjoint representation of  $H$ , when restricted to  $\text{SL}(2, \mathbb{C}) \cdot \text{Spin}(12-a, a)$ , decomposes as the adjoint representation of  $\text{SL}(2, \mathbb{C}) \cdot \text{Spin}(12-a, a)$  and  $2 \otimes 2 \otimes V$  by decomposition (6.2). The Killing form on  $H$  restricts to a positive multiple of the Killing form on  $\text{SL}(2, \mathbb{C}) \cdot \text{Spin}(12-a, a)$  (as can be seen over  $\mathbb{C}$  by the explicit formula on p. E-14 of (26))—i.e., has signature  $-44$  or  $-12$  for  $a = 1$  or 3—and a form of signature  $\pm 2(12-2a)$  on  $2 \otimes 2 \otimes V$ ; the sum of these has signature 0,  $-24$ , or  $-64$  since  $a = 1$  or 3. Comparing the two lists verifies that  $H$  is isogenous to  $\text{SO}(12, 4)$ .

The Killing form on  $H$  has signature  $-24$ . The invariant bilinear form on the half-spin representation is hyperbolic (because  $H$  is isogenous to spin of an isotropic quadratic form of dimension divisible by 8, see (29, 1.1)). As a representation of  $H$ ,  $\text{Lie}(E)$  is a sum of these two representations, and we conclude that the Killing form on  $\text{Lie}(E)$  has signature  $-24$ , as claimed.  $\square$

*Remark 72* We can determine the centralizer and the real form of  $E_8$  also in the excluded case in the proof where both  $\mathfrak{sl}_2$ 's have index 2. As in Lemma 62, the

centralizer is a real form of  $\mathrm{Sp}_{4,\mathbb{C}} \times \mathrm{Sp}_{4,\mathbb{C}}$ . Decomposition (6.6) shows that complex conjugation interchanges the two  $\mathrm{Sp}_{4,\mathbb{C}}$  terms, so the centralizer is  $R(\mathrm{Sp}_{4,\mathbb{C}})$ . Complex conjugation interchanges the irreducible representations appearing in Eq. (1.1) in pairs (contributing 0 to the signature of the Killing form  $\kappa_E$  of  $E$ ), except for  $2 \otimes 2 \otimes V_{2,2}$ , which has dimension  $8^2$ . This last piece breaks up into a 36-dimensional even subspace, and a 28-dimensional odd subspace, contributing 8 to the signature of  $\kappa_E$  and proving that the resulting real form of  $E_8$  is the split one.

## 8 No Theory of Everything in a Real Form of $E_8$

In the decomposition of  $\mathrm{Lie}(E) \otimes \mathbb{C}$  from Eq. (1.1), the integers  $m, n$  are positive, so (ToE2) implies

$$V_{m,n} = 0 \quad \text{if } m \geq 4 \text{ or } n \geq 4. \quad (\text{ToE2}')$$

We prove the following strengthening of the real case of Theorem 12:

**Lemma 81** *If subgroups  $\mathrm{SL}(2, \mathbb{C})$  and  $G$  of a real form  $E$  of  $E_8$  satisfy (ToE1) and (ToE2'), then  $V_{1,2}$  is a self-conjugate representation of  $G$ , i.e., (ToE3) fails.*

*Proof* As in the proof of Proposition 71, over the complex numbers we get two copies of  $\mathfrak{sl}_2$  that embed in  $E_8$  with the same index, which is 1 or 2.

If the index is 1, we are in the case of that proposition. The  $-1$ -eigenspace in  $\mathrm{Lie}(E)$  (of the element  $-1$  in the center of  $\mathrm{SL}(2, \mathbb{C})$ ) is a real representation of  $\mathrm{SL}(2, \mathbb{C}) \cdot G$ , and  $G$  is contained in a copy of  $\mathrm{Spin}(12 - a, a)$  for  $a = 1, 3$ , or  $5$ . As in the proof of the proposition, there is a representation  $W$  of  $\mathrm{SL}(2, \mathbb{C}) \times \mathrm{Spin}(12 - a, a)$  defined over  $\mathbb{R}$  that is isomorphic to

$$(2 \otimes 1 \otimes S_+) \oplus (1 \otimes 2 \otimes S_-)$$

over  $\mathbb{C}$ . Now  $G$  is contained in the maximal compact subgroup of  $\mathrm{Spin}(12 - a, a)$ , i.e.,  $\mathrm{Lie}(G)$  is a subalgebra of  $\mathfrak{so}(11)$ ,  $\mathfrak{so}(9) \times \mathfrak{so}(3)$ , or  $\mathfrak{so}(7) \times \mathfrak{so}(5)$ . The restriction of the two half-spin representations of  $\mathrm{Spin}(12 - a, a)$  to the compact subalgebra are equivalent (25, p. 264), and we see that in each case the restriction is *quaternionic*. (To see this, one uses the standard fact that the spin representation of  $\mathfrak{so}(2\ell + 1)$  is real for  $\ell \equiv 0, 3 \pmod{4}$  and quaternionic for  $\ell \equiv 1, 2 \pmod{4}$ .) That is, the restrictions of  $S_+$ ,  $S_-$ , and their complex conjugates to the maximal compact subgroup are all equivalent (over  $\mathbb{C}$ ), hence the same is true for their further restrictions to  $G$ , and (ToE3) fails.

If the index is 2, then  $G$  is contained in a real form of  $\mathrm{Sp}_{4,\mathbb{C}} \times \mathrm{Sp}_{4,\mathbb{C}}$  by Lemma 62. When we decompose  $\mathfrak{e}_8$  as in Eq. (1.1), we find  $V_{2,1}$  and  $V_{1,2}$  as in decomposition (6.6). As complex conjugation interchanges these two representations, it follows that complex conjugation interchanges the two  $\mathrm{Sp}_{4,\mathbb{C}}$  factors, i.e., the centralizer of  $\mathrm{SL}(2, \mathbb{C})$  has as identity component the transfer  $R(\mathrm{Sp}_{4,\mathbb{C}})$  of  $\mathrm{Sp}_{4,\mathbb{C}}$ . Its maximal compact subgroup is the compact form of  $\mathrm{Sp}_{4,\mathbb{C}}$  (also known as  $\mathrm{Spin}(5)$ ), all of whose irreducible representations are self-conjugate. Therefore, (ToE3) fails.  $\square$

*Remark 82* It is worthwhile noting that, in each of the three cases in Proposition 71 (the three cases where (ToE2) holds), it is possible to embed  $G_{\text{SM}}$  in the centralizer, thus showing that (ToE1) is satisfied. Given such an embedding, a simple computation verifies explicitly that  $S_+$  has a self-conjugate structure as a representation of  $G_{\text{SM}}$ .

First consider  $\text{Spin}(11, 1)$ . There is an obvious embedding of  $G_{\text{GUT}} := \text{Spin}(10)$ . Under this embedding,  $S_+$  decomposes as the direct sum of the two half-spinor representations, i.e., as a generation and an anti-generation.

For  $\text{Spin}(7, 5)$ , there is an obvious embedding of the Pati-Salam group,  $G_{\text{GUT}} := (\text{Spin}(6) \times \text{Spin}(4))/(\mathbb{Z}/2\mathbb{Z})$ . Again,  $S_+$  decomposes as the direct sum of a generation and an anti-generation.

Finally,  $\text{Spin}(3, 9)$  contains  $(\text{SU}(3) \times \text{SU}(2) \times \text{SU}(2) \times \text{U}(1))/(\mathbb{Z}/6\mathbb{Z})$  as a subgroup. Under this subgroup,

$$S_+ = (3, 2, 2)_{1/6} \oplus (\bar{3}, 2, 2)_{-1/6} + (1, 2, 2)_{-1/2} + (1, 2, 2)_{1/2},$$

where the subscript indicates the  $\text{U}(1)$  weights, and the overall normalization is chosen to agree with the physicists' convention for the weights of the Standard Model's  $\text{U}(1)_Y$ . Embedding the  $\text{SU}(2)$  of the Standard Model in one of the two  $\text{SU}(2)$ s, we obtain an embedding of  $G_{\text{SM}} \subset \text{Spin}(3, 9)$  where, again  $S_+$  has a self-conjugate structure as a representation of  $G_{\text{SM}}$ .

## 9 No Theory of Everything in Complex $E_8$

We now complete the proof of Theorem 12 by proving the following strengthening of the complex case.

**Lemma 91** *If subgroups  $\text{SL}(2, \mathbb{C})$  and  $G$  of  $R(E_{8, \mathbb{C}})$  satisfy (ToE1) and (ToE2'), then  $V_{1,2}$  is a self-conjugate representation of  $G$ , i.e., (ToE3) fails.*

First, recall the definition of the transfer  $R(H_{\mathbb{C}})$  of a complex group  $H_{\mathbb{C}}$  as described, e.g., in (30, 2.1.2). Its complexification can be viewed as  $H_{\mathbb{C}} \times H_{\mathbb{C}}$ , where complex conjugation acts via

$$\overline{(h_1, h_2)} = (\bar{h}_2, \bar{h}_1).$$

One can view  $R(H_{\mathbb{C}})$  as the subgroup of the complexification consisting of elements fixed by complex conjugation.

Now consider an inclusion  $\phi : \text{SL}(2, \mathbb{C}) = R(\text{SL}_{2, \mathbb{C}}) \hookrightarrow R(E_{8, \mathbb{C}})$ . Complexifying, we identify  $R(\text{SL}_{2, \mathbb{C}}) \otimes \mathbb{C}$  with  $\text{SL}_{2, \mathbb{C}} \times \text{SL}_{2, \mathbb{C}}$  and similarly for  $R(E_{8, \mathbb{C}})$ , and write out  $\phi$  as

$$\phi(h_1, h_2) = (\phi_1(h_1)\phi_2(h_2), \psi_1(h_1)\psi_2(h_2)) \quad (9.1)$$

for some homomorphisms  $\phi_1, \phi_2, \psi_1, \psi_2 : \text{SL}_{2, \mathbb{C}} \rightarrow E_{8, \mathbb{C}}$ . As  $\phi$  is defined over  $\mathbb{R}$ , we have:

$$\phi(h_1, h_2) = \overline{\phi(\bar{h}_2, \bar{h}_1)} = (\overline{\psi_1(\bar{h}_2)\psi_2(\bar{h}_1)}, \overline{\phi_1(\bar{h}_2)\phi_2(\bar{h}_1)}),$$

and it follows that  $\psi_1(h_1) = \overline{\phi_2(\bar{h}_1)}$  and  $\psi_2(h_2) = \overline{\phi_1(\bar{h}_2)}$ . Conversely, given any two homomorphisms  $\phi_1, \phi_2 : \text{SL}_{2, \mathbb{C}} \rightarrow E_{8, \mathbb{C}}$  (over  $\mathbb{C}$ ) with commuting images, the same equations define a homomorphism  $\phi : \text{SL}(2, \mathbb{C}) \rightarrow R(E_{8, \mathbb{C}})$  defined over  $\mathbb{R}$ .

*Proof of Lemma 91* Write  $Z$  for the identity component of the centralizer of the image of the map  $\phi_1 \times \phi_2 : \mathrm{SL}_{2,\mathbb{C}} \times \mathrm{SL}_{2,\mathbb{C}} \rightarrow E_{8,\mathbb{C}}$  from Eq. (9.1). Clearly,  $G$  is contained in the transfer  $R(Z)$  of  $Z$ . In each of the cases below, we verify that

$$Z \text{ is semisimple and } -1 \text{ is in the Weyl group of } Z. \quad (9.2)$$

It follows from this that the maximal compact subgroup of  $R(Z)$  is the compact real form  $Z_{\mathbb{R}}$  of  $Z$  and that  $Z_{\mathbb{R}}$  is an *inner* form. Hence every irreducible representation of  $Z_{\mathbb{R}}$  is real or quaternionic; hence every representation of  $Z_{\mathbb{R}}$  is self-conjugate. That is, (ToE3) fails, which is the desired contradiction.

*Case 1  $\phi_1$  or  $\phi_2$  is trivial.* Consider the easiest-to-understand case where  $\phi_1$  or  $\phi_2$  is the zero map, say  $\phi_2$ . In the notation of Eq. (9.1),  $\phi(h_1, h_2) = (\phi_1(h_1), \phi_1(\overline{h_2}))$ , i.e.,  $\phi$  is the transfer of the homomorphism  $\phi_1 : \mathrm{SL}_{2,\mathbb{C}} \rightarrow E_{8,\mathbb{C}}$ . By Proposition 53,  $\phi_1$  has index 1 or 2. If  $\phi_1$  has index 1, then  $Z$  is simple of type  $E_7$  by Example 54, hence Property (9.2) holds. If  $\phi_1$  has index 2, then  $\mathrm{Lie}(Z)$  is isomorphic to  $\mathfrak{so}_{13,\mathbb{C}}$  by Lemma 61, and again Property (9.2) holds.

*Case 2 Neither  $\phi_1$  nor  $\phi_2$  is trivial.* Now suppose that neither  $\phi_1$  nor  $\phi_2$  is trivial. Again, Proposition 53 implies that  $\phi_1$  and  $\phi_2$  have Dynkin index 1 or 2.

If  $\phi_1$  and  $\phi_2$  both have index 1, then (over  $\mathbb{C}$ ) the homomorphism  $\phi_1 \times \phi_2$  is the one from the proof of Proposition 71 and  $Z$  is the standard  $D_6$  subgroup of  $E_{8,\mathbb{C}}$  and Property (9.2) holds.

Now suppose that  $\phi_1$  and  $\phi_2$  both have index 2. As  $\phi$  is an injection, it is not possible that  $\phi_1$  and  $\phi_2$  both vanish on  $-1 \in \mathrm{SL}_{2,\mathbb{C}}$ , and it follows from the proof of Lemma 62 that  $\phi_1 \times \phi_2$  is an injection as in the statement of Lemma 62. In particular,  $Z$  has Lie algebra  $\mathfrak{sp}_{4,\mathbb{C}} \times \mathfrak{sp}_{4,\mathbb{C}}$  of type  $B_2 \times B_2$  and Property (9.2) holds. Note that (ToE2) fails in this case by Table (6.5).

Suppose finally that  $\phi_1$  has index 1 and  $\phi_2$  has index 2. We conjugate so that  $\phi_2(\mathfrak{sl}_2)$  is the copy of  $\mathfrak{sl}_2$  from Example 52, and (by Lemma 41 for the centralizer  $\mathfrak{so}_{13}$  of  $\phi_2(\mathfrak{sl}_2)$ ) we can take  $\phi_1(\mathfrak{sl}_2)$  to be a copy of  $\mathfrak{sl}_2$  generated by the highest root of  $\mathfrak{so}_{13}$ . Calculating as described in 55 gives the following table of multiplicities for the irreducible representation  $m \otimes n$  of  $\mathfrak{sl}_2 \times \mathfrak{sl}_2$  in  $\mathfrak{e}_8$ :

$$\begin{array}{c|cccc} & 1 & 2 & 3 & m \\ \hline 1 & 39 & 18 & 1 & \\ n \ 2 & 32 & 16 & 0 & \\ 3 & 10 & 2 & 0 & \end{array} \quad (9.3)$$

In particular, the  $A_1 \times B_4$  subgroup of  $\mathrm{Spin}_{13}$  that centralizes the image of  $\phi_1 \times \phi_2$  is all of the identity component  $Z$  of the centralizer in  $E_8$ . Again Property (9.2) holds. (Of course, Table (9.3) shows that (ToE2) fails.)  $\square$

## 10 Relaxing (ToE2) to (ToE2')

Combining Lemmas 81 and 91 gives a proof not only of Theorem 12, but of the following stronger statement.

**Theorem 101** *There are no subgroups  $\mathrm{SL}(2, \mathbb{C}) \cdot G$  satisfying (ToE1), (ToE2'), and (ToE3) in the (transfer of the) complex  $E_8$  or any real form of  $E_8$ .*

We retained hypothesis (ToE2) in the Introduction because that is what is demanded by physics. Technically, it is *possible* for  $V_{2,3}$  and  $V_{3,2}$  to be nonzero in an interacting theory—so (ToE2) is false but (ToE2') still holds—but only in the presence of local supersymmetry (i.e., in supergravity theories) (31). Lisi's framework is not compatible with local supersymmetry, so we excluded this possibility above.

For real forms of  $E_8$ , weakening (ToE2) to (ToE2') only adds the case of  $E_{8(8)}$ , with  $G_{\max} = \mathrm{Spin}(5)$ , where we find

$$V_{3,2} \simeq V_{2,3} = 4, \quad V_{2,1} \simeq V_{1,2} = 4 \oplus 16, \quad (10.1)$$

and we have indicated the irreducible representations of  $\mathrm{Spin}(5)$  by their dimensions. Because the gravitinos transform nontrivially under  $G_{\max}$  and because of their multiplicity, the only consistent possibility would be a gauged  $\mathcal{N} = 4$  supergravity theory (for a recent review of such theories, see (32)). Unfortunately, the rest of the matter content (it suffices to look at  $V_{2,1}$ ) is not compatible with  $\mathcal{N} = 4$  supersymmetry. Even if it were,  $\mathcal{N} = 4$  supersymmetry would, of course, necessitate that the theory be non-chiral, making it unsuitable as a candidate Theory of Everything.

To summarize the results of this section, the previous subsection, and Remark 72, weakening (ToE2) to (ToE2') adds only three additional entries to Table 1.2.

$E$	$G_{\max}$	$V_{3,2}$	$V_{2,1}$	
$E_{8(8)}$	$\mathrm{Spin}(5)$	4	$4 \oplus 16$	
$R(E_{8,C})$	$\mathrm{Spin}(5) \times \mathrm{Spin}(5)$	$(4, 1) \oplus (1, 4)$	$(4, 5) \oplus (5, 4)$	·
$R(E_{8,C})$	$\mathrm{SU}(2) \times \mathrm{Spin}(9)$	$(2, 1)$	$(2, 9) \oplus (2, 16)$	

(10.2)

In each case the fermion representations,  $V_{2,1} \simeq V_{1,2}$  and  $V_{3,2} \simeq V_{2,3}$ , are pseudoreal representations of  $G_{\max}$ .

## 11 Conclusion

In Subsect. 3.3 above, we observed by an easy dimension count that no proposed Theory of Everything constructed using subgroups of a real form  $E$  of  $E_8$  has a sufficient number of weight vectors in the  $-1$ -eigenspace to identify with all known fermions. The proof of our Theorem 12 was quite a bit more complicated, but it also gives much more. It shows that you cannot obtain a *chiral* gauge theory for *any* candidate ToE subgroup of  $E$ , whether  $E$  is a real form or the complex form of  $E_8$ . In particular, it is impossible to obtain even the 1-generation Standard Model (in the sense of Definition 25) in this fashion.

**Acknowledgements.** The second author thanks Fred Helenius for pointing out a reference and Patrick Brosnan for suggesting this project. The authors' research was supported by the National Science Foundation under grant nos. PHY-0455649 (Distler) and DMS-0653502 (Garibaldi).

## References

1. Lisi, A.G.: *An exceptionally simple theory of everything*. <http://arxiv.org/abs/0711.0770> V1 [hep-th], 2007
2. Borel, A.: *Linear Algebraic Groups*. Vol. **126** of Graduate Texts in Mathematics, 2nd ed., New York: Springer-Verlag, 1991
3. Serre, J.-P.: *Local Fields*. Vol. **67** of Graduate Texts in Mathematics. New York-Berlin: Springer, 1979
4. J. Tits (1971) Représentations linéaires irréductibles d’un groupe réductif sur un corps quelconque *J. Reine Angew. Math.* **247** 196 – 220
5. H. Georgi S. Glashow (1974) Unity of all elementary particle forces *Phys. Rev. Lett.* **32** 438 – 441
6. J. Pati A. Salam (1974) Lepton number as the fourth color *Phys. Rev.* **D10** 275 – 289
7. Distler, J.: *Superconnections for dummies*. Weblog entry, May 12, 2008, available at <http://golem.ph.utexas.edu/~distler/blog/archives/001680.html>
8. Streater, R.F., Wightman, A.S.: *PCT, Spin and Statistics, and All That*. Princeton Landmarks in Mathematics and Physics, Princeton, NJ: Princeton University Press, 2000
9. Weinberg, S.: *The Quantum Theory of Fields*. 3 volumes. Cambridge: Cambridge University Press, 1995-2000
10. ’t Hooft, G.: In: *Recent Developments in Gauge Theories*, Cargèse France, ’t Hooft, G., Itzykson, C., Jaffe, A., Lehmann, H., Mitter, P.K., Singer, I., Stora, R., eds., no. **59** in Proceedings of the Nato Advanced Study Series B, London: Plenum Press, 1980
11. Lisi, A.G.: *First person: A. Garrett Lisi*, *Financial Times*, (April 25, 2009), available at <http://www.ft.com/cms/s/2/ebead98a-2d71-11de-9eba-00144feabdc0.html>
12. Helgason, S.: *Differential Geometry, Lie Groups, and Symmetric Spaces*. vol. **34** of Graduate Studies in Mathematics, Providence, RI: Amer. Math. Soc., 2001
13. J.-P. Serre (2004) On the values of the characters of compact Lie groups *Oberwolfach Reports* **1** 1 666 – 667
14. Elashvili, A., Kac, V., Vinberg, E.: *On exceptional nilpotents in semisimple Lie algebras*. <http://arxiv.org/abs/0812.1571>, V1 [math.GR], 2008
15. Dynkin, E.: Semisimple subalgebras of semisimple Lie algebras. Amer. Math. Soc. Transl. (2) **6**, 111–244, (1957) [Russian original: Mat. Sbornik N.S. **30(72)**, 349–462, (1952)]
16. Bourbaki, N.: *Lie Groups and Lie Algebras*: Chapters 7–9. Berlin: Springer-Verlag, 2005
17. B. Gross G. Nebe (2004) Globally maximal arithmetic groups *J. Algebra* **272** 2 625 – 642
18. Humphreys, J.: *Introduction to Lie Algebras and Representation Theory*. Vol. **9** of Graduate Texts in Mathematics. Third printing, revised Berlin-Heidelberg-New York: Springer-Verlag, 1980
19. Collingwood, D., McGovern, W.: *Nilpotent Orbits in Semisimple Lie Algebras*

- bras*. New York: Van Nostrand Reinhold, 1993
20. Onishchik, A., Vinberg, E.: *Lie Groups and Lie Algebras III*. Vol. **41** of Encyclopaedia Math. Sci. Berlin-Heidelberg-New York: Springer, 1994
21. Carter, R.: *Finite Groups of Lie Type: Conjugacy Classes and Complex Characters*. New York: Wiley-Interscience, 1985
22. Bourbaki, N. *Lie Groups and Lie Algebras: Chapters 4–6*, Berlin: Springer-Verlag, 2002
23. N. Vavilov (2004) Do it yourself structure constants for Lie algebras of type  $E_\ell$  *J. Math. Sci. (N.Y.)* **120** 4 1513 – 1548
24. A. Èlašvili (1975) Centralizers of nilpotent elements in semisimple Lie algebras *Sakharth. SSR Mecn. Akad. Math. Inst. Šrom.* **46** 109 – 132
25. McKay, W., Patera, J.: *Tables of Dimensions, Indices, and Branching Rules for Representations of Simple Lie Algebras*. Vol. **69** of Lecture Notes in Pure and Applied Mathematics. New York: Marcel Dekker Inc., 1981
26. Springer, T., Steinberg, R.: Conjugacy classes. In: *Seminar on Algebraic Groups and Related Finite Groups (The Institute for Advanced Study, Princeton, N.J., 1968/69)*, Vol. **131** of Lecture Notes in Math., Berlin: Springer, 1970, pp. 167–266
27. Steinberg, R.: *Lectures on Chevalley Groups*. New Haven, CT: Yale University, 1968
28. Tits, J.: *Tabellen zu den einfachen Lie Gruppen und ihren Darstellungen*. Vol. **40** of Lecture Notes in Mathematics. Berlin: Springer-Verlag, 1967
29. R. Garibaldi (2001) Clifford algebras of hyperbolic involutions *Math. Zeit.* **236** 321 – 349
30. Platonov, V., Rapinchuk, A.: *Algebraic Groups and Number Theory*. Boston, MA: Academic Press, 1994
31. M.T. Grisaru H.N. Pendleton (1977) Soft spin 3/2 fermions require gravity and supersymmetry *Phys. Lett.* **B67** 323
32. J. Schon M. Weidner (2006) Gauged  $N=4$  supergravities *JHEP* **05** 034