



ELSEVIER

ScienceDirect

Nuclear Physics B ●● (●●●) ●●●●●

NUCLEAR
PHYSICS Bwww.elsevier.com/locate/nuclphysb

Higher anomalies, higher symmetries, and cobordisms III: QCD matter phases anew

Zheyuan Wan^a, Juven Wang^{b,c,*}^a *Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China*^b *Center of Mathematical Sciences and Applications, Harvard University, Cambridge, MA 02138, USA*^c *School of Natural Sciences, Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA*

Received 12 March 2020; accepted 8 April 2020

Editor: Clay Córdova

Abstract

We explore QCD₄ quark matter, the μ -T (chemical potential-temperature) phase diagram, possible 't Hooft anomalies, and topological terms, via non-perturbative tools of cobordism theory and higher anomaly matching. We focus on quarks in 3-color and 3-flavor on bi-fundamentals of SU(3), then analyze the continuous and discrete global symmetries and pay careful attention to finite group sectors. We input constraints from $T = CP$ or CT time-reversal symmetries, implementing QCD on unorientable spacetimes and distinct topology. Examined phases include the high T QGP (quark-gluon plasma/liquid), the low T ChSB (chiral symmetry breaking), 2SC (2-color superconductivity) and CFL (3-color-flavor locking superconductivity) at high density. We introduce a possibly useful but only *approximate* higher anomaly, involving discrete 0-form axial and 1-form mixed chiral-flavor-locked center symmetries, matched by the above four QCD phases. We also enlist as much as possible, but without identifying all of, 't Hooft anomalies and topological terms relevant to Symmetry Protected/Enriched Topological states (SPTs/SETs) of gauged SU(2) or SU(3) QCD_d-like matter theories in general in any spacetime dimensions $d = 2, 3, 4, 5$ via cobordism.

© 2020 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author at: Center of Mathematical Sciences and Applications, Harvard University, Cambridge, MA 02138, USA.

E-mail addresses: wanzheyuan@mail.tsinghua.edu.cn (Z. Wan), jw@cmsa.fas.harvard.edu (J. Wang).

URL: <http://sns.ias.edu/~juven/> (J. Wang).

<https://doi.org/10.1016/j.nuclphysb.2020.115016>

0550-3213/© 2020 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

Contents

1 Introduction and summary 3

1.1. Physics in QCD quark matter 3

1.2. Approximate higher anomaly constraint on the QCD phase diagram 9

1.3. Mathematical primer 13

2. QCD symmetries, anomalies and topological terms without time-reversal 14

2.1. Chiral symmetry breaking $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V)}{\mathbb{Z}_{3c,v} \times \mathbb{Z}_{3f,v}}$ as $(\text{Spin}(d) \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3})$ 15

2.2. 3-Color-flavor locking superconductivity $SU(3)_{C+L+R} \times \mathbb{Z}_{2,V}^F$ as $\text{Spin} \times SU(3)$ 19

2.3. 2-Color superconductivity: $\frac{([SU(2c)_{V,rg}] \times SU(2f)_{L,ud} \times SU(2f)_{R,ud} \times U(1f)_{V,s} \times U(1c)_{V,b})}{\mathbb{Z}_{2,V}^F}$ as $\text{Spin} \times \mathbb{Z}_2 \text{Spin}(4) \times U(1) \times U(1)$ 20

2.4. Quark gluon plasma/liquid $\frac{([SU(3)_V] \times SU(3)_L \times SU(3)_R \times U(1)_V)}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V}}$ as $\text{Spin}(d) \times \mathbb{Z}_2 \frac{U(3)_L \times U(3)_R}{(\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,A} \times U(1)_A)}$ 23

3. QCD symmetries, anomalies and topological terms with time-reversal 26

3.1. Chiral symmetry breaking $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V) \times \mathbb{Z}_4^T}{\mathbb{Z}_{3c,v} \times \mathbb{Z}_{3f,v} \times \mathbb{Z}_{2,v}^F}$ 28

3.1.1. $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V) \times \mathbb{Z}_4^{TF}}{\mathbb{Z}_{3c,v} \times \mathbb{Z}_{3f,v} \times \mathbb{Z}_{2,v}^F}$ as $(\text{Pin}^+(d) \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3})$ and $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V) \times \mathbb{Z}_2^T}{\mathbb{Z}_{3c,v} \times \mathbb{Z}_{3f,v}}$ as $(\text{Pin}^-(d) \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3})$ 28

3.2. 3-Color-flavor locking superconductivity $SU(3)_{C+L+R} \times \mathbb{Z}_4^T$ 29

3.2.1. $SU(3)_{C+L+R} \times \mathbb{Z}_4^{TF}$ as $\text{Pin}^+ \times SU(3)$ 29

3.2.2. $SU(3)_{C+L+R} \times \mathbb{Z}_{2,V}^F \times \mathbb{Z}_2^T$ as $\text{Pin}^- \times SU(3)$ 31

3.3. 2-Color superconductivity: $\frac{([SU(2c)_{V,rg}] \times SU(2f)_{L,ud} \times SU(2f)_{R,ud} \times U(1f)_{V,s} \times U(1c)_{V,b}) \times \mathbb{Z}_4^T}{(\mathbb{Z}_{2,V}^F)^2}$ 32

3.3.1. $\frac{([SU(2c)_{V,rg}] \times SU(2f)_{L,ud} \times SU(2f)_{R,ud} \times U(1f)_{V,s} \times U(1c)_{V,b}) \times \mathbb{Z}_4^{TF}}{(\mathbb{Z}_{2,V}^F)^2}$ as $\text{Pin}^+ \times \mathbb{Z}_2 \text{Spin}(4) \times U(1) \times U(1)$ 32

3.3.2. $\frac{([SU(2c)_{V,rg}] \times SU(2f)_{L,ud} \times SU(2f)_{R,ud} \times U(1f)_{V,s} \times U(1c)_{V,b}) \times \mathbb{Z}_2^T}{\mathbb{Z}_{2,V}^F}$ as $\text{Pin}^- \times \mathbb{Z}_2 \text{Spin}(4) \times U(1) \times U(1)$ 33

3.4. Quark gluon plasma/liquid $\frac{([SU(3)_V] \times SU(3)_L \times SU(3)_R \times U(1)_V) \times \mathbb{Z}_4^T}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V} \times \mathbb{Z}_{2,V}^F}$ 35

3.4.1. $\frac{([SU(3)_V] \times SU(3)_L \times SU(3)_R \times U(1)_V) \times \mathbb{Z}_4^{TF}}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V} \times \mathbb{Z}_{2,V}^F}$ as $\text{Pin}^+ \times \mathbb{Z}_2 \frac{U(3)_L \times U(3)_R}{(\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,A} \times U(1)_A)}$ and $\frac{([SU(3)_V] \times SU(3)_L \times SU(3)_R \times U(1)_V) \times \mathbb{Z}_2^T}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V}}$ as $\text{Pin}^- \times \mathbb{Z}_2 \frac{U(3)_L \times U(3)_R}{(\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,A} \times U(1)_A)}$ 35

Acknowledgements 39

References 39

1. Introduction and summary

1.1. Physics in QCD quark matter

We are made of atoms, which are made out of quarks, the particles¹ of Quantum Chromodynamics (QCD) vacuum, plus some electrons. The majority of our mass is from the mass of nuclei. While about the 1%~2% of our mass is from the Higgs condensate, the surprising significant around 98%~99% of our mass is from QCD chiral condensate.² Meanwhile, we live in the chiral symmetry breaking (ChSB) phase of QCD vacuum. In order to investigate the nature of QCD matter and its vacuum structure, it is helpful to move out from this particular vacuum (ChSB) to other new foreign phases. In condensed matter language, we try to explore other unfamiliar foreign phases outside the familiar domestic phase, away from the ground state (i.e., vacuum) we live in, by tuning parameters in the QCD phase diagram (see recent selected reviews [1–3]). Namely, we should explore different new vacua or ground state structures and their excitation spectra.

In this work, we will look at some simplified ideal toy models. One model is that quarks are nearly massless and on bi-fundamentals of SU(3). Here the quarks are the Dirac spinors in 3+1 dimensional spacetime (we denoted as 3+1D or 4d). We will consider various types of curved spacetime manifolds with different topology and with Spin structure, Pin⁺, or Pin⁻ or other twisted structures.³ In physics language, quarks are 4d Dirac fermions, in the fundamental representation 3 of SU(3) color gauge group and fundamental representation of SU(3) flavor global symmetry group. We denote them as the representation (Rep) 3_c in SU(3_c)_V for color and 3_f in SU(3_f)_V for flavor, where subindex V stands for the vector symmetry. Follow the notations in [4], for the Euclidean path integral, we have the schematic partition function

$$\begin{aligned} & \int [Da][D\psi][D\bar{\psi}] \exp\left(-S_{E,\text{Dirac},(\mathbb{D}_a+m_q+\mu)} - S_{\text{YM}} - S_\theta\right) \\ & \equiv \int [Da][D\psi][D\bar{\psi}] \exp\left(-\int_{M_4} d^4x_E \sqrt{\det g} \left(\sum_q \bar{\psi}_q (\mathbb{D}_a + m_q + \mu) \psi_q\right)\right. \\ & \quad \left.- \int_{M_4} \left(\frac{1}{g^2} \text{Tr}(F_a \wedge \star F_a) + \frac{i\theta}{8\pi^2} \text{Tr}(F_a \wedge F_a)\right)\right). \end{aligned} \quad (1.1)$$

This path integral describes an SU(N_c) = SU(3_c) gauge theory with 1-form gauge field a and 2-form field strength F_a for the (exact) color $N_c = 3$ with three colors in fundamental: red (r),

¹ In particle physics, quarks are *elementary particles*. In condensed matter viewpoint, it may be beneficial to alternatively view the quarks as *quasiparticles*, quasi-excitations out of certain vacuum.

² The up quark u bare mass is about 2.2 MeV/ c^2 and the down quark d bare mass is about 4.79 MeV/ c^2 . So the bare mass of proton from uud quarks is about 9.4 MeV/ c^2 and the bare mass of neutron from udd quarks is about 11.9 MeV/ c^2 . But the rest mass of proton is about 938.3 MeV/ c^2 and the rest mass of neutron is about 939.6 MeV/ c^2 . Since nearly all of our (atomic) mass is concentrated in the nucleons, based on the ratio between the bare mass of bare quarks to the rest mass of proton/neutron is about 1%~2% contributed from the Higgs effect, this shows about the remained 98%~99% of the mass of baryonic matter is in fact the QCD binding energy.

³ We consider the smooth differentiable manifolds with a metric g as spacetime – if the fermions/spinor can live on them, we require Spin structure; if we require time-reversal $T = CP$, or CT or other reflection symmetries, we require Pin⁺, Pin⁻ or other semi-direct (\times) product or twisted structures between the spacetime tangent bundle TM and the gauge bundle E_G of the gauge group G . See more in the main text and see an overview of our setting in [4].

Table 1

Pairing of quark-quark condensate for 2SC (2-color superconductivity) and CFL (3-color-flavor locking superconductivity). The L and R are for left/right-handed spinors.

SC	Spin s	Momentum (angular/orbital) \vec{L}, \vec{k} L/R (Parity, J^P)	Color (SU(2)/SU(3))	Flavor (SU(2)/SU(3))	Condensate $\langle qq \rangle$
2SC	singlet ($\uparrow\downarrow - \downarrow\uparrow$) (ud, rg) 1 in $2 \otimes 2 = 1 \oplus 3$	0, 0 LL \pm RR even/odd; 0^\pm	($rg - gr$) singlet: 1 in $2 \otimes 2 = 1 \oplus 3$ color-anti-triplet: A in $3 \otimes 3 = \bar{3}_A \oplus 6_S$	($ud - du$) 1 in $2 \otimes 2 = 1 \oplus 3$ A in $3 \otimes 3 = \bar{3}_A \oplus 6_S$	$\langle \psi_{\alpha i}^T C \gamma^5 \psi_{\beta j} \rangle_{2SC}$ $\propto \epsilon_{\alpha\beta A} \epsilon_{ij B} \delta_3^A \delta_B^3 = \epsilon_{\alpha\beta 3} \epsilon_{ij 3}$
CFL	singlet ($\uparrow\downarrow - \downarrow\uparrow$) 1 in $2 \otimes 2 = 1 \oplus 3$	0, 0 LL \pm RR even/odd; 0^\pm	$\bar{3}_A$ in $3 \otimes 3 = \bar{3}_A \oplus 6_S$ not-SU(3)-singlet	$\bar{3}_A$ in $3 \otimes 3 = \bar{3}_A \oplus 6_S$	$\langle \psi_{\alpha i}^T C \gamma^5 \psi_{\beta j} \rangle_{CFL}$ $\propto \epsilon_{\alpha\beta A} \epsilon_{ij B} \delta_B^A$

Table 2

Charge conjugate matrix $C = i\gamma^2\gamma^0$ is unitary. All pairings are T -symmetry invariant. Under Parity P : $L \leftrightarrow R$ and $k \leftrightarrow -k$, but spin up/down (\uparrow / \downarrow) \leftrightarrow (\uparrow / \downarrow) is invariant. Under Time Reversal T : $L/R \leftrightarrow L/R$ invariant, but $k \leftrightarrow -k$, and the spin up/down flips (\uparrow / \downarrow) \leftrightarrow (\downarrow / \uparrow).

	Pairing function	Wavefunction	Parity P	Spin or orbital
$\psi^\dagger \Delta C \gamma^5 \psi^*$	$\Delta_k = \Delta_{-k}$	$(L, L\rangle + R, R\rangle) k, -k\rangle(\uparrow\downarrow\rangle - \downarrow\uparrow\rangle)$	Even	s wave/singlet
$\psi^\dagger \Delta C \gamma^5 \psi^*$	$\Delta_k = -\Delta_{-k}$	$(L, L\rangle - R, R\rangle) k, -k\rangle(\uparrow\downarrow\rangle + \downarrow\uparrow\rangle)$	Even	p wave/triplet
$\psi^\dagger \Delta C \psi^*$	$\Delta_k = \Delta_{-k}$	$(L, L\rangle - R, R\rangle) k, -k\rangle(\uparrow\downarrow\rangle - \downarrow\uparrow\rangle)$	Odd	s wave/singlet
$\psi^\dagger \Delta C \psi^*$	$\Delta_k = -\Delta_{-k}$	$(L, L\rangle + R, R\rangle) k, -k\rangle(\uparrow\downarrow\rangle + \downarrow\uparrow\rangle)$	Odd	p wave/triplet

green (g), and blue (b). It also describes the (approximate) $N_f = 3$ -flavor in fundamental for fermions ψ_q , where we choose the lightest bare quarks in nature: the up quark (u), the down quark (d), and the strange quark (s). At the massless limit $m_q = 0$, the topological θ -term can be absorbed by axial $U(1)_A$ rotations of quarks, and we may set $\theta = 0$. The fermion ψ_q are quarks that carry quantum number (denoted the quark quantum number q) of color (c), flavor (f), the spacetime self-rotational spin (s), whose quark-pairing can also carry the spatial momentum \vec{k} , angular momentum \vec{L} , and parity quantum number P , etc.⁴ See Table 1 and 2 for some examples of quantum numbers for different quark-pairing condensates. In Eq. (1.1), we can tune the temperature $T = \beta_E = \frac{1}{L_E}$ proportional to the inverse size of the Euclidean time circle L_E , and we can also tune the chemical potential μ , which changes the density of quark matter.

The standard lore from the pioneer studies of quark matter [1,2] teaches us that four dominant phases occur at different regions of QCD phase diagram drawn in the μ - T (chemical potential v.s. temperature) axes. Below we aim to revisit some of these four phases, and applying modern perspectives of symmetries and anomalies to constrain these quantum systems.

⁴ For convenience, sometime we specify the color (c) quantum number as the first subindex α and the flavor (f) quantum number as the second subindex i :

$$\psi_q \equiv \psi_{\alpha i}. \tag{1.2}$$

See Table 1 and 2 for some examples of quantum numbers for different quark-pairing condensates.

- (I) *Global Symmetries*: For symmetries, we explore and exhaust both continuous and discrete global symmetries and pay special attention to finite group sectors.
- (II) *Higher Symmetries*: For gauge theories, there are extended operators of lines and surfaces, etc. They can also carry quantum numbers thus also *charged* under the higher generalized global symmetries [5]. There are also corresponding symmetry generators as *charge* operators. We also need to pay attention to higher symmetries.
- (III) *Anomalies and Higher Anomalies*: Given the global symmetry, there can be potential obstructions to couple the symmetry to background gauge field or to subsequently gauging the symmetry — the phenomena are known as the 't Hooft anomalies [6]. For higher symmetries, there are also associated higher 't Hooft anomalies. By anomalies, we mean to include both
- **Perturbative local anomalies** calculable from perturbative Feynman diagram loop calculations, classified by the integer group \mathbb{Z} classes (or the so-called free classes). Selective examples include:
 - (1): Perturbative fermionic anomalies from chiral fermions with U(1) symmetry, originated from Adler-Bell-Jackiw (ABJ) anomalies [7,8] with \mathbb{Z} classes.
 - (2): Perturbative gravitational anomalies [9].
 - **Non-perturbative global anomalies**, classified by finite groups such as \mathbb{Z}_N (or the so-called torsion classes). Some selective examples from QFT or gravity include:
 - (1): An SU(2) anomaly of Witten in 4d or in 5d [10] with a \mathbb{Z}_2 class, which is a gauge anomaly.
 - (2): A new SU(2) anomaly in 4d or in 5d [11] with another \mathbb{Z}_2 class, which is a mixed gauge-gravity anomaly.
 - (3): Non-perturbative bosonic anomalies from finite-group-symmetry bosonic systems [12,13].
 - (4): Higher 't Hooft anomalies for a pure 4d SU(2) YM theory with a second-Chern-class topological term [14–16] (the $SU(2)_{\theta=\pi}$ YM): The higher anomaly involves a discrete 0-form \mathbb{Z}_2^T time-reversal symmetry and a 1-form center $\mathbb{Z}_{2,[1]}$ -symmetry. The first anomaly is discovered in [14]; later the anomaly is refined via a mathematical well-defined 5d co/bordism invariant as its invertible Topological Quantum Field Theories (iTQFTs) topological term, with additional new anomaly for four siblings of YM [15,16].
 - (5): Global gravitational anomalies [17].
- (IV) *Symmetry Protected Topological states (SPTs)/Symmetry Enriched Topologically ordered states (SETs) or Higher SPTs/SETs*: Quantum systems (usually the quantum vacuum or the ground state) can be protected by global symmetry in a topological way. These are known as the interacting generalizations of topological insulators (TI) and topological superconductors (TSC) [18–22] known as the SPTs for interacting bosons and interacting fermions (see the overview [23–25]). Ref. [26–28] propose mathematical theories of cobordism classifying these SPTs and their low energy iTQFTs. In the context that we require to apply is the SU(N) and time-reversal symmetry generalization of SPTs studied in Ref. [4] suitable for Yang-Mills and QCD systems. In this work, we will apply a generalized cobordism theory including the higher-SPTs classifications (given by higher classifying spaces and higher symmetries) based on the computations and tools in [29].

By keeping in minds and utilizing the above modern concepts of quantum systems and QFTs beyond Ginzburg-Landau symmetry-breaking paradigm, here we revisit the standard lore of four




Fig. 1. We revisit the QCD₄ matter phases in the schematic μ - T (chemical potential-temperature) phase diagram (mainly focus on $N_c = N_f = 3$): The high T QGP (quark-gluon plasma/liquid), the low T ChSB (chiral symmetry breaking), 2SC (2-color superconductivity) and CFL (3-color-flavor locking superconductivity) at high density. We do *not* attempt to address the nature of phase transitions in this figure, thus we intentionally make some of the phase boundaries blur (i.e., in this work, we do *not* attempt to address whether the phase boundary should be a crossover, or a first order or a second order continuous phase transition). The $\mathbb{Z}_{6,A}$ or $\mathbb{Z}_{4,A}$ means that part of the discrete axial symmetry (A) is broken: $\mathbb{Z}_{2N_f,A}$. In general, if the G_{sym} group is broken, we denote it as G_{sym} .




Fig. 2. Follow Fig. 1 (mainly focus on $N_c = N_f = 3$), but now we include possible time-reversal symmetries, which can be any reasonable \mathbb{Z}_2 -reflection symmetry by putting the Euclidean QCD₄ path integral on an unorientable spacetime. Here we choose a semi-direct product of $\times \mathbb{Z}_4^T$, where $\mathbb{Z}_4^T \supset \mathbb{Z}_2^F$. The semi-direct product is more general, which also includes the direct product case of $\times \mathbb{Z}_4^{CT}$ as the CT symmetry.

QCD quark matter phases [1,2] and list down their global symmetries⁵ in Fig. 1 (without time-reversal symmetry) and Fig. 2 (with certain time-reversal symmetries):

1. QGP (quark-gluon plasma/liquid) at high T :

For general N_c and N_f , we have the symmetry:

$$\frac{[SU(N_c)V] \times SU(N_f)_L \times SU(N_f)_R \times U(1)_V}{\mathbb{Z}_{N_c,V} \times \mathbb{Z}_{N_f,V}}, \tag{1.3}$$

⁵ It is worthwhile to emphasize the old literature may happen to pay less attention to the finite group and discrete sectors. However, the finite group and discrete sectors are important for topological terms and non-perturbative global anomalies later we compute from the cobordism group. Thus, we aim to be as precise as possible writing down the global symmetries, see also [4].

- 1 where the $[SU(N_c)_V]$ color is gauged as a gauge group.⁶ The $U(1)_V/\mathbb{Z}_{N_c,V}$ is the vector
 2 symmetry associated to the baryon number B conservation. The $SU(N_f)_L$ and $SU(N_f)_R$
 3 are the left/right-handed Weyl spinor $SU(N_f)$ flavor symmetries under the projection of
 4 $P_{L/R} = \frac{1 \mp \gamma^5}{2}$.⁷ We will mainly focus on $N_c = N_f = 3$.
 5 2. ChSB (chiral symmetry breaking) at low T and at lower densities and low μ :
 6 For general N_c and N_f , we have the symmetry:
 7

$$\frac{([SU(N_c)_V] \times SU(N_f)_V \times U(1)_V)}{\mathbb{Z}_{N_c,V} \times \mathbb{Z}_{N_f,V}}, \quad (1.4)$$

- 8 where the $[SU(N_c)_V]$ color is gauged. Due to the chiral condensate $\langle \bar{\psi}\psi \rangle \neq 0$ in this vac-
 9 uum, the $SU(N_f)_L$ and $SU(N_f)_R$ flavor symmetries of the left/right-handed Weyl spinor
 10 are broken down to the diagonal vector subgroup $SU(N_f)_V$. We will mainly focus on
 11 $N_c = N_f = 3$. In ChSB, by the spontaneously symmetry breaking (SSB) from QGP, we
 12 gain 8 pseudo-Goldstone bosons (as mesons: $\pi^0, \pi^\pm, K^0, \bar{K}^0, \eta, K^\pm$), while there is one
 13 massive η' meson. If the ChSB further forms the nucleon superfluid, the $U(1)_V$ has SSB,
 14 and we gain 1 more Goldstone boson.
 15 3. 2SC (2-color superconductivity) at low T and at intermediate densities and μ :
 16
 17
 18
 19

$$\frac{[SU(2_c)_{V,rg}] \times SU(2_f)_{L,ud} \times SU(2_f)_{R,ud} \times U(1)_{V,s} \times U(1)_{V,b}}{\mathbb{Z}_{2,V}^F}, \quad (1.5)$$

- 20 where the $[SU(2_c)_V]$ color is gauged thus there is an $SU(2)$ gauge theory. The 2SC pairing
 21 is shown in Table 1. The $\mathbb{Z}_{2,V}^F$ is the fermion parity symmetry, which is a vector symmetry.
 22 The 2SC pairs 2-flavor $u-d$ and 2-color $r-g$ both into $SU(2)$ singlets as a color supercon-
 23 ductor. Since the $[SU(3)_c]$ is broken to $[SU(2_c)]$, this results in 5 massive gluons. There are
 24 4 gapped quasiparticle fermions in the 2SC Bogoliubov basis. There are 5 un-paired and
 25 gapless quasiparticle fermion in the 2SC Bogoliubov basis. Thus many symmetries are still
 26 intact.
 27 4. CFL (3-color-flavor locking superconductivity) at low T and at high density and high μ :
 28 The $SU(3)_{C+L+R}$ means that the $SU(3_c)$ are locked and rotated in the opposite manner as
 29 the diagonal of $SU(3_f)_L \times SU(3_f)_R$. This is a CFL superconductor, with 9 gapped ($8 \oplus$
 30 1) quasiparticle fermion in the CFL Bogoliubov basis. By comparing the QGP and CFL
 31 phases, we see that the continuous Lie group generators broken down from 25 generators
 32 ($8 \times 3 + 1$) in QGP to the only 8 generators of $SU(3)_{C+L+R}$. The missing 17 generators can
 33 be accounted via: 7 massive gluons and 1 mixture of a photon+gluon gauge boson, there are
 34 ($8+1$) Goldstone bosons. The 8 unbroken $SU(3)_{C+L+R} \supset [U(1)]_{\bar{Q}}$ actually contains another
 35 mixture of the photon+gluon gauge boson. Including the fermion parity symmetry $\mathbb{Z}_{2,V}^F$, we
 36 have the global symmetry:
 37
 38
 39
 40
 41

$$SU(3)_{C+L+R} \times \mathbb{Z}_{2,V}^F, \quad (1.6)$$

42
 43
 44
 45 ⁶ The $[G_g]$ specifies that G_g is dynamically gauged.

46 ⁷ It is worthwhile mentioning that the discrete axial symmetry $\mathbb{Z}_{2N_f,A}$ that are not broken via the ABJ effect, still
 47 remains and sits inside: $\frac{[SU(N_c)_V] \times SU(N_f)_L \times SU(N_f)_R \times U(1)_V}{\mathbb{Z}_{N_c,V} \times \mathbb{Z}_{N_f,V}} \supset \mathbb{Z}_{2N_f,A}$.

although there is a part of the global symmetry containing the electromagnetic $[U(1)]_{\tilde{Q}}$ which is gauged not global.⁸

By including time reversal symmetries into the QCD system, we can choose any suitable outer automorphism of the color gauge or flavor global symmetry group as a \mathbb{Z}_2 -time reversal symmetry, which is a \mathbb{Z}_2 -reflection symmetry by putting the Euclidean QCD₄ path integral on an unorientable spacetime.

In fact, several recent works have attempted to study QCD matter phases based on the languages of higher symmetries and anomalies above. We should quickly overview some of these pioneer works:

- (1). Whether the color superconductivity can be topological in some way was questioned in Ref. [30]. What Ref. [30] concerns is the topological insulators (TI) / topological superconductors (TSC) in the free non-interacting quadratic mean-field Hamiltonian systems. Thus the classifications in Ref. [30] are only either 0 or \mathbb{Z} classes for mean-field free fermion systems. The new input in our context is that we consider fully interacting systems and enlist possible SPTs for these QCD matter phases by cobordism group classifications.
- (2). Ref. [31–34] explores a related system of 4d adjoint quantum chromodynamics (QCD₄) with an SU(2) gauge group and two massless adjoint Weyl fermions. Higher symmetries and higher anomalies play an important role. Depend on the complex mass parameters of fermions, we can land onto different phases, and there are interesting quantum phase transitions between bulk phases. There are implications and constraints for 3+1D deconfined quantum critical points (dQCP), quantum spin liquids (QSL) or fermionic liquids in condensed matter, and constraints on 3+1D ultraviolet-infrared (UV-IR) duality. See Ref. [34] for an overview of the proposed phases at quantum critical points.
- (3). Ref. [35,36] explores quantum phase transitions between Landau ordering phase transitions but beyond the Landau paradigm, for example, due to the effects of topological θ -terms. Ref. [36] suggests that SU(2) QCD₄ with large odd number of flavors of quarks could be a direct second order phase transition between two phases of U(1) gauge theories as well as between a U(1) gauge theory and a trivial vacuum (e.g. a Landau symmetry-breaking gapped paramagnet). The gauge group is enhanced to be non-Abelian at and only at the transition. It is characterized as Gauge Enhanced Quantum Critical Points.
- (4). Ref. [37–42] employs global anomalies or global inconsistency of QCD₄ matter to constrain its QCD phase diagram, including either QCD zero-T phase or thermal phase. Ref. [37–42], implicitly or explicitly, suggests that there are obstructions or anomalies to simultaneously gauge or preserve both (1) a discrete axial symmetry and (2) a discrete 1-form color or flavor center symmetry (e.g. twisted discrete flavor-symmetry background boundary conditions or chemical potential μ), and/or (3) under certain twisted boundary conditions — so preferably and naively either symmetries must be broken. In fact, there are still possibilities that symmetric gapped TQFTs can be constructed for some of these anomalies, for example, based on the symmetry extension method [43] or higher-symmetry extension method [34]. In any case, Ref. [37–42] can rule out trivial gapped phases under certain circumstances.

⁸ In fact, since the strong forces are much dominant than the electromagnetism, we will focus on the $[SU(3_c)]$ gauge sectors from the strong forces first, and treat the gauge sectors of electromagnetism $U(1)_{EM}$ separately later.

(5). *Quark-hadron continuity outside the Ginzburg-Landau paradigm*: Quark-hadron continuity [44] asserts that hadronic matter superfluid phase is continuously connected to color-superconductor without phase transitions when the μ increases. This proposal is based on Ginzburg-Landau theory where two sides of phases have the similar symmetry breaking patterns and the similar gapless and gapped energetic spectrum. Ref. [45] questions the quark-hadron continuity to be invalid, by suggesting there must be a phase transition due to the topological fractionalization of excitations are different. Ref. [46] re-analyzes the scenario based on higher-symmetry is not spontaneously broken, and found that the quark-hadron continuity is still plausible.

In the remained of this article, we like to point out a higher anomaly involving discrete 0-form axial chiral and 1-form mixed chiral-flavor-locked center symmetries that can be matched by the above four QCD phases. Then we will give a quick mathematical introduction of tools we used in Sec. 1.3. After then, we will list down various data and tables computed from cobordism theory, with a view toward the applications of QCD matter phases, to be studied in the future [47]. The QCD_d matter symmetries, anomalies, and topological terms without time-reversal symmetry, classified by the cobordism theory, are studied in Sec. 2 via a cobordism theory. The QCD_d matter symmetries, anomalies, and topological terms with time-reversal symmetry, classified by the cobordism theory, are studied in Sec. 3, in general in any spacetime dimensions $d = 2, 3, 4, 5$ via a cobordism theory.

1.2. Approximate higher anomaly constraint on the QCD phase diagram

In this section, we point out a higher 't Hooft anomaly involving discrete 0-form axial and 1-form mixed chiral-flavor-locked center symmetries can be matched by the above four QCD phases. Our approach is related but still somehow different from Ref. [37–42].

For simplicity, we will set $N_c = N_f = N$ below. If $N_c \neq N_f$, we just need to replace the N below to their greatest common divisor $\text{gcd}(N_c, N_f)$ and make some moderate but a straightforward generalization.

First, under some *assumptions* that we will comment later, we hope to point out that there is an *approximate* 1-form electric-magnetic (e-m) global symmetry $\mathbb{Z}_{N_{cf},[1]}$ that mixed between 1-form color and flavor (CF) center symmetry: $\mathbb{Z}_{N_{cf},[1]}$. To recall, focus on the kinematics of the UV path integral of QFT,

(●1) If we do not have quarks in the fundamental Rep of the color gauge group $[\text{SU}(N_c)_V]$, then we have the 1-form electric symmetry $\mathbb{Z}_{N_c,[1]}$ whose charged objects are the color $\text{SU}(N_c)$ fundamental Wilson line. If we have quarks in the fundamental Rep of $[\text{SU}(N_c)_V]$, then the $\mathbb{Z}_{N_c,[1]}$ is broken explicitly.

(●2) By looking at the largest symmetry at QCD phases — Eq. (1.3)'s QGP symmetry: $\frac{[\text{SU}(N_c)_V] \times \text{SU}(N_f)_L \times \text{SU}(N_f)_R \times \text{U}(1)_V}{\mathbb{Z}_{N_c,V} \times \mathbb{Z}_{N_f,V}}$ where $[\text{SU}(N_c)_V]$ is gauged, we still are left with part of the projective special unitary symmetry $\frac{\text{SU}(N_f)_V}{\mathbb{Z}_{N_f,V}} = \text{PSU}(N_f)_V$ from the flavor symmetry. We can regard it as a 1-form magnetic symmetry that can be coupled to the flavor symmetry background probed fields $\frac{\text{SU}(N_f)_V}{\mathbb{Z}_{N_f,V}} = \text{PSU}(N_f)_V$, by using the Stiefel-Whitney $w_2(\text{PSU}(N_f)_V)$. This idea is implemented already in [38–42].

(●3) Now we combine the above two 1-form e and m symmetries into a diagonal 1-form symmetry, and name it as 1-form electric-magnetic (e-m) global symmetry $\mathbb{Z}_{N_{cf},[1]}$. We define this

1 1-form e-m color-flavor-locked global symmetry $\mathbb{Z}_{N_{cf},[1]}$ as the diagonal symmetry that rotates
 2 oppositely the color-fundamental (W_c) and the flavor-fundamental (W_f) Wilson line along the
 3 1d curve γ^1 , called

$$4 \quad W_c W_f = \exp(q_c \oint_{\gamma^1} a_c) \exp(q_f \oint_{\gamma^1} a_f), \quad (1.7)$$

8 where the a_c is. The $(W_c W_f)$ have ends that can be opened by having bi-fundamental quark and
 9 anti-quark at each of two ends. Let us call the 2-dimensional surface operator that is the 1-form
 10 $\mathbb{Z}_{N_{cf},[1]}$ symmetry generator on 2d area Σ^2 as:

$$12 \quad U \sim (U_{e,c})(U_{m,f})^{-1} \sim \exp\left(\iint_{\Sigma^2} \Lambda\right) \exp\left(-\iint_{\Sigma^2} w_2(\text{PSU}(N_f)V)\right), \quad (1.8)$$

15 Here we write down the electric and magnetic 2-surface operators based on the conventions and
 16 notations in [16]. So that even if the bi-fundamental quarks can open up the color-fundamental
 17 and the flavor-fundamental Wilson line, it will *not* be charged under the 1-form e-m $\mathbb{Z}_{N_{cf},[1]}$
 18 symmetry, because the obtained phase is exactly cancelled:

$$20 \quad \langle (U)(W_c W_f) \rangle |_{\#_{U,(W_c W_f)}=1} = \exp(i \frac{2\pi}{N}) \exp(-i \frac{2\pi}{N}) \langle (U)(W_c W_f) \rangle |_{\#_{U,(W_c W_f)}=0} \\ 21 \quad = 1 \cdot \langle (U)(W_c W_f) \rangle |_{\#_{U,(W_c W_f)}=0} \quad (1.9)$$

23 when the 2d U surface and the 1d combined $W_c W_f$ line have a nontrivial linking number

$$25 \quad \#_{U,(W_c W_f)} = 1$$

27 in a 4d spacetime.⁹ Importantly, although bi-fundamental quarks can open up the $(W_c W_f)$ line,
 28 physically this means that this (approximate) 1-form e-m $\mathbb{Z}_{N_{cf},[1]}$ symmetry is not broken (ex-
 29 plicitly and kinematically) by the quark which must sit as bi-fundamental Rep in QCD₄!

30 However the 1-form e-m $\mathbb{Z}_{N_{cf},[1]}$ symmetry still acts nontrivially on the color-fundamental
 31 Wilson line because this linking

$$33 \quad \langle (U)(W_c) \rangle |_{\#_{U,(W_c W_f)}=1} = \exp(i \frac{2\pi}{N}) \langle (U)(W_c) \rangle |_{\#_{U,(W_c W_f)}=0} \quad (1.10)$$

35 shows W_c is charged under 1-form $\mathbb{Z}_{N_{cf},[1]}$ with a charge $\exp(i \frac{2\pi}{N})$. Similarly, the 1-form e-
 36 m $\mathbb{Z}_{N_{cf},[1]}$ symmetry still acts nontrivially on the flavor-fundamental Wilson line because this
 37 linking

$$39 \quad \langle (U)(W_f) \rangle |_{\#_{U,(W_c W_f)}=1} = \exp(-i \frac{2\pi}{N}) \langle (U)(W_f) \rangle |_{\#_{U,(W_c W_f)}=0} \quad (1.11)$$

41 shows W_f is charged under 1-form $\mathbb{Z}_{N_{cf},[1]}$ symmetry with a charge $\exp(-i \frac{2\pi}{N})$. We propose
 42 that there is an *approximate* anomaly mixing between the discrete $\mathbb{Z}_{2N_f,A}$ axial (chiral) sym-
 43 metry and the 1-form e-m color-flavor-locked $\mathbb{Z}_{N_{cf},[1]}$ global symmetry. We call the 2-form \mathbb{Z}_N

46 ⁹ The illustration and TQFT calculations of such linking number, link invariants and link configurations in the space-
 47 time picture, e.g. $\#_{(\Sigma^2, \gamma^1)} = 1$ on a 4-sphere S^4 , can be found in [48,49].

valued background field of 1-form $\mathbb{Z}_{N_{cf},[1]}$ as $B_{cf}^{(2)}$. We find that there is an anomaly captured by the $\mathbb{Z}_{2N_f,A}$ axial (chiral) symmetry transformation, such that the partition function \mathbf{Z} gains a fractionalized term

$$\mathbf{Z} \xrightarrow{\mathbb{Z}_{2N_f,A} \text{ transformation}} \mathbf{Z} \cdot \exp\left(\frac{-iN}{2\pi} \int_{M_4} (B_{cf}^{(2)} \wedge B_{cf}^{(2)})\right). \tag{1.12}$$

The fractionalized term $\exp\left(\frac{-iN}{2\pi} \int_{M_4} (B_{cf}^{(2)} \wedge B_{cf}^{(2)})\right)$ is not a 4d SPTs but a fractionalized 4d SPTs, not a counter term, thus cannot be absorbed into 4d, and should be regarded as a 5d higher-SPTs/higher-iTQFT – which in fact is an indicator of the 4d higher ’t Hooft anomaly of the QCD₄.

The caveat however is that this 4d higher ’t Hooft anomaly of the QCD₄ is only *approximate*. The subtlety is that it is only a precise anomaly if we also “gauge” the $\frac{SU(N_f)_V}{\mathbb{Z}_{N_f,V}}$ sector into the

$$\left[\frac{SU(N_c)_V \times SU(N_f)_V}{\mathbb{Z}_{\text{gcd}(N_c, N_f), V}} \right]$$

gauge theory. The disadvantages of our *approximate* anomaly (1.12) are that the flavor Wilson lines are only probed but not dynamical objects, so we do not really have the 1-form symmetry unless we at least *weakly* gauge the flavor symmetry.¹⁰ Thus another interpretation of our gauge theory and anomaly is indeed the *bi-fundamental gauge theory* in 3+1 dimensions studied in [50,51].

In comparison, other approaches in Ref. [37–42] also have their own disadvantages. For example, Ref. [39] derives a different kind of anomaly *only under a certain twisted flavor boundary condition* in 4d:

$$\mathbf{Z} \xrightarrow{\mathbb{Z}_{2N_f,A} \text{ transformation}} \mathbf{Z} \cdot \exp\left(\frac{-iN}{2\pi} \int_{M_4} (B_c^{(2)} \wedge B_f^{(2)})\right) \tag{1.13}$$

and its dimensional reduction in 3d

$$\mathbf{Z} \xrightarrow{\mathbb{Z}_{2N_f,A} \text{ transformation}} \mathbf{Z} \cdot \exp\left(\frac{-iN}{2\pi} \int_{M_4} (B_c^{(1)} \wedge B_f^{(2)})\right), \tag{1.14}$$

where B_c and B_f are color/flavor background fields respectively. Ref. [38] derives constraints that have limitations on the chemical potential and boundary conditions as well.

In Fig. 3, we show how our approximate anomaly in (1.12) is still required and can be matched by the four QCD₄ matter phases. We use the triple data with three inputs

$$(\mathbb{Z}_{2N_f,A}; \mathbb{Z}_{N_c\text{-shift}}, \mathbb{Z}_{N_{cf},[1]}) \tag{1.15}$$

which in Fig. 3 can be also denoted as

$$(\mathbb{Z}_{2N_f,A}; \mathbb{Z}_{N_c\text{-shift}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)}), \tag{1.16}$$

¹⁰ *Weakly* gauge the flavor symmetry is thus related to a certain twisted flavor boundary condition studied in, for examples, Ref. [38,39].




Fig. 3. Follow Fig. 1 and Fig. 2, QCD₄ matter phases shown here include the high T QGP (quark-gluon plasma/liq-uid), the low T ChSB (chiral symmetry breaking), 2SC (2-color superconductivity) and CFL (3-color-flavor locking superconductivity) at high density. The four phases can be matched and cancelled by our approximate anomaly (1.12),

$Z \xrightarrow{\mathbb{Z}_{2N_f, A} \text{ transformation}} Z \cdot \exp\left(\frac{-iN}{2\pi} \int_{M_4} (B_{cf}^{(2)} \wedge B_{cf}^{(2)})\right)$. The four phases can also be matched and cancelled by the anomaly (1.14), $Z \xrightarrow{\mathbb{Z}_{2N_f, A} \text{ transformation}} Z \cdot \exp\left(\frac{-iN}{2\pi} \int_{M_4} (B_c^{(1)} \wedge B_f^{(2)})\right)$ introduced in Ref. [38]. We use the triple data $(\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_c\text{-shift}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)})$ to label the discrete 0-form axial symmetry $\mathbb{Z}_{2N_f, A}$, a dimensionally reduced color-shift $\mathbb{Z}_{N_c\text{-shift}}^{(0)}$ symmetry introduced in Ref. [38], and the 1-form mixed chiral-flavor-locked center symmetry $\mathbb{Z}_{N_{cf}}^{(1)}$ that we introduce.

where the upper index (1) in $\mathbb{Z}_{N_{cf}}^{(1)}$ indicates it is truly a 1-form symmetry in (1.12), while the upper index (0) in $\mathbb{Z}_{N_c\text{-shift}}^{(0)}$ indicates that it is dimensionally reduced from a 1-form symmetry $\mathbb{Z}_{N_c\text{-shift}, [1]}$ to a 0-form symmetry. The 1-form color-shift symmetry $\mathbb{Z}_{N_c\text{-shift}, [1]}$ is introduced in Ref. [38].

Let us indicate how the higher 't Hooft anomalies in (1.12) and in (1.14) can be matched by breaking some of the global symmetries in the triple data $(\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_{cf}}^{(1)}, \mathbb{Z}_{N_c\text{-shift}}^{(0)})$. Our notations are that if the G_{sym} is broken, we denote it as G_{sym} ; if the G_{sym} is preserved, we would either indicate G_{sym} remained, or simply omit the symbol as we did in the Fig. 3.

1. QGP (quark-gluon plasma/liquid) at high T:

$$(\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_c\text{-shift}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)}) \xrightarrow{\text{SSB}} (\mathbb{Z}_{2N_f, A}; \cancel{\mathbb{Z}_{N_c\text{-shift}}^{(0)}}, \cancel{\mathbb{Z}_{N_{cf}}^{(1)}})$$

shown in Fig. 3 QGP. (1.17)

2. ChSB (chiral symmetry breaking) at low T and at lower densities and low μ :

$$(\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_c\text{-shift}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)}) \xrightarrow{\text{SSB}} (\cancel{\mathbb{Z}_{2N_f, A}} \rightarrow \mathbb{Z}_2^F; \mathbb{Z}_{N_c\text{-shift}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)})$$

shown in Fig. 3 ChSB. (1.18)

3. 2SC (2-color superconductivity) at low T and at intermediate densities and μ :

$$(\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_c\text{-shift}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)}) \xrightarrow{\text{SSB}} (\cancel{\mathbb{Z}_{2N_f, A}}; \cancel{\mathbb{Z}_{N_c\text{-shift}}^{(0)}}, \cancel{\mathbb{Z}_{N_{cf}}^{(1)}})$$

shown in Fig. 3 2SC. (1.19)

4. CFL (3-color-flavor locking superconductivity) at low T and at high density and high μ :

$$(\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_{c\text{-shift}}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)}) \xrightarrow{\text{SSB}} (\mathbb{Z}_{2N_f, A}; \mathbb{Z}_{N_{c\text{-shift}}}^{(0)}, \mathbb{Z}_{N_{cf}}^{(1)})$$

(1.20)

What we have shown above is that higher 't Hooft anomalies in (1.12) and in (1.14) can indeed be matched by four phases via breaking some of the global symmetries. We thus can constrain other possible QCD phases via the proposed approximate anomaly (1.12), based on higher 't Hooft anomaly matching and cancellation.

1.3. Mathematical primer

In this article, we use spectral sequences (Adams spectral sequence, Atiyah-Hirzebruch spectral sequence, and Serre spectral sequence) to compute several cobordism groups which appear in QCD matter phases (QGP, ChSB, 2SC and CFL in Sec. 1.1). See [4,29,52] for a primer.

We aim to compute the cobordism group Ω_d^G for $d \leq 5$ where G is the gauge group of QCD matter phases (QGP, ChSB, 2SC and CFL).

By the generalized Pontryagin-Thom isomorphism,

$$\Omega_d^G = \pi_d(MTG) \tag{1.21}$$

which identifies the cobordism group Ω_d^G with the homotopy group of the Madsen-Tillmann spectrum MTG .

We have the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}_p}^{s,t}(H^*(Y, \mathbb{Z}_p), \mathbb{Z}_p) \Rightarrow \pi_{t-s}(Y)^\wedge_p. \tag{1.22}$$

Here \mathcal{A}_p is the mod p Steenrod algebra, Y is any spectrum. For any finitely generated abelian group G , $G_p^\wedge = \lim_{n \rightarrow \infty} G/p^n G$ is the p -completion of G . In particular, \mathcal{A}_2 is generated by Steenrod squares Sq^i .

In our cases, the cobordism groups only have 2-torsion and 3-torsion.

We will use Adams spectral sequence to compute the 2-torsion part of the cobordism groups (we consider $Y = MTG$ in Adams spectral sequence, and we focus on $p = 2$).

We will use Serre spectral sequence and Atiyah-Hirzebruch spectral sequence to compute the 3-torsion part of the cobordism groups. In our cases, in order to compute the 3-torsion part of the cobordism groups, we need only to compute the cobordism group $\Omega_d^{\text{SO}}(\text{BG}')$ for some group G' .

We have the Atiyah-Hirzebruch spectral sequence

$$H_p(\text{BG}', \Omega_q^{\text{SO}}) \Rightarrow \Omega_{p+q}^{\text{SO}}(\text{BG}'). \tag{1.23}$$

Since

$$\Omega_d^{\text{SO}} = \begin{cases} \mathbb{Z} & d = 0 \\ 0 & d = 1 \\ 0 & d = 2 \\ 0 & d = 3 \\ \mathbb{Z} & d = 4 \\ \mathbb{Z}_2 & d = 5 \end{cases}, \tag{1.24}$$

we need to know the integral homology groups $H_p(BG', \mathbb{Z})$. In order to obtain this data, we compute the integral cohomology groups $H^p(BG', \mathbb{Z})$ using Serre spectral sequence (we find a fibration of which BG' is the total space).

For example, if $MTG = MSpin \wedge X$ where X is any spectrum, by Corollary 5.1.2 of [53], we have

$$\text{Ext}_{\mathcal{A}_2}^{s,t}(\mathbb{H}^*(MSpin \wedge X, \mathbb{Z}_2), \mathbb{Z}_2) = \text{Ext}_{\mathcal{A}_2(1)}^{s,t}(\mathbb{H}^*(X, \mathbb{Z}_2), \mathbb{Z}_2) \tag{1.25}$$

for $t - s < 8$. Here $\mathcal{A}_2(1)$ is the subalgebra of \mathcal{A}_2 generated by Sq^1 and Sq^2 .

So for the dimension $d = t - s < 8$, we have

$$\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(\mathbb{H}^*(X, \mathbb{Z}_2), \mathbb{Z}_2) \Rightarrow (\Omega_{t-s}^G)^{\wedge}_2. \tag{1.26}$$

The $\mathbb{H}^*(X, \mathbb{Z}_2)$ is an $\mathcal{A}_2(1)$ -module whose internal degree t is given by the $*$.

Our computation of E_2 pages of $\mathcal{A}_2(1)$ -modules is based on Lemma 11 of [29]. More precisely, we find a short exact sequence of $\mathcal{A}_2(1)$ -modules $0 \rightarrow L_1 \rightarrow L_2 \rightarrow L_3 \rightarrow 0$, then apply Lemma 11 of [29] to compute $\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(L_2, \mathbb{Z}_2)$ by the data of $\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(L_1, \mathbb{Z}_2)$ and $\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(L_3, \mathbb{Z}_2)$. Our strategy is choosing L_1 to be the direct sum of suspensions of \mathbb{Z}_2 on which Sq^1 and Sq^2 act trivially, then we take L_3 to be the quotient of L_2 by L_1 . We can use this procedure again and again until $\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(L_3, \mathbb{Z}_2)$ is determined.

2. QCD symmetries, anomalies and topological terms without time-reversal

Before we consider the cobordism theory and co/bordism group of the following four QCD matter phases, we need to convert our notations to involve both the internal symmetry and the spacetime symmetry. Here are the results of conversions for the high T QGP (quark-gluon plasma/liquid), the low T ChSB (chiral symmetry breaking), 2SC (2-color superconductivity) and CFL (3-color-flavor locking superconductivity) at high density.

QGP:

$$\begin{aligned} & \frac{[\text{SU}(N_c)_V] \times \text{SU}(N_f)_L \times \text{SU}(N_f)_R \times \text{U}(1)_V}{\mathbb{Z}_{N_c, V} \times \mathbb{Z}_{N_f, V}} \\ & := \left(\text{Spin}(d) \times_{\mathbb{Z}_2} \frac{\text{SU}(3)_L \times \text{SU}(3)_R \times \text{U}(1)_V}{(\mathbb{Z}_3)^2} \right) \\ & = \left(\text{Spin}(d) \times_{\mathbb{Z}_2} \frac{\text{U}(3)_L \times \text{U}(3)_R}{(\mathbb{Z}_3)^2 \times \text{U}(1)_A} \right) \times \mathbb{Z}_{3, L} \times \mathbb{Z}_{3, R} \\ & = \left(\text{Spin}(d) \times_{\mathbb{Z}_2} \frac{\text{U}(3)_L \times \text{U}(3)_R}{(\mathbb{Z}_{3, V} \times \mathbb{Z}_{3, A} \times \text{U}(1)_A)} \right). \tag{2.1} \end{aligned}$$

Here $\text{U}(1)_V = \frac{\text{U}(1)_L \times \text{U}(1)_R}{\text{U}(1)_A}$, $\text{U}(3)_L = \frac{\text{SU}(3)_L \times \text{U}(1)_L}{\mathbb{Z}_{3, L}}$, $\text{U}(3)_R = \frac{\text{SU}(3)_R \times \text{U}(1)_R}{\mathbb{Z}_{3, R}}$ where L/R for

$P_{L/R} = \frac{1 \pm \gamma^5}{2}$ ChSB:

$$\begin{aligned} & \frac{([\text{SU}(N_c)_V] \times \text{SU}(N_f)_V \times \text{U}(1)_V)}{\mathbb{Z}_{N_c, V} \times \mathbb{Z}_{N_f, V}} \\ & := \left(\text{Spin}(d) \times_{\mathbb{Z}_2} \frac{\text{SU}(3) \times \text{U}(1)}{(\mathbb{Z}_3)^2} \right) = \left(\text{Spin}(d) \times_{\mathbb{Z}_2} \frac{\text{U}(3)}{\mathbb{Z}_3} \right). \tag{2.2} \end{aligned}$$

2SC:

$$\frac{[\mathrm{SU}(2_c)_{V,rg}] \times \mathrm{SU}(2_f)_{L,ud} \times \mathrm{SU}(2_f)_{R,ud} \times \mathrm{U}(1_f)_{V,s} \times \mathrm{U}(1_c)_{V,b}}{\mathbb{Z}_{2,V}^F} := \left(\mathrm{Spin}(d) \times_{\mathbb{Z}_2} (\mathrm{SU}(2) \times \mathrm{SU}(2)) \right) \times (\mathrm{U}(1) \times \mathrm{U}(1)). \quad (2.3)$$

CFL:

$$\mathrm{SU}(3)_{C+L+R} \times \mathbb{Z}_{2,V}^F := \mathrm{Spin}(d) \times \mathrm{SU}(3)_{C+L+R}. \quad (2.4)$$

2.1. Chiral symmetry breaking $\frac{([\mathrm{SU}(3)_V] \times \mathrm{SU}(3)_V \times \mathrm{U}(1)_V)}{\mathbb{Z}_{3c,V} \times \mathbb{Z}_{3f,V}}$ as $(\mathrm{Spin}(d) \times_{\mathbb{Z}_2} \frac{\mathrm{U}(3)}{\mathbb{Z}_3})$

For ChSB with a global symmetry:

$$\frac{([\mathrm{SU}(N_c)_V] \times \mathrm{SU}(N_f)_V \times \mathrm{U}(1)_V)}{\mathbb{Z}_{N_c,V} \times \mathbb{Z}_{N_f,V}} := \left(\mathrm{Spin}(d) \times_{\mathbb{Z}_2} \frac{\mathrm{SU}(3) \times \mathrm{U}(1)}{(\mathbb{Z}_3)^2} \right) = \left(\mathrm{Spin}(d) \times_{\mathbb{Z}_2} \frac{\mathrm{U}(3)}{\mathbb{Z}_3} \right).$$

We have a fibration

$$\begin{array}{ccc} \mathrm{BU}(3) & & (2.5) \\ \downarrow & & \\ \mathrm{B}\left(\frac{\mathrm{U}(3)}{\mathbb{Z}_3}\right) & \longrightarrow & \mathrm{B}^2\mathbb{Z}_3. \end{array}$$

Hence we have the Serre spectral sequence, see Fig. 4.

$$\mathrm{H}^p(\mathrm{B}^2\mathbb{Z}_3, \mathrm{H}^q(\mathrm{BU}(3), \mathbb{Z})) \Rightarrow \mathrm{H}^{p+q}\left(\mathrm{B}\left(\frac{\mathrm{U}(3)}{\mathbb{Z}_3}\right), \mathbb{Z}\right). \quad (2.6)$$

We have

$$\mathrm{H}^*(\mathrm{BU}(3), \mathbb{Z}) = \mathbb{Z}[c_1, c_2, c_3] \quad (2.7)$$

and

$$\mathrm{H}^p(\mathrm{B}^2\mathbb{Z}_3, \mathbb{Z}) = \begin{cases} \mathbb{Z} & p = 0 \\ 0 & p = 1 \\ 0 & p = 2 \\ \mathbb{Z}_3 & p = 3 \\ 0 & p = 4 \\ \mathbb{Z}_3 & p = 5 \\ 0 & p = 6 \\ \mathbb{Z}_9 & p = 7 \end{cases}. \quad (2.8)$$

There is another approach: we have a fibration

$$\begin{array}{ccc} \mathrm{BPSU}(3) & & (2.9) \\ \downarrow & & \\ \mathrm{B}\left(\frac{\mathrm{U}(3)}{\mathbb{Z}_3}\right) & \longrightarrow & \mathrm{BU}(1). \end{array}$$

1	6	\mathbb{Z}^3	0	0	\mathbb{Z}_3^3	0	\mathbb{Z}_3^3	0	\mathbb{Z}_9^3	1
2										2
3										3
4	5	0	0	0	0	0	0	0	0	4
5										5
6										6
7										7
8	4	\mathbb{Z}^2	0	0	\mathbb{Z}_3^2	0	\mathbb{Z}_3^2	0	\mathbb{Z}_9^2	8
9										9
10										10
11	3	0	0	0	0	0	0	0	0	11
12										12
13										13
14										14
15	2	\mathbb{Z}	0	0	\mathbb{Z}_3	0	\mathbb{Z}_3	0	\mathbb{Z}_9	15
16										16
17										17
18	1	0	0	0	0	0	0	0	0	18
19										19
20										20
21										21
22	0	\mathbb{Z}	0	0	\mathbb{Z}_3	0	\mathbb{Z}_3	0	\mathbb{Z}_9	22
23										23
24		0	1	2	3	4	5	6	7	24
25										25

Fig. 4. Serre spectral sequence for the fibration $BU(3) \rightarrow B(\frac{U(3)}{\mathbb{Z}_3}) \rightarrow B^2\mathbb{Z}_3$. The arrow from (0,4) to (5,0) is a nontrivial differential by comparison with the Serre spectral sequence of the fibration $BSU(3) \rightarrow BPSU(3) \rightarrow B^2\mathbb{Z}_3$. There is no differential from (0,2) to (3,0) since the $3d \mathbb{Z}_3$ survives the spectral sequence in Fig. 5.

Hence we have the Serre spectral sequence, see Fig. 5.

$$H^p(BU(1), H^q(BPSU(3), \mathbb{Z})) \Rightarrow H^{p+q}(B(\frac{U(3)}{\mathbb{Z}_3}), \mathbb{Z}). \tag{2.10}$$

We have

$$H^*(BU(1), \mathbb{Z}) = \mathbb{Z}[c_1] \tag{2.11}$$

and

$$H^p(BPSU(3), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p = 0 \\ 0 & p = 1 \\ 0 & p = 2 \\ \mathbb{Z}_3 & p = 3 \\ \mathbb{Z} & p = 4 \\ 0 & p = 5 \\ \mathbb{Z} & p = 6 \end{cases}. \tag{2.12}$$

In Fig. 5, we find that the $3d \mathbb{Z}_3$ survives the spectral sequence, so in Fig. 4, there is no nontrivial differential from (0,2) to (3,0). Since the differential is a derivation, we conclude that




Fig. 5. Serre spectral sequence for the fibration $BPSU(3) \rightarrow B(\frac{U(3)}{\mathbb{Z}_3}) \rightarrow BU(1)$.

in Fig. 4 the dashed arrow from (0,4) to (3,2) does not actually exist. So there is a \mathbb{Z}_3 in 5d survives the spectral sequence, thus in Fig. 5, the dashed arrow from (0,4) to (2,3) also does not actually exist.

So

$$H^p(B(\frac{U(3)}{\mathbb{Z}_3}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p=0 \\ 0 & p=1 \\ \mathbb{Z} & p=2 \\ \mathbb{Z}_3 & p=3 \\ \mathbb{Z}^2 & p=4 \\ \mathbb{Z}_3 & p=5 \\ \mathbb{Z}^3 & p=6 \end{cases} \quad (2.13)$$

and

$$H_p(B(\frac{U(3)}{\mathbb{Z}_3}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p=0 \\ 0 & p=1 \\ \mathbb{Z} \times \mathbb{Z}_3 & p=2 \\ 0 & p=3 \\ \mathbb{Z}^2 \times \mathbb{Z}_3 & p=4 \\ 0 & p=5 \\ \mathbb{Z}^3 \times ? & p=6 \end{cases} \quad (2.14)$$

1	5	\mathbb{Z}_2							
2									
3									
4	4	\mathbb{Z}	0	$\mathbb{Z} \times \mathbb{Z}_3$					
5									
6									
7									
8	3	0	0	0	0				
9									
10									
11	2	0	0	0	0	0			
12									
13									
14									
15	1	0	0	0	0	0	0		
16									
17									
18	0	\mathbb{Z}	0	$\mathbb{Z} \times \mathbb{Z}_3$	0	$\mathbb{Z}^2 \times \mathbb{Z}_3$	0	$\mathbb{Z}^3 \times ?$	
19									
20									
21		0	1	2	3	4	5	6	

Fig. 6. Atiyah-Hirzebruch spectral sequence for $\Omega_d^{SO \times \frac{U(3)}{\mathbb{Z}_3}}$. Here ? is an undetermined 3-torsion group.

Here ? is an undetermined 3-torsion group.

By the Atiyah-Hirzebruch spectral sequence, we have

$$H_p(B(\frac{U(3)}{\mathbb{Z}_3}), \Omega_q^{SO}) \Rightarrow \Omega_{p+q}^{SO}(B(\frac{U(3)}{\mathbb{Z}_3})). \tag{2.15}$$

See Fig. 6.

Since the localization of $\text{Spin} \times_{\mathbb{Z}_2} \frac{U(3)}{\mathbb{Z}_3}$ and $\text{SO} \times \frac{U(3)}{\mathbb{Z}_3}$ at the prime 3 are the same, the 3-torsion part of $\Omega_d^{\text{Spin} \times_{\mathbb{Z}_2} \frac{U(3)}{\mathbb{Z}_3}}$ and $\Omega_d^{\text{SO} \times \frac{U(3)}{\mathbb{Z}_3}}$ are the same.

Since the localization of $\text{Spin} \times_{\mathbb{Z}_2} \frac{U(3)}{\mathbb{Z}_3}$ and $\text{Spin} \times_{\mathbb{Z}_2} U(3)$ at the prime 2 are the same, the 2-torsion part of $\Omega_d^{\text{Spin} \times_{\mathbb{Z}_2} \frac{U(3)}{\mathbb{Z}_3}}$ and $\Omega_d^{\text{Spin} \times_{\mathbb{Z}_2} U(3)}$ are the same.

Also since $U(3) = \text{SU}(3) \times_{\mathbb{Z}_3} U(1)$, the localization of $\text{Spin} \times_{\mathbb{Z}_2} U(3)$ and $\text{Spin} \times_{\mathbb{Z}_2} (U(1) \times \text{SU}(3))$ at the prime 2 are the same, the 2-torsion part of $\Omega_d^{\text{Spin} \times_{\mathbb{Z}_2} U(3)}$ and $\Omega_d^{\text{Spin} \times_{\mathbb{Z}_2} (U(1) \times \text{SU}(3))} = \Omega_d^{\text{Spin}^c \times \text{SU}(3)}$ are the same.

We have $MT(\text{Spin}^c \times \text{SU}(3)) = M\text{Spin} \wedge \Sigma^{-2} MU(1) \wedge (\text{BSU}(3))_+$.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^*(\text{BSU}(3), \mathbb{Z}_2), \mathbb{Z}_2) \Rightarrow \Omega_{t-s}^{\text{Spin}^c \times \text{SU}(3)}. \tag{2.16}$$

We have

$$H^*(\text{BSU}(3), \mathbb{Z}_2) = \mathbb{Z}_2[c_2, c_3] \tag{2.17}$$




Fig. 7. The $\mathcal{A}_2(1)$ -module structure of $H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 6.




Fig. 8. $\Omega_{2*}^{\text{Spin}^c \times \text{SU}(3)}$.

where c_i is the Chern class of the $SU(3)$ bundle.

By Thom isomorphism,

$$H^{*+2}(MU(1), \mathbb{Z}_2) = \mathbb{Z}_2[c_1]U \tag{2.18}$$

where c_1 is the Chern class of the $U(1)$ bundle and U is the Thom class.

The $\mathcal{A}_2(1)$ -module structure of $H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 6 and the E_2 page are shown in Fig. 7, 8.

So there is no 2-torsion in $\Omega_d^{\text{Spin} \times \mathbb{Z}_2 \times \frac{U(3)}{\mathbb{Z}_3}}$.

Combine the 2-torsion and 3-torsion results, we have in Table 3.

2.2. 3-Color-flavor locking superconductivity $SU(3)_{C+L+R} \times \mathbb{Z}_{2,V}^F$ as $\text{Spin} \times \text{SU}(3)$

We have $MT(\text{Spin} \times \text{SU}(3)) = M\text{Spin} \wedge (\text{BSU}(3))_+$.

Table 3

Bordism group. The notation $e(A, B)$ denotes a group extension of A by B , that is, a group that fits into the following short exact sequence $0 \rightarrow B \rightarrow e(A, B) \rightarrow A \rightarrow 0$.

d	Bordism group	
	$\Omega_d^{\text{Spin} \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}}$	generators
0	\mathbb{Z}	
1	0	
2	$\mathbb{Z} \times \mathbb{Z}_3$	
3	0	
4	$e(\mathbb{Z}^2 \times \mathbb{Z}_3, \mathbb{Z})$	
5	0	



Fig. 9. The $\mathcal{A}_2(1)$ -module structure of $H^*(BSU(3), \mathbb{Z}_2)$ below degree 6.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(H^*(BSU(3), \mathbb{Z}_2), \mathbb{Z}_2) \Rightarrow \Omega_{t-s}^{\text{Spin} \times \text{SU}(3)}. \tag{2.19}$$

We have

$$H^*(BSU(3), \mathbb{Z}_2) = \mathbb{Z}_2[c_2, c_3] \tag{2.20}$$

where c_i is the Chern class of the $SU(3)$ bundle (Table 4).

The $\mathcal{A}_2(1)$ -module structure of $H^*(BSU(3), \mathbb{Z}_2)$ below degree 6 and the E_2 page are shown in Fig. 9, 10.

2.3. 2-Color superconductivity: $\frac{[SU(2_c)_{V,rg}] \times SU(2_f)_{L,ud} \times SU(2_f)_{R,ud} \times U(1_f)_{V,s} \times U(1_c)_{V,b}}{\mathbb{Z}_{2,V}^F}$ as

$$\text{Spin} \times_{\mathbb{Z}_2} \text{Spin}(4) \times U(1) \times U(1)$$

We have $MT(\text{Spin} \times_{\mathbb{Z}_2} \text{Spin}(4) \times U(1) \times U(1)) = M\text{Spin} \wedge \Sigma^{-4}M\text{SO}(4) \wedge (\text{BU}(1) \times \text{BU}(1))_+$.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\begin{aligned} \text{Ext}_{\mathcal{A}_2(1)}^{s,t}(H^{*+4}(M\text{SO}(4), \mathbb{Z}_2) \otimes H^*(\text{BU}(1), \mathbb{Z}_2) \otimes H^*(\text{BU}(1), \mathbb{Z}_2), \mathbb{Z}_2) \\ \Rightarrow \Omega_{t-s}^{\text{Spin} \times \mathbb{Z}_2 \text{Spin}(4) \times U(1) \times U(1)}. \end{aligned} \tag{2.21}$$

By Thom isomorphism, we have




Fig. 10. $\Omega_*^{\text{Spin} \times \text{SU}(3)}$.

Table 4

Bordism group. $\tilde{\eta}$ is a mod 2 index of 1d Dirac operator. Arf is a 2d Arf invariant. σ is the signature of manifold. Note that $c_3 = \text{Sq}^2 c_2 = (w_2 + w_1^2)c_2 = 0 \pmod 2$ on Spin 6-manifolds.

d	Bordism group	
	$\Omega_d^{\text{Spin} \times \text{SU}(3)}$	generators
0	\mathbb{Z}	
1	\mathbb{Z}_2	$\tilde{\eta}$
2	\mathbb{Z}_2	Arf
3	0	
4	\mathbb{Z}^2	$\frac{\sigma}{16}, c_2$
5	0	
6	\mathbb{Z}	$\frac{c_3}{2}$

$$H^{*+4}(MSO(4), \mathbb{Z}_2) = \mathbb{Z}_2[w'_2, w'_3, w'_4]U \tag{2.22}$$

where w'_i is the Stiefel-Whitney class of the $SO(4)$ bundle and U is the Thom class.

We also have

$$H^*(BU(1), \mathbb{Z}_2) = \mathbb{Z}_2[c_1] \tag{2.23}$$

where c_1 is the first Chern class of the $U(1)$ bundle (Table 5).

The $\mathcal{A}_2(1)$ -module structure of $H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2)$ below degree 5 and the E_2 page are shown in Fig. 11, 12.




Fig. 11. The $\mathcal{A}_2(1)$ -module structure of $H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2)$ below degree 5.

Table 5

Bordism group. Here e_i is the Euler class, p_i is the Pontryagin class, c_i is the Chern class. Here $w'_i = w_i(SO(4))$, $p'_1 = p_1(SO(4))$, $e'_i = e_i(SO(4))$. $\tilde{\eta}$ is the mod 2 index of 1d Dirac operator, Arf is the Arf invariant. The ? is an undetermined cobordism invariant which also appears in [52].

d	Bordism group	
	$\Omega_d^{\text{Spin} \times \mathbb{Z}_2 \text{ Spin}(4) \times \text{U}(1) \times \text{U}(1)}$	generators
0	\mathbb{Z}	
1	0	
2	\mathbb{Z}^2	c_1, c'_1
3	0	
4	\mathbb{Z}^6	$p'_1, e'_4, c_1^2, c_1'^2, c_1 c_1', ?$
5	\mathbb{Z}_2^3	$w'_2 w'_3, w'_4 \tilde{\eta}, w'_3 \text{Arf}(?)$




Fig. 12. $\Omega_*^{\text{Spin} \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)}$

2.4. Quark gluon plasma/liquid $\frac{[\text{SU}(3)_V] \times \text{SU}(3)_L \times \text{SU}(3)_R \times \text{U}(1)_V}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V}}$ as $\text{Spin}(d) \times \mathbb{Z}_2 \frac{\text{U}(3)_L \times \text{U}(3)_R}{(\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,A} \times \text{U}(1)_A)}$

Since the localization of $\text{Spin} \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1)}$ and $\text{SO} \times \frac{\text{U}(3) \times \text{U}(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1)}$ at the prime 3 are the same, the 3-torsion of $\Omega_d^{\text{Spin} \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1)}}$ and $\Omega_d^{\text{SO} \times \frac{\text{U}(3) \times \text{U}(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1)}}$ are the same.

Since the localization of $\text{Spin} \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1)}$ and $\text{Spin} \times \mathbb{Z}_2 (\text{U}(1) \times \text{SU}(3) \times \text{SU}(3))$ at the prime 2 are the same, the 2-torsion of $\Omega_d^{\text{Spin} \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1)}}$ and $\Omega_d^{\text{Spin}^c \times \text{SU}(3) \times \text{SU}(3)}$ are the same.

We have $MT(\text{Spin}^c \times \text{SU}(3) \times \text{SU}(3)) = M\text{Spin} \wedge \Sigma^{-2} M\text{U}(1) \wedge (\text{BSU}(3) \times \text{BSU}(3))_+$.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\begin{aligned} \text{Ext}_{\mathcal{A}_2(1)}^{s,t}(\mathbb{H}^{*+2}(M\text{U}(1), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2), \mathbb{Z}_2) \\ \Rightarrow \Omega_{t-s}^{\text{Spin}^c \times \text{SU}(3) \times \text{SU}(3)}. \end{aligned} \tag{2.24}$$

We have

$$\mathbb{H}^*(\text{BSU}(3) \times \text{BSU}(3), \mathbb{Z}_2) = \mathbb{Z}_2[c_2, c_3, c'_2, c'_3] \tag{2.25}$$

where c_i is the Chern class of the $\text{SU}(3)$ bundle.

By Thom isomorphism,

$$\mathbb{H}^{*+2}(M\text{U}(1), \mathbb{Z}_2) = \mathbb{Z}_2[c_1]U \tag{2.26}$$

where c_1 is the Chern class of the $\text{U}(1)$ bundle and U is the Thom class.

The $\mathcal{A}_2(1)$ -module structure of $\mathbb{H}^{*+2}(M\text{U}(1), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2)$ below degree 6 and the E_2 page are shown in Fig. 13, 14.




Fig. 13. The $\mathcal{A}_2(1)$ -module structure of $H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 6.




Fig. 14. $\Omega_*^{Spin^c \times SU(3) \times SU(3)}$.

So there is no 2-torsion in $\Omega_d^{Spin \times \mathbb{Z}_2 \times \frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}}$.

We have a fibration

$$BU(3) \times BU(3) \tag{2.27}$$



$$B\left(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}\right) \longrightarrow B^2\mathbb{Z}_3 \times B^2\mathbb{Z}_3 \times B^2U(1).$$

Hence we have the Serre spectral sequence, see Fig. 15.

$$\begin{aligned} & H^p(B^2\mathbb{Z}_3 \times B^2\mathbb{Z}_3 \times B^2U(1), H^q(BU(3) \times BU(3), \mathbb{Z})) \\ & \Rightarrow H^{p+q}\left(B\left(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}\right), \mathbb{Z}\right). \end{aligned} \tag{2.28}$$

We have

$$H^*(BU(3) \times BU(3), \mathbb{Z}) = \mathbb{Z}[c_1, c_2, c_3, c'_1, c'_2, c'_3] \tag{2.29}$$




Fig. 15. Serre spectral sequence for the fibration $BU(3) \times BU(3) \rightarrow B(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}) \rightarrow B^2\mathbb{Z}_3 \times B^2\mathbb{Z}_3 \times B^2U(1)$. The dashed arrows are possible differentials.

and

$$H^p(B^2\mathbb{Z}_3, \mathbb{Z}) = \begin{cases} \mathbb{Z} & p=0 \\ 0 & p=1 \\ 0 & p=2 \\ \mathbb{Z}_3 & p=3 \\ 0 & p=4 \\ \mathbb{Z}_3 & p=5 \\ 0 & p=6 \\ \mathbb{Z}_9 & p=7 \end{cases} \quad (2.30)$$

and

$$H^p(B^2U(1), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p=0 \\ 0 & p=1 \\ 0 & p=2 \\ \mathbb{Z} & p=3 \\ 0 & p=4 \\ 0 & p=5 \\ \mathbb{Z}_2 & p=6 \end{cases} \quad (2.31)$$

By Künneth formula,

$$H^p(B^2\mathbb{Z}_3 \times B^2\mathbb{Z}_3 \times B^2U(1), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p = 0 \\ 0 & p = 1 \\ 0 & p = 2 \\ \mathbb{Z} \times \mathbb{Z}_3^2 & p = 3 \\ 0 & p = 4 \\ \mathbb{Z}_3^3 & p = 5 \\ \mathbb{Z}_2 \times \mathbb{Z}_3^3 & p = 6 \end{cases} \quad (2.32)$$

Compared with the 2-torsion result, we find that the differential from (0,2) to (3,0) kills one \mathbb{Z} since the 2d cobordism group contains only one \mathbb{Z} , and the differential from (0,4) to (3,2) kills two \mathbb{Z} since the 4d cobordism group contains four \mathbb{Z} while one \mathbb{Z} is from Ω_4^{SO} .

So

$$H^p(B(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p = 0 \\ 0 & p = 1 \\ \mathbb{Z} & p = 2 \\ \mathbb{Z}_3 \text{ or } \mathbb{Z}_3^2 & p = 3 \\ \mathbb{Z}_3^3 & p = 4 \\ ? & p = 5 \end{cases} \quad (2.33)$$

and

$$H_p(B(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & p = 0 \\ 0 & p = 1 \\ \mathbb{Z} \times \mathbb{Z}_3 \text{ or } \mathbb{Z} \times \mathbb{Z}_3^2 & p = 2 \\ 0 & p = 3 \\ \mathbb{Z}_3^3 \times ? & p = 4 \\ 0 & p = 5 \end{cases} \quad (2.34)$$

Here ? is an undetermined 3-torsion group.

By the Atiyah-Hirzebruch spectral sequence, we have

$$H_p(B(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}), \Omega_q^{SO}) \Rightarrow \Omega_{p+q}^{SO}(B(\frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)})). \quad (2.35)$$

See Fig. 16.

Combine the 2-torsion and 3-torsion results, we have in Table 6.

3. QCD symmetries, anomalies and topological terms with time-reversal

Now we consider putting the QCD matters on the smooth differentiable and unorientable spacetime manifolds – if the fermions/spinor can live on them, we require Spin structure; if we require time-reversal $T = CP$, or CT or other reflection symmetries, we require Pin^+ , Pin^- or other semi-direct (\rtimes) product or twisted structures between the spacetime tangent bundle TM and the gauge bundle E_G of the gauge group G . See more in the main text and see an overview of our setting in [4]. Follow Fig. 1 and Fig. 2, we can choose any suitable outer automorphism of the color gauge or flavor global symmetry group as possible time-reversal symmetries, which can be any reasonable \mathbb{Z}_2 -reflection symmetry. This implies putting the Euclidean QCD_4 path integral on an unorientable spacetime. The most general case is a semi-direct product $\rtimes \mathbb{Z}_4^T$, which is all allowed total group made from the exact sequence:

1	5	\mathbb{Z}_2						
2								
3								
4	4	\mathbb{Z}	0					
5								
6								
7								
8	3	0	0	0				
9								
10								
11	2	0	0	0	0			
12								
13								
14								
15	1	0	0	0	0	0		
16								
17								
18	0	\mathbb{Z}	0	$\mathbb{Z} \times \mathbb{Z}_3^n$	0	$\mathbb{Z}^3 \times ?$	0	
19								
20								
21			0	1	2	3	4	5
22								

Fig. 16. Atiyah-Hirzebruch spectral sequence for $\Omega_d^{SO \times \frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}}$. Here $n = 1$ or 2 , while $?$ is an undetermined 3-torsion group.

Table 6
Bordism group. The notation $e(A, B)$ denotes a group extension of A by B , that is, a group that fits into the following short exact sequence $0 \rightarrow B \rightarrow e(A, B) \rightarrow A \rightarrow 0$. Here $n = 1$ or 2 , while $?$ is an undetermined 3-torsion group.

d	Bordism group	
	$\Omega_d^{Spin \times \mathbb{Z}_2 \times \frac{U(3) \times U(3)}{\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1)}}$	generators
0	\mathbb{Z}	
1	0	
2	$\mathbb{Z} \times \mathbb{Z}_3^n$	
3	0	
4	$e(\mathbb{Z}^3 \times ?, \mathbb{Z})$	
5	0	

$$1 \rightarrow \mathbb{Z}_2^F \rightarrow \mathbb{Z}_4^T \rightarrow \mathbb{Z}_2^T \rightarrow 1.$$

Here we will only focus on two cases: The direct product $\times \mathbb{Z}_4^T$ which implies the Pin^+ structure, where $\mathbb{Z}_4^T \supset \mathbb{Z}_2^F$. We may also denote such a $\mathbb{Z}_4^T := \mathbb{Z}_4^{TF}$ to indicate it includes \mathbb{Z}_2^F as a normal subgroup. The direct product $\times \mathbb{Z}_2^T \times \mathbb{Z}_2^F$ which implies the Pin^- structure. For other possible time-reversal symmetries, we leave them in a future work [47].

Table 7

Bordism group. Here c_1 is the Chern class of the $U(1)$ bundle, c_2 is the Chern class of the $SU(3)$ bundle, ABK is the Arf-Brown-Kervaire invariant.

Bordism group			
d	$\Omega_d^{\text{Pin}^c \times \text{SU}(3)} = \Omega_d^{\text{Pin}^+ \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}} = \Omega_d^{\text{Pin}^- \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}}$		generators
0	\mathbb{Z}_2		
1	0		
2	\mathbb{Z}_4		$ABK \text{ mod } 4$
3	0		
4	$\mathbb{Z}_2^2 \times \mathbb{Z}_8$		$c_2 \text{ mod } 2, c_1^2 \text{ mod } 2, (c_1 \text{ mod } 2)ABK$
5	0		

3.1. Chiral symmetry breaking $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V) \times \mathbb{Z}_4^T}{\mathbb{Z}_{3c,V} \times \mathbb{Z}_{3f,V} \times \mathbb{Z}_{2,V}^F}$

3.1.1. $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V) \times \mathbb{Z}_4^T}{\mathbb{Z}_{3c,V} \times \mathbb{Z}_{3f,V} \times \mathbb{Z}_{2,V}^F}$ as $(\text{Pin}^+(d) \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3})$ and $\frac{([SU(3)_V] \times SU(3)_V \times U(1)_V) \times \mathbb{Z}_2^T}{\mathbb{Z}_{3c,V} \times \mathbb{Z}_{3f,V}}$ as $(\text{Pin}^-(d) \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3})$

Since the localization of $\text{Pin}^\pm \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}$ and $O \times \frac{U(3)}{\mathbb{Z}_3}$ at the prime 3 are the same, so the 3-torsion of $\Omega_d^{\text{Pin}^\pm \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}}$ and $\Omega_d^{O \times \frac{U(3)}{\mathbb{Z}_3}}$ are the same, hence there is no 3-torsion in $\Omega_d^{\text{Pin}^\pm \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}}$.

Since the localization of $\text{Pin}^\pm \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}$ and $\text{Pin}^\pm \times \mathbb{Z}_2 (U(1) \times SU(3))$ at the prime 2 are the same, so the 2-torsion of $\Omega_d^{\text{Pin}^\pm \times \mathbb{Z}_2 \frac{U(3)}{\mathbb{Z}_3}}$ and $\Omega_d^{\text{Pin}^\pm \times \text{SU}(3)}$ are the same.

We have $MT(\text{Pin}^c \times \text{SU}(3)) = M\text{Spin} \wedge \Sigma^{-2}MU(1) \wedge \Sigma^{-1}MO(1) \wedge (\text{BSU}(3))_+$.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\begin{aligned} \text{Ext}_{\mathcal{A}_2(1)}^{s,t}(\mathbb{H}^{*+2}(MU(1), \mathbb{Z}_2) \otimes \mathbb{H}^{*+1}(MO(1), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2), \mathbb{Z}_2) \\ \Rightarrow \Omega_{t-s}^{\text{Pin}^c \times \text{SU}(3)}. \end{aligned} \tag{3.1}$$

We have

$$\mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2) = \mathbb{Z}_2[c_2, c_3] \tag{3.2}$$

where c_i is the Chern class of the $SU(3)$ bundle.

By Thom isomorphism,

$$\mathbb{H}^{*+2}(MU(1), \mathbb{Z}_2) = \mathbb{Z}_2[e_1]U \tag{3.3}$$

where c_1 is the Chern class of the $U(1)$ bundle and U is the Thom class.

Also by Thom isomorphism,

$$\mathbb{H}^{*+1}(MO(1), \mathbb{Z}_2) = \mathbb{Z}_2[w_1]V \tag{3.4}$$

where w_1 is the Stiefel-Whitney class of the $O(1)$ bundle and V is the Thom class.

The $\mathcal{A}_2(1)$ -module structure of $\mathbb{H}^{*+2}(MU(1), \mathbb{Z}_2) \otimes \mathbb{H}^{*+1}(MO(1), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2)$ below degree 6 and the E_2 page are shown in Fig. 17, 18.

Combine the 2-torsion and 3-torsion results, we have in Table 7.




Fig. 17. The $\mathcal{A}_2(1)$ -module structure of $H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 6.




Fig. 18. $\Omega_*^{\text{Pin}^c \times \text{SU}(3)}$.

3.2. 3-Color-flavor locking superconductivity $SU(3)_{C+L+R} \times \mathbb{Z}_4^T$

3.2.1. $SU(3)_{C+L+R} \times \mathbb{Z}_4^{TF}$ as $\text{Pin}^+ \times \text{SU}(3)$

We have $MT(\text{Pin}^+ \times \text{SU}(3)) = MT\text{Pin}^+ \wedge (\text{BSU}(3))_+$.

$MT\text{Pin}^+ = M\text{Spin} \wedge \Sigma^1 MTO(1)$.

By Künneth formula,

$$H^*(\Sigma^1 MTO(1) \wedge (\text{BSU}(3))_+, \mathbb{Z}_2) = H^{*-1}(MTO(1), \mathbb{Z}_2) \otimes H^*(\text{BSU}(3), \mathbb{Z}_2). \quad (3.5)$$




Fig. 19. The $\mathcal{A}_2(1)$ -module structure of $H^{*-1}(MTO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 5.




Fig. 20. $\Omega_*^{Pin^+ \times SU(3)}$

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(H^{*-1}(MTO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2), \mathbb{Z}_2) \Rightarrow \Omega_{t-s}^{Pin^+ \times SU(3)}. \quad (3.6)$$

The $\mathcal{A}_2(1)$ -module structure of $H^{*-1}(MTO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 5 and the E_2 page are shown in Fig. 19, 20 (Table 8).

Table 8

Bordism group. Here w_1 is the Stiefel-Whitney class of the tangent bundle, $\tilde{\eta}$ is the mod 2 index of 1d Dirac operator, Arf is the Arf invariant, η is the 4d eta invariant, c_2 is the Chern class of the SU(3) bundle.

d	Bordism group	
	$\Omega_d^{\text{Pin}^+ \times \text{SU}(3)}$	generators
0	\mathbb{Z}_2	
1	0	
2	\mathbb{Z}_2	$w_1 \tilde{\eta}$
3	\mathbb{Z}_2	$w_1 \text{Arf}$
4	$\mathbb{Z}_2 \times \mathbb{Z}_{16}$	$c_2 \pmod{2}, \eta$
5	0	




Fig. 21. The $\mathcal{A}_2(1)$ -module structure of $H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 5.

3.2.2. $SU(3)_{C+L+R} \times \mathbb{Z}_{2,V}^F \times \mathbb{Z}_2^T$ as $\text{Pin}^- \times \text{SU}(3)$

We have $MT(\text{Pin}^- \times \text{SU}(3)) = MT\text{Pin}^- \wedge (BSU(3))_+$.

$MT\text{Pin}^- = M\text{Spin} \wedge \Sigma^{-1}MO(1)$.

By Künneth formula,

$$H^*(\Sigma^{-1}MO(1) \wedge (BSU(3))_+, \mathbb{Z}_2) = H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2). \tag{3.7}$$

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2), \mathbb{Z}_2) \Rightarrow \Omega_{t-s}^{\text{Pin}^- \times \text{SU}(3)}. \tag{3.8}$$

The $\mathcal{A}_2(1)$ -module structure of $H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 5 and the E_2 page are shown in Fig. 21, 22 (Table 9).




Fig. 22. $\Omega_*^{\text{Pin}^- \times \text{SU}(3)}$

Table 9

Bordism group. Here $\tilde{\eta}$ is the mod 2 index of 1d Dirac operator, ABK is the Arf-Brown-Kervaire invariant, c_2 is the Chern class of the $\text{SU}(3)$ bundle.

d	Bordism group	
	$\Omega_d^{\text{Pin}^- \times \text{SU}(3)}$	generators
0	\mathbb{Z}_2	
1	\mathbb{Z}_2	$\tilde{\eta}$
2	\mathbb{Z}_8	ABK
3	0	
4	\mathbb{Z}_2	$c_2 \pmod{2}$
5	0	

3.3. 2-Color superconductivity: $\frac{([\text{SU}(2_c)_{V,rg}] \times \text{SU}(2_f)_{L,ud} \times \text{SU}(2_f)_{R,ud} \times \text{U}(1_f)_{V,s} \times \text{U}(1_c)_{V,b}) \times \mathbb{Z}_4^T}{(\mathbb{Z}_{2,V}^F)^2}$

3.3.1. $\frac{([\text{SU}(2_c)_{V,rg}] \times \text{SU}(2_f)_{L,ud} \times \text{SU}(2_f)_{R,ud} \times \text{U}(1_f)_{V,s} \times \text{U}(1_c)_{V,b}) \times \mathbb{Z}_4^{TF}}{(\mathbb{Z}_{2,V}^F)^2}$

$as\text{Pin}^+ \times_{\mathbb{Z}_2} \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)$

We have $MT(\text{Pin}^+ \times_{\mathbb{Z}_2} \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)) = MTP\text{in}^+ \wedge \Sigma^{-4}MSO(4) \wedge (\text{BU}(1) \times \text{BU}(1))_+ = M\text{Spin} \wedge \Sigma^1 M\text{TO}(1) \wedge \Sigma^{-4}MSO(4) \wedge (\text{BU}(1) \times \text{BU}(1))_+$.

We have the constraint $w_2 = w'_2$ where w_i is the Stiefel-Whitney class of the tangent bundle, w'_i is the Stiefel-Whitney class of the $\text{SO}(4)$ bundle.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(\mathbb{H}^{*-1}(M\text{TO}(1), \mathbb{Z}_2) \otimes \mathbb{H}^{*+4}(MSO(4), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BU}(1), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BU}(1), \mathbb{Z}_2), \mathbb{Z}_2)$$

$$\Rightarrow \Omega_{t-s}^{\text{Pin}^+ \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)} \quad (3.9)$$

By Thom isomorphism, we have

$$H^{*+4}(MSO(4), \mathbb{Z}_2) = \mathbb{Z}_2[w'_2, w'_3, w'_4]U \quad (3.10)$$

where w'_i is the Stiefel-Whitney class of the SO(4) bundle and U is the Thom class.

Also by Thom isomorphism, we have

$$H^{*-1}(MTO(1), \mathbb{Z}_2) = \mathbb{Z}_2[w_1]V \quad (3.11)$$

where w_1 is the Stiefel-Whitney class of the O(1) bundle V_1 , and V is the Thom class of $-V_1$.

We also have

$$H^*(BU(1), \mathbb{Z}_2) = \mathbb{Z}_2[c_1] \quad (3.12)$$

where c_1 is the first Chern class of the U(1) bundle.

The $\mathcal{A}_2(1)$ -module structure of $H^{*-1}(MTO(1), \mathbb{Z}_2) \otimes H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2)$ below degree 5 and the E_2 page are shown in Fig. 23, 24 (Table 10).

$$3.3.2. \frac{([\text{Ext}_{\mathcal{A}_2(1)}^{s,t}(\text{SU}(2_c)_{V,rg}] \times \text{SU}(2_f)_{L,ud} \times \text{SU}(2_f)_{R,ud} \times \text{U}(1_f)_{V,s} \times \text{U}(1_c)_{V,b}) \times \mathbb{Z}_2^T}{\mathbb{Z}_2^F}_{2,V}$$

$as\text{Pin}^- \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)$

We have $MT(\text{Pin}^- \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)) = MT\text{Pin}^- \wedge \Sigma^{-4}MSO(4) \wedge (BU(1) \times BU(1))_+ = M\text{Spin} \wedge \Sigma^{-1}MO(1) \wedge \Sigma^{-4}MSO(4) \wedge (BU(1) \times BU(1))_+$.

We have the constraint $w_2 + w_1^2 = w'_2$ where w_i is the Stiefel-Whitney class of the tangent bundle, w'_i is the Stiefel-Whitney class of the SO(4) bundle.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\begin{aligned} & \text{Ext}_{\mathcal{A}_2(1)}^{s,t}(H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \\ & \quad \otimes H^*(BU(1), \mathbb{Z}_2), \mathbb{Z}_2) \\ & \Rightarrow \Omega_{t-s}^{\text{Pin}^- \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)} \end{aligned} \quad (3.13)$$

By Thom isomorphism, we have

$$H^{*+4}(MSO(4), \mathbb{Z}_2) = \mathbb{Z}_2[w'_2, w'_3, w'_4]U \quad (3.14)$$

where w'_i is the Stiefel-Whitney class of the SO(4) bundle and U is the Thom class.

Also by Thom isomorphism, we have

$$H^{*+1}(MO(1), \mathbb{Z}_2) = \mathbb{Z}_2[w_1]V \quad (3.15)$$

where w_1 is the Stiefel-Whitney class of the O(1) bundle V_1 , and V is the Thom class of V_1 .

We also have

$$H^*(BU(1), \mathbb{Z}_2) = \mathbb{Z}_2[c_1] \quad (3.16)$$

where c_1 is the first Chern class of the U(1) bundle.

The $\mathcal{A}_2(1)$ -module structure of $H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2)$ below degree 5 and the E_2 page are shown in Fig. 25, 26 (Table 11).




Fig. 23. The $\mathcal{A}_2(1)$ -module structure of $H^{*-1}(MTO(1), \mathbb{Z}_2) \otimes H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2)$ below degree 5.




Fig. 24. $\Omega_*^{\text{Pin}^+ \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)}$

Table 10

Bordism group. Here w_i is the Stiefel-Whitney class of the tangent bundle, w'_i is the Stiefel-Whitney class of the $\text{SO}(4)$ bundle, c_1 (c'_1) is the Chern class of the $\text{U}(1)$ bundle, $\eta_{\text{Spin}(4)}$ is similar to the $\eta_{\text{SU}(2)}$ defined in [4].

Bordism group		
d	$\Omega_d^{\text{Pin}^+ \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)}$	generators
0	\mathbb{Z}_2	
1	0	
2	\mathbb{Z}_2^3	$w_1^2, c_1 \text{ mod } 2, c'_1 \text{ mod } 2$
3	0	
4	$\mathbb{Z}_2^7 \times \mathbb{Z}_4$	$w_4, w_2'^2, w_1^2 c_1, w_1^2 c'_1, c_1^2 \text{ mod } 2, c_1'^2 \text{ mod } 2, c_1 c'_1 \text{ mod } 2, \eta_{\text{Spin}(4)}$
5	\mathbb{Z}_2	$w_2 w'_3$

3.4. Quark gluon plasma/liquid $\frac{([\text{SU}(3)_V] \times \text{SU}(3)_L \times \text{SU}(3)_R \times \text{U}(1)_V) \rtimes \mathbb{Z}_4^T}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V} \times \mathbb{Z}_{2,V}^T}$

3.4.1. $\frac{([\text{SU}(3)_V] \times \text{SU}(3)_L \times \text{SU}(3)_R \times \text{U}(1)_V) \times \mathbb{Z}_4^{TF}}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V} \times \mathbb{Z}_{2,V}^T}$ as $\text{Pin}^+ \times \mathbb{Z}_2 \frac{\text{U}(3)_L \times \text{U}(3)_R}{(\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,A} \times \text{U}(1)_A)}$ and $\frac{([\text{SU}(3)_V] \times \text{SU}(3)_L \times \text{SU}(3)_R \times \text{U}(1)_V) \times \mathbb{Z}_2^T}{\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,V}}$ as $\text{Pin}^- \times \mathbb{Z}_2 \frac{\text{U}(3)_L \times \text{U}(3)_R}{(\mathbb{Z}_{3,V} \times \mathbb{Z}_{3,A} \times \text{U}(1)_A)}$

Since the localization of $\text{Pin}^\pm \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}$ and $\text{O} \times \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}$ at the prime 3 are the same, the 3-torsion of $\Omega_d^{\text{Pin}^\pm \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}}$ and $\Omega_d^{\text{O} \times \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}}$ are the same, hence there is no 3-torsion in $\Omega_d^{\text{Pin}^\pm \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}}$.




Fig. 25. The $\mathcal{A}_2(1)$ -module structure of $H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^{*+4}(MSO(4), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2) \otimes H^*(BU(1), \mathbb{Z}_2)$ below degree 5.




Fig. 26. $\Omega_*^{\text{Pin}^- \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)}$

Table 11

Bordism group. Here w_i is the Stiefel-Whitney class of the tangent bundle, w'_i is the Stiefel-Whitney class of the $\text{SO}(4)$ bundle, c_1 (c'_1) is the Chern class of the $\text{U}(1)$ bundle, $\tilde{\eta}$ is the mod 2 index of 1d Dirac operator, Arf is the Arf invariant.

Bordism group		
d	$\Omega_d^{\text{Pin}^- \times \mathbb{Z}_2 \text{Spin}(4) \times \text{U}(1) \times \text{U}(1)}$	generators
0	\mathbb{Z}_2	
1	0	
2	\mathbb{Z}_2^3	$w_1^2, c_1 \text{ mod } 2, c'_1 \text{ mod } 2$
3	0	
4	\mathbb{Z}_2^9	$w_1^4, w'_4, w_2^2, w_1^2 c_1, w_1^2 c'_1, w_2 c_1, w_2 c'_1, c_1 c'_1 \text{ mod } 2, (w_1^3 + w_1 w'_2 + w'_3) \tilde{\eta}$
5	\mathbb{Z}_2^3	$w'_2 w'_3, w'_4 \tilde{\eta}, (w_1^3 + w_1 w'_2 + w'_3) \text{Arf}$

Since the localization of $\text{Pin}^\pm \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}$ and $\text{Pin}^\pm \times \mathbb{Z}_2 (\text{U}(1) \times \text{SU}(3) \times \text{SU}(3))$ at the prime 2 are the same, the 2-torsion of $\Omega_d^{\text{Pin}^\pm \times \mathbb{Z}_2 \frac{\text{U}(3) \times \text{U}(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \text{U}(1))}}$ and $\Omega_d^{\text{Pin}^\pm \times \text{SU}(3) \times \text{SU}(3)}$ are the same.

We have $MT(\text{Pin}^c \times \text{SU}(3) \times \text{SU}(3)) = M\text{Spin} \wedge \Sigma^{-2} MU(1) \wedge \Sigma^{-1} MO(1) \wedge (\text{BSU}(3) \times \text{BSU}(3))_+$.

For $t - s < 8$, since there is no odd torsion, we have the Adams spectral sequence

$$\begin{aligned} & \text{Ext}_{\mathcal{A}_2(1)}^{s,t} (\mathbb{H}^{*+2}(MU(1), \mathbb{Z}_2) \otimes \mathbb{H}^{*+1}(MO(1), \mathbb{Z}_2) \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2) \\ & \quad \otimes \mathbb{H}^*(\text{BSU}(3), \mathbb{Z}_2), \mathbb{Z}_2) \\ & \Rightarrow \Omega_{t-s}^{\text{Pin}^c \times \text{SU}(3) \times \text{SU}(3)}. \end{aligned} \tag{3.17}$$

We have

$$\mathbb{H}^*(\text{BSU}(3) \times \text{BSU}(3), \mathbb{Z}_2) = \mathbb{Z}_2[c_2, c_3, c'_2, c'_3] \tag{3.18}$$




Fig. 27. The $\mathcal{A}_2(1)$ -module structure of $H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 6.




Fig. 28. $\Omega_*^{\text{Pin}^c} \times \text{SU}(3) \times \text{SU}(3)$.

where c_i is the Chern class of the $SU(3)$ bundle.

By Thom isomorphism,

$$H^{*+2}(MU(1), \mathbb{Z}_2) = \mathbb{Z}_2[c_1]U \tag{3.19}$$

where c_1 is the Chern class of the $U(1)$ bundle and U is the Thom class.

Table 12

Bordism group. Here c_1 is the Chern class of the $U(1)$ bundle, c_2 (c'_2) is the Chern class of the $SU(3)$ bundle, ABK is the Arf-Brown-Kervaire invariant.

		Bordism group	
	d	$\Omega_d^{\text{Pin}^c \times \text{SU}(3) \times \text{SU}(3)}$ $= \Omega_d^{\text{Pin}^+ \times \mathbb{Z}_2} \frac{U(3) \times U(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1))}$ $= \Omega_d^{\text{Pin}^- \times \mathbb{Z}_2} \frac{U(3) \times U(3)}{(\mathbb{Z}_3 \times \mathbb{Z}_3 \times U(1))}$	generators
0	\mathbb{Z}_2		
1	0		
2	\mathbb{Z}_4	ABK mod 4	
3	0		
4	$\mathbb{Z}_2^3 \times \mathbb{Z}_8$	$c_2 \pmod 2, c'_2 \pmod 2, c_1^2 \pmod 2, (c_1 \pmod 2) \text{ABK}$	
5	0		

Also by Thom isomorphism,

$$H^{*+1}(MO(1), \mathbb{Z}_2) = \mathbb{Z}_2[w_1]V \tag{3.20}$$

where w_1 is the Stiefel-Whitney class of the $O(1)$ bundle and V is the Thom class.

The $\mathcal{A}_2(1)$ -module structure of $H^{*+2}(MU(1), \mathbb{Z}_2) \otimes H^{*+1}(MO(1), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2) \otimes H^*(BSU(3), \mathbb{Z}_2)$ below degree 6 and the E_2 page are shown in Fig. 27, 28.

Combine the 2-torsion and 3-torsion results, we have in Table 12.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors are listed in the alphabetical order by the standard convention. JW thanks Kantaro Ohmori, Pavel Putrov, Nathan Seiberg, Edward Witten, and Yunqin Zheng for conversations. JW are also grateful to many other colleagues for helpful conversations, at Institute for Advanced Study, MIT, Princeton University, Harvard University, National Taiwan University and University of Tokyo. Part of this work was also reported by JW at the Aspen Center for Physics during “Field Theory Dualities and Strongly Correlated Matter,” March 18-24, 2018 under the title: “Time Reversal, $SU(N)$ Yang Mills, and Topological Phases” at http://www.its.caltech.edu/~motrunch/Aspen2018_Dualities/. ZW acknowledges previous supports from NSFC grants 11431010 and 11571329. ZW is supported by the Shuimu Tsinghua Scholar Program. JW was supported by NSF Grant PHY-1606531 at IAS. This work is also supported by NSF Grant DMS-1607871 “Analysis, Geometry and Mathematical Physics” and Center for Mathematical Sciences and Applications at Harvard University.

References

[1] K. Rajagopal, F. Wilczek, The Condensed matter physics of QCD, in: M. Shifman, B. Ioffe (Eds.), At the frontier of particle physics, in: Handbook of QCD, vol. 1–3, 2000, pp. 2061–2151, arXiv:hep-ph/0011333.

- [2] M.G. Alford, A. Schmitt, K. Rajagopal, T. Schäfer, Color superconductivity in dense quark matter, *Rev. Mod. Phys.* 80 (2008 Oct) 1455–1515, arXiv:0709.4635.
- [3] K. Fukushima, T. Hatsuda, The phase diagram of dense QCD, *Rep. Prog. Phys.* 74 (2011 Jan) 014001, arXiv:1005.4814.
- [4] M. Guo, P. Putrov, J. Wang, Time reversal, SU(N) Yang-Mills and cobordisms: interacting topological superconductors/insulators and quantum spin liquids in 3+1D, arXiv:1711.11587, 2017 Nov.
- [5] D. Gaiotto, A. Kapustin, N. Seiberg, B. Willett, Generalized global symmetries, *J. High Energy Phys.* 02 (2015) 172, arXiv:1412.5148.
- [6] G. 't Hooft, Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking, *NATO ASI Ser., Ser. B: Phys.* 59 (1980) 135.
- [7] S.L. Adler, Axial vector vertex in spinor electrodynamics, *Phys. Rev.* 177 (1969) 2426–2438.
- [8] J.S. Bell, R. Jackiw, A PCAC puzzle: $\pi^0 \rightarrow \gamma\gamma$ in the σ model, *Nuovo Cimento A* 60 (1969) 47–61.
- [9] L. Alvarez-Gaume, E. Witten, Gravitational anomalies, *Nucl. Phys. B* 234 (1984) 269.
- [10] E. Witten, An SU(2) anomaly, *Phys. Lett. B* 117 (1982) 324–328.
- [11] J. Wang, X.-G. Wen, E. Witten, A new SU(2) anomaly, *J. Math. Phys.* 60 (2019) 052301, arXiv:1810.00844.
- [12] J. Wang, L.H. Santos, X.-G. Wen, Bosonic anomalies, induced fractional quantum numbers and degenerate zero modes: the anomalous edge physics of symmetry-protected topological states, *Phys. Rev. B* 91 (2015) 195134, arXiv:1403.5256.
- [13] A. Kapustin, R. Thorngren, Anomalies of discrete symmetries in various dimensions and group cohomology, arXiv:1404.3230.
- [14] D. Gaiotto, A. Kapustin, Z. Komargodski, N. Seiberg, Theta, time reversal, and temperature, *J. High Energy Phys.* 05 (2017) 091, arXiv:1703.00501.
- [15] Z. Wan, J. Wang, Y. Zheng, New higher anomalies, SU(N) Yang-Mills gauge theory and $\mathbb{C}\mathbb{P}^{N-1}$ sigma model, arXiv:1812.11968.
- [16] Z. Wan, J. Wang, Y. Zheng, Quantum 4d Yang-Mills theory and time-reversal symmetric 5d higher-gauge topological field theory, *Phys. Rev. D* 100 (2019) 085012, arXiv:1904.00994.
- [17] E. Witten, Global gravitational anomalies, *Commun. Math. Phys.* 100 (1985) 197.
- [18] M.Z. Hasan, C.L. Kane, Colloquium: topological insulators, *Rev. Mod. Phys.* 82 (2010 Oct) 3045–3067, arXiv:1002.3895.
- [19] X.-L. Qi, S.-C. Zhang, Topological insulators and superconductors, *Rev. Mod. Phys.* 83 (2011 Oct) 1057–1110, arXiv:1008.2026.
- [20] A. Schnyder, S. Ryu, A. Furusaki, A.W.W. Ludwig, Classification of topological insulators and superconductors in three spatial dimensions, *Phys. Rev. B* 78 (2008) 195125.
- [21] A.Y. Kitaev, Periodic table for topological insulators and superconductors, *AIP Conf. Proc.* 1134 (22) (2009).
- [22] X.-G. Wen, Symmetry-protected topological phases in noninteracting fermion systems, *Phys. Rev. B* 85 (2012 Feb) 085103, arXiv:1111.6341.
- [23] X. Chen, Z.-C. Gu, Z.-X. Liu, X.-G. Wen, Symmetry protected topological orders and the group cohomology of their symmetry group, *Phys. Rev. B* 87 (2013 Apr) 155114, arXiv:1106.4772.
- [24] T. Senthil, Symmetry protected topological phases of quantum matter, *Annu. Rev. Condens. Matter Phys.* 6 (2015) 299, arXiv:1405.4015.
- [25] X.-G. Wen, Zoo of quantum-topological phases of matter, arXiv:1610.03911, 2016 Oct.
- [26] A. Kapustin, Symmetry protected topological phases, anomalies, and cobordisms: beyond group cohomology, arXiv:1403.1467, 2014 Mar.
- [27] A. Kapustin, R. Thorngren, A. Turzillo, Z. Wang, Fermionic symmetry protected topological phases and cobordisms, *J. High Energy Phys.* 12 (2015) 052, arXiv:1406.7329.
- [28] D.S. Freed, M.J. Hopkins, Reflection positivity and invertible topological phases, arXiv:1604.06527, 2016 Apr.
- [29] Z. Wan, J. Wang, Higher anomalies, higher symmetries, and cobordisms I: classification of higher-symmetry-protected topological states and their boundary fermionic/bosonic anomalies via a generalized cobordism theory, *Ann. Math. Sci. Appl.* 4 (2019) 107–311, arXiv:1812.11967.
- [30] Y. Nishida, Is a color superconductor topological?, *Phys. Rev. D* 81 (2010 Apr) 074004, arXiv:1001.2555.
- [31] M.M. Anber, E. Poppitz, Two-flavor adjoint QCD, *Phys. Rev. D* 98 (2018) 034026, arXiv:1805.12290.
- [32] C. Cordova, T.T. Dumitrescu, Candidate phases for SU(2) adjoint QCD₄ with two flavors from $\mathcal{N} = 2$ supersymmetric Yang-Mills theory, arXiv:1806.09592.
- [33] Z. Bi, T. Senthil, Adventure in topological phase transitions in 3+1 -D: non-Abelian deconfined quantum criticalities and a possible duality, *Phys. Rev. X* 9 (2019) 021034, arXiv:1808.07465.
- [34] Z. Wan, J. Wang, Adjoint QCD₄, deconfined critical phenomena, symmetry-enriched topological quantum field theory, and higher symmetry-extension, *Phys. Rev. D* 99 (2019) 065013, arXiv:1812.11955.

- [35] Z. Bi, E. Lake, T. Senthil, Landau ordering phase transitions beyond the Landau paradigm, arXiv:1910.12856.
- [36] J. Wang, Y.-Z. You, Y. Zheng, Gauge enhanced quantum criticality and time reversal domain wall: SU(2) Yang-Mills dynamics with topological terms, arXiv:1910.14664.
- [37] A. Cherman, S. Sen, M. Unsal, M.L. Wagman, L.G. Yaffe, Order parameters and color-flavor center symmetry in QCD, arXiv:1706.05385, 2017 June.
- [38] H. Shimizu, K. Yonekura, Anomaly constraints on deconfinement and chiral phase transition, arXiv:1706.06104, 2017 June.
- [39] Y. Tanizaki, Y. Kikuchi, T. Misumi, N. Sakai, Anomaly matching for phase diagram of massless \mathbb{Z}_N -QCD, arXiv:1711.10487, 2017 Nov.
- [40] Y. Tanizaki, Anomaly constraint on massless QCD and the role of Skyrmions in chiral symmetry breaking, J. High Energy Phys. 08 (171) (2018), arXiv:1807.07666.
- [41] K. Yonekura, Anomaly matching in QCD thermal phase transition, J. High Energy Phys. 05 (2019) 062, arXiv:1901.08188.
- [42] M.M. Anber, E. Poppitz, On the baryon-color-flavor (BCF) anomaly in vector-like theories, J. High Energy Phys. 11 (2019) 063, arXiv:1909.09027.
- [43] J. Wang, X.-G. Wen, E. Witten, Symmetric gapped interfaces of SPT and SET states: systematic constructions, arXiv:1705.06728, 2017 May.
- [44] T. Schafer, F. Wilczek, Continuity of quark and hadron matter, Phys. Rev. Lett. 82 (1999) 3956–3959, arXiv:hep-ph/9811473.
- [45] A. Cherman, S. Sen, L.G. Yaffe, Anyonic particle-vortex statistics and the nature of dense quark matter, Phys. Rev. D 100 (2019) 034015, arXiv:1808.04827.
- [46] Y. Hirono, Y. Tanizaki, Quark-hadron continuity beyond the Ginzburg-Landau paradigm, Phys. Rev. Lett. 122 (2019) 212001, arXiv:1811.10608.
- [47] Work to appear.
- [48] P. Putrov, J. Wang, S.-T. Yau, Braiding statistics and link invariants of bosonic/fermionic topological quantum matter in 2+1 and 3+1 dimensions, Ann. Phys. 384 (2017 Sept) 254–287, arXiv:1612.09298.
- [49] J. Wang, X.-G. Wen, S.-T. Yau, Quantum statistics and spacetime topology: quantum surgery formulas, Ann. Phys. 409 (2019) 167904, arXiv:1901.11537.
- [50] Y. Tanizaki, Y. Kikuchi, Vacuum structure of bifundamental gauge theories at finite topological angles, J. High Energy Phys. 06 (2017) 102, arXiv:1705.01949.
- [51] A. Karasik, Z. Komargodski, The bi-fundamental gauge theory in 3+1 dimensions: the vacuum structure and a cascade, J. High Energy Phys. 05 (2019) 144, arXiv:1904.09551.
- [52] Z. Wan, J. Wang, Beyond standard models and grand unifications: anomalies, topological terms and dynamical constraints via cobordisms, arXiv:1910.14668, 2019 Oct.
- [53] A. Beaudry, J.A. Campbell, A guide for computing stable homotopy groups, arXiv:1801.07530, 2018 Jan.