

Modeling of neutron-star mergers: a review while awaiting gravitational-wave detection

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 J. Phys.: Conf. Ser. 759 012004

(<http://iopscience.iop.org/1742-6596/759/1/012004>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 188.184.3.52

This content was downloaded on 21/11/2016 at 09:25

Please note that [terms and conditions apply](#).

You may also be interested in:

[On the robustness of the r-process in neutron-star mergers against variations of nuclear masses](#)

J J Mendoza-Temis, M R Wu, G Martínez-Pinedo et al.

[Collective excitations in neutron-star crusts](#)

N Chamel, D Page and S Reddy

[Early Advanced LIGO binary neutron-star sky localization and parameter estimation](#)

C P L Berry, B Farr, W M Farr et al.

[Maximum pulsar mass, equation of state and structure of neutron-star cores](#)

P Haensel and J L Zdunik

[Symmetry energy from nuclear masses and neutron-star observations using generalised Skyrme functionals](#)

N Chamel, A F Fantina, J M Pearson et al.

[THE FATE OF Cyg X-1: AN EMPIRICAL LOWER LIMIT ON BLACK-HOLE--NEUTRON-STAR MERGER RATE](#)

Krzysztof Belczynski, Tomasz Bulik and Charles Bailyn

[Recent progress and some open questions in nuclear astrophysics](#)

K Langanke and G Martínez-Pinedo

Modeling of neutron-star mergers: a review while awaiting gravitational-wave detection

Luca Baiotti

Graduate School of Science, Osaka University, Toyonaka, 560-0043, Japan

E-mail:

Abstract.

I review some recent results on simulations of mergers of binary neutron stars, highlighting some of the several significant advances published in the literature in the years 2013 to 2015.

1. Introduction

We are living in an exciting time for research on binary neutron-star (BNS) mergers. In fact, we are in a very dynamical phase of research, in which many accomplishments have been achieved (especially in the years 2013-2015), while many more need to be achieved to describe such fascinating physical phenomena. The first detection of gravitational waves [1] adds excitement to the field. Such first detection concerned a binary black-hole coalescence, but measurements of BNS mergers are also expected in the next months.

The details of the genesis of BNS systems are still unclear, but the general picture seems well accepted [2]. The more massive star of the binary explodes as a supernova once it has evolved off the main sequence and through its giant phase. The remnant of such an explosion becomes the first neutron star of the binary, with a potential recoil. Subsequently, the secondary star evolves off the main sequence and explodes as a supernova becoming the second neutron star.

Many astronomical observations revealed that BNS indeed exist [3]. The inspiral and merger of two neutron stars in binary orbit is the inevitable fate of close-binary evolution, since their angular momentum is dissipated through the emission of gravitational radiation. Here is a brief description of their evolution. As the inspiral progressively speeds up and the two neutron stars approach each other, tidal waves produced by the tidal interaction become visible at their surface. At the merger, the two stars collide with a rather large impact parameter. A *vortex sheet* (or *shear interface*) develops, where the tangential component of the velocity exhibits a discontinuity. This condition is known to be unstable to very small perturbations and it can develop a *Kelvin-Helmholtz instability*, which curls the interface forming a series of vortices at all wavelengths [4, 5]. Even if this instability is purely hydrodynamical and it is likely to be important only for binaries with very similar masses, it can have strong consequences if the stars have magnetic fields. It has been shown in most simulations [6, 7, 8, 9, 10, 11] that, in the presence of an initially poloidal magnetic field, this instability may lead to an exponential growth of the toroidal component. Such a growth is the result of the exponentially rapid formation of vortices that curl magnetic field lines that were initially purely poloidal. The exponential growth



caused by the Kelvin–Helmholtz instability leads to an overall amplification of the magnetic field of up to a few orders of magnitude [10, 11].

After the merger, unless the progenitor stars have very small masses or for some particular equations of state (EoSs), the compact object newly formed is expected to collapse to a black hole, either promptly or after a certain amount of time. If the collapse is not prompt, the merged object is expected to be temporarily a hypermassive neutron star (HMNS), namely a neutron star whose rest mass exceeds the maximum rest mass of nonspinning or uniformly spinning neutron stars and is sustained against collapse by differential rotation. The merged object oscillates violently, with a dominant $m = 2$ non-axisymmetric character [12]. Dissipative effects like viscosity, magnetic fields or gravitational-wave emission bring then the star towards a configuration which is unstable to gravitational collapse. The time interval between merger and collapse can be roughly estimated to be of the order of 1–100 s or more. The large uncertainty in determining this is due particularly to the difficulty of performing accurate numerical simulations over these very long time-scales. During this time, if the merged object has a sufficiently high ratio of rotational to gravitational binding energy, it could also become dynamically unstable to nonlinear instabilities leading to a bar-mode deformation [13, 14, 15]. Both magnetic fields [7, 10, 11] and radiative processes [16, 17, 18, 9, 19] can modify significantly the lifetime of the HMNS and hence change the properties of the black hole and of the surrounding torus. In most cases when a black hole is formed some amount of matter remains outside of it. This leads to the formation of an accretion torus that may be rather dense ($10^{12} - 10^{13} \text{ g cm}^{-3}$) and extended horizontally for tens of kilometers and vertically for a few tens of kilometers.

BNS mergers are expected (i) to be significant sources of gravitational radiation, not only during the inspiral, but also during and after the merger, (ii) to be possible progenitors for short-gamma-ray bursts (GRBs), whose short rise times suggest that their central sources have to be highly relativistic objects, and (iii) to be the possible sources of other electromagnetic (e.g. macronovae [20, 21]) and neutrino emission.

The typical scenario leading to GRBs assumes that a system composed of a rotating black hole and a surrounding massive torus is formed after the merger. If the disc had a mass $\gtrsim 0.01 - 0.1 M_{\odot}$, it could provide the large amount of energy observed in SGRBs, either through neutrino processes or by extracting the rotational energy of the black hole via magnetic fields [22, 23, 8]. Such a configuration is expected to form in most BNS mergers.

As far as gravitational-wave emission is concerned, BNS systems are expected to produce signals of amplitude large enough to be relevant for Earth-based gravitational-wave detectors such as LIGO [24], Virgo [25], or Kagra [26]. They are also expected to be sufficiently frequent sources: Advanced interferometric detectors are in fact expected to observe them at a rate of $\sim 0.4 - 400$ events per year, with a most realistic rate of ~ 40 per year [27]. Among other results, observations of gravitational waves, photons, and neutrinos emitted during and after BNS mergers will give strong indications about the EoS of matter at nuclear densities, which cannot be probed in laboratories on Earth and is not fully understood at the theoretical level [28].

During the inspiral phase, when the two stars are still well separated, sophisticated post-Newtonian expansions offer a good approximate description of the dynamics, but only full numerical simulations can describe the highly nonlinear regime of the merger and post-merger phases. The final goal of numerical computations may be a simulation that includes the solution of the Einstein equations, the relativistic hydrodynamic and (resistive) magnetohydrodynamics (MHD) equations, EoSs based on microphysical calculations, neutrino and photon radiation transport, nuclear-reaction networks. On top of all this, codes should possibly implement high-order, high-accuracy numerical methods and run fast enough to allow parameter-space exploration. This is a goal still very far in the future. But when we achieve all this, the reward will be great, namely an accurate interpretative and predictive description of the processes related to BNS systems and involving and causing short GRBs, jet formation and emission,

particle formation and emission, other electromagnetic emission like that of macronovae, heavy-element abundance. As far as the study and interpretation of gravitational-wave observations is concerned, we may be able to obtain satisfactory results even without the full machinery introduced above. In fact, the interesting gravitational radiation is probably emitted before neutrino, radiation transport, and nuclear reactions become important. The evolution of the merged object, instead, depends on a large number of details, such as the mass-ratio of the initial stars, the EoS, the neutrino physics and precise energy transport, the dynamical relevance of the magnetic field, etc.

Several groups are working on BNS simulations with their own independent codes. Most of the codes solve the full Einstein Equations without approximations (except for the truncation error and the errors associated to numerical methods), but there are two notable exceptions, namely the code of Rosswog [29, 30, 31, 32] and the code of Janka/Bauswein [33, 34, 35]. These two smoothed-particle-hydrodynamics (SPH) codes make theoretical approximations to the evolution of gravity / spacetime but are among the most advanced in treating microphysical processes.

The first successful fully general-relativistic simulations of BNS mergers were presented at the end of the past century in [36] and huge advances have been made since. Several groups have developed numerical codes for general-relativistic simulations. All of these codes share the following capabilities:

- Implementation and use of high-order finite-differencing techniques¹ for the solution of the Einstein equations
- Implementation and use of high-resolution shock-capturing (HRSC) methods for the solution of the GRHD/GRMHD equations
- Implementation and use of adaptive mesh-refinement (AMR) techniques that provide higher resolution around the orbiting stars and the merged object
- Use of consistent initial data representing a system of BNS in quasi-circular orbits²
- Accurate evolution of matter and spacetime (including long-term evolutions of the formed black holes and accretion discs), thanks to the numerical methods mentioned above
- Analysis of the properties of the black holes produced in the merger, *e.g.* through the calculation of trapped surfaces
- Extraction of gravitational waveforms (usually through computation of the Weyl tensor)

Intense work is ongoing and first satisfactory results are being obtained on linking future gravitational-wave observations to physical properties of the emitting system (*e.g.* relating the main frequency of postmerger oscillations to the neutron-star masses) [41, 42, 43, 44, 45, 46, 47, 48] and on heavy-element production and macronovae [49, 31, 50, 32, 51, 52, 53, 54, 55, 56, 57, 58, 59, 19, 60]. There are then still issues that are very open, like the accurate dynamics of magnetic fields after the merger [8, 10, 11] (and before the merger for resistive MHD [61, 62, 63, 64, 65]), neutrino treatment [16, 17, 66, 18, 67, 68, 9, 69, 70, 71, 19] and photon radiation transport [70].

Most of the general-relativistic codes [36, 72, 73, 74, 10, 75, 76, 77, 78, 79, 61, 80, 81, 82, 83, 62, 84, 85, 86, 87] solve the field equations in the Baumgarte-Shapiro-Shibata-Nakamura formulation [88, 89]. The remainder [90, 91, 92, 93] use the generalized-harmonic formulation [94, 95]. Most groups use finite-difference methods for the metric evolution and

¹ One group is developing a code based on pseudo-spectral techniques, but no publication on BNS mergers has been produced yet, except the preliminary studies of [37, 38]. It is named the *SpEC-hydro* code and employs a dual-grid method in which the metric fields are solved using pseudospectral methods on a multidomain spectral grid, and the matter fields are solved using high-resolution shock-capturing finite volume schemes on a Cartesian finite-difference grid.

² Some studies focus on purpose on different types of initial data [39, 40, 31, 32].

conservative, HRSC schemes for the hydrodynamics evolution. In all cases, these unigrid algorithms are then interfaced with some sort of AMR.

Some groups have implemented the MHD equations in full general relativity; since these codes all make use of conservative HRSC methods, their main differences are in how they meet the challenge of preserving the $\text{div}\cdot\mathbf{B} = 0$ constraint, especially in AMR³. Besides MHD, the other major advances in the physical modelling for numerical relativity codes have been in the arena of microphysics. While the polytropic or ideal-fluid EoSs have been the community standard for quite some time, most codes now allow for a nuclear theory-based EoSs and/or use various parameterized, piecewise polytropic EoSs inspired by the range of plausible nuclear EoSs [96]. Some groups have also started to account for neutrino transport via simplified leakage schemes [16, 17, 66, 18, 67, 68, 9, 69] and are also studying formulations for more accurate treatments [70, 71, 19], which show much promise for numerical relativity simulations with neutrino physics beyond the leakage approximation.

2. Status of models, techniques, and results for general-relativistic simulations

General relativistic hydrodynamics simulations of BNS started being performed more than 15 years ago [72, 36, 97, 98]. Even if nowadays many state-of-the-art codes are able to solve more physical equations (for magnetic fields, neutrinos, radiation, etc.), simulations involving only general-relativistic hydrodynamics (*pure GRHD*) are still the benchmark for any new code and the necessary testbed for more advanced codes. Furthermore, in many cases, results obtained with pure GRHD - notably inspiral gravitational waveforms - are a sufficiently good description of BNS systems. During the merger and after a single object forms, possibly surrounded by a massive disc, it is necessary to consider additional physical components in order to have a satisfactory description.

One particularly important aspect of BNS simulations is the EoS. While detectable differences between simulations that employed different EoSs already appear during the inspiral [96, 99], it is really the post-merger phase that is markedly different. Hence, an accurate description of the post-merger evolution is essential to extract information concerning the neutron-star interior structure. These particularly interesting parts of the waveforms that are more likely to yield fundamental clues to the physics beyond nuclear density are at very high frequencies and thus probably only marginally detectable by detectors like advanced LIGO. Third-generation detectors, such as ET [100], may provide the first realistic opportunity to use gravitational waves to decipher the stellar structure and EoS [101].

First attempts to single out the effects of the EoS on the inspiral and merger were done in Refs. [102, 103, 74, 104]. The authors there focus mostly on the dynamics of equal-mass binaries, as these are thought to be the most common [105] and are easier and faster to compute, since symmetries of the configuration can be exploited to save computational resources. In Baiotti et al. [104], an investigation was made using two analytic EoSs: a *cold*, *i.e.* polytropic, EoS, or a *hot*, *i.e.* ideal-fluid (also called *simple fluid* or *gamma-law*), EoS. Although the polytropic EoS is isentropic and thus unrealistic for describing the post-merger evolution, it provides an adequate description of the inspiral phase, during which the neutron stars are expected to interact only gravitationally [106]. The most marked differences between *cold* and *hot* EoSs are in the post-merger phase.

The study of the effect of realistic EoSs in GRHD simulations has been subsequently especially brought forward by the Shibata group⁴, starting from Ref. [107], where they employ the Akmal-Pandharipande-Ravenhall (APR) EoS [108] augmented, after the onset of merger, by a *hot* part through an ideal-fluid EoS. In another work of the same group [109], the authors studied the

³ Straightforward finite-difference evolutions of the magnetic field would generically lead to magnetic monopoles and, hence, unphysical behavior.

⁴ Other groups have implemented the ability to use tabulated EoSs on their codes, see [68, 9]

dependence of the dynamical behavior of BNS mergers on the EoS of the supernuclear-density matter with piecewise-polytropic EoSs [110]. Table I in [109] usefully summarizes the piecewise polytropic parameters of several realistic EoSs.

In a work published soon after, the same group showed results of simulations performed by incorporating both nucleonic and hyperonic finite-temperature EoS [17, 18]. Reported results are of interest, even if the hyperonic EoS used here is limited by the fact that it only takes Λ hyperons into account and that it cannot produce a stable neutron star with realistic masses⁵. It was found that also for the hyperonic EoS, a HMNS is first formed after the merger and subsequently collapses to a black hole. The radius of such a HMNS decreases in time because of the increase of the mass fraction of hyperons and the consequent decrease in supporting pressure. Such a shrinking is noticeably larger than the one simply due to angular-momentum loss through gravitational-wave emission that is present also in nucleonic EoSs. These dynamics are clearly visible in the gravitational-wave signal, whose characteristic peak frequency has an increase of 20%-30% during the HMNS evolution. By contrast, for the nucleonic EoS the peak gravitational-wave frequency in the HMNS phase is approximately constant. The authors stress that their results raise a warning about using the peak frequency of the gravitational-wave spectrum to extract information of the neutron-star matter, because it may evolve and so make the relation of the peak frequency with the HMNS structure ambiguous. Finally, the authors found that the torus mass for the hyperonic EoS is smaller than that for the nucleonic EoS. This makes hyperonic EoSs less favorite for the description of short gamma-ray bursts.

The most recent works of the Shibata group on EoSs are Ref. [44, 99], where they use their AMR SACRA code [74] to perform a large number of simulations with a variety of EoSs and mass ranges. On the basis of their results they derive a fitting formula for the quasiperiodic gravitational waveforms, which they say may be used for the analysis of future observed gravitational-wave signals. They choose two types of EoSs: piecewise polytropic EoSs [110] and the tabulated Shen EoS [114]. In both cases, they add approximate finite-temperature effects to the cold EoSs through an additional ideal-fluid term.

They put in evidence that the universal features of gravitational waves emitted by HMNSs include, in order: (i) a peak in frequency and amplitude soon after the merger starts, (ii) a decrease in amplitude during the merger and a new increase when the HMNS forms, (iii) a damped oscillation of the frequency during the HMNS phase lasting for several oscillation periods and eventually settling to an approximately constant value (although a long-term secular change associated with the change of the state of HMNSs is always present), (iv) a final decrease in the amplitude during the HMNS phase, either monotonical or with modulations. Based on this, they find an optimal 13-parameter fitting function. Using the fitted functions it may be possible to constrain the neutron-star radius with errors of about 1 km.

3. Outlook

In this article, because of limited space, I have been able to review only one topic about BNS merger simulations, namely the influence of the EoS. The phenomenology and physics of BNS mergers, however, are much richer. Simulations of BNS merger and post-merger phases will soon be connected to observations of gravitational waves and will allow us to interpret them and therefore to uncover the mysteries of the physical and astrophysical properties of relativistic stars.

References

- [1] Abbott B P, Abbott R, Abbott T D, Abernathy M R, Acernese F, Ackley K, Adams C, Adams T, Addesso P, Adhikari R X and et al 2016 *Physical Review Letters* **116** 061102 (*Preprint* 1602.03837)

⁵ Hyperonic EoSs, which are generally soft, are currently disfavored by the observation of a $2M_{\odot}$ star [111, 112]. However, the existence of exotic phases in neutron stars remains unconstrained [113].

- [2] Kalogera V, Belczynski K, Kim C, O’Shaughnessy R and Willems B 2007 *Physics Reports* **442** 75–108 (Preprint [arXiv:astro-ph/0612144](#))
- [3] Kramer M, Lyne A, Burgay M, Possenti A, Manchester R, Camilo F, McLaughlin M, Lorimer D, D’Amico N, Joshi B, Reynolds J and Freire P 2004 The double pulsar – a new testbed for relativistic gravity *Binary Pulsars* ed Rasio and Stairs (Chicago: PSAP)
- [4] Chandrasekhar S 1981 *Hydrodynamic and hydromagnetic stability* (New York, USA: Dover Edition)
- [5] Bodo G, Massaglia S, Ferrari A and Trussoni E 1994 *Astron. Astrophys.* **283** 655–676
- [6] Price D J and Rosswog S 2006 *Science* **312** 719–722 (Preprint [astro-ph/0603845](#))
- [7] Giacomazzo B, Rezzolla L and Baiotti L 2011 *Phys. Rev. D* **83** 044014
- [8] Rezzolla L, Giacomazzo B, Baiotti L, Granot J, Kouveliotou C and Aloy M A 2011 *Astrophys. J. Letters* **732** L6 (Preprint [1101.4298](#))
- [9] Neilsen D W, Liebling S L, Anderson M, Lehner L, O’Connor E and Palenzuela C 2014 *Phys. Rev. D* **89** 104029 (Preprint [1403.3680](#))
- [10] Kiuchi K, Kyutoku K, Sekiguchi Y, Shibata M and Wada T 2014 *Phys. Rev. D* **90** 041502 (Preprint [1407.2660](#))
- [11] Kiuchi K, Cerdá-Durán P, Kyutoku K, Sekiguchi Y and Shibata M 2015 *Phys. Rev. D* **92** 124034 (Preprint [1509.09205](#))
- [12] Stergioulas N, Bauswein A, Zagkouris K and Janka H T 2011 *Mon. Not. R. Astron. Soc.* **418** 427–436 (Preprint [1105.0368](#))
- [13] Shibata M, Baumgarte T W and Shapiro S L 2000 *Astrophys. J.* **542** 453–463 (Preprint [astro-ph/0005378](#))
- [14] Baiotti L, de Pietri R, Manca G M and Rezzolla L 2007 *Phys. Rev. D* **75** 044023 (Preprint [astro-ph/0609473](#))
- [15] Franci L, De Pietri R, Dionysopoulou K and Rezzolla L 2013 *Journal of Physics Conference Series* **470** 012008 (Preprint [1309.6549](#))
- [16] Sekiguchi Y, Kiuchi K, Kyutoku K and Shibata M 2011 *Physical Review Letters* **107** 051102 (Preprint [1105.2125](#))
- [17] Sekiguchi Y, Kiuchi K, Kyutoku K and Shibata M 2011 *Phys. Rev. Lett.* **107** 211101 (Preprint [1110.4442](#))
- [18] Sekiguchi Y, Kiuchi K, Kyutoku K and Shibata M 2012 *Progress of Theoretical and Experimental Physics* **2012** 01A304 (Preprint [1206.5927](#))
- [19] Sekiguchi Y, Kiuchi K, Kyutoku K and Shibata M 2015 *Phys. Rev. D* **91** 064059 (Preprint [1502.06660](#))
- [20] Li L X and Paczyński B 1998 *Astrophys. J.* **507** L59–L62 (Preprint [astro-ph/9807272](#))
- [21] Metzger B D, Martínez-Pinedo G, Darbha S, Quataert E, Arcones A, Kasen D, Thomas R, Nugent P, Panov I V and Zinner N T 2010 *Mon. Not. R. Astron. Soc.* **406** 2650–2662 (Preprint [1001.5029](#))
- [22] Paczynski B 1986 *Astrophys. J. Lett.* **308** L43–L46
- [23] Eichler D, Livio M, Piran T and Schramm D N 1989 *Nature* **340** 126–128
- [24] Abadie J, Abbott B P, Abbott R, Abernathy M, Adams C, Adhikari R, Ajith P, Allen B, Allen G, Amador Ceron E and et al 2010 *Nuclear Instruments and Methods in Physics Research A* **624** 223–240 (Preprint [1007.3973](#))
- [25] Accadia T et al. 2011 *Class. Quantum Grav.* **28** 114002
- [26] Kuroda K and LCGT Collaboration 2010 *Class. Quantum Grav.* **27** 084004
- [27] Abadie J, Abbott B P, Abbott R, Abernathy M, Accadia T, Acernese F, Adams C, Adhikari R, Ajith P, Allen B and et al 2010 *Class. Quantum Grav.* **27** 173001 (Preprint [1003.2480](#))
- [28] Lattimer J M and Prakash M 2007 *Phys. Rep.* **442** 109–165 (Preprint [arXiv:astro-ph/0612440](#))
- [29] Rosswog S, Davies M B, Thielemann F K and Piran T 2000 *Astron. Astrophys.* **360** 171–184 (Preprint [astro-ph/0005550](#))
- [30] Rosswog S 2010 *Journal of Computational Physics* **229** 8591–8612 (Preprint [0907.4890](#))
- [31] Rosswog S, Piran T and Nakar E 2013 *Mon. Not. R. Astron. Soc.* **430** 2585–2604 (Preprint [1204.6240](#))
- [32] Piran T, Nakar E and Rosswog S 2013 *Mon. Not. R. Astron. Soc.* **430** 2121–2136 (Preprint [1204.6242](#))
- [33] Oechslin R, Rosswog S and Thielemann F K 2002 *Phys. Rev. D* **65** 103005 (Preprint [gr-qc/0111005](#))
- [34] Oechslin R, Janka H T and Marek A 2007 *Astron. Astrophys.* **467** 395–409 (Preprint [astro-ph/0611047](#))
- [35] Bauswein A, Janka H and Oechslin R 2010 *Phys. Rev. D* **82** 084043 (Preprint [1006.3315](#))
- [36] Shibata M and Uryū K 2000 *Phys. Rev. D* **61** 064001 (Preprint [gr-qc/9911058](#))
- [37] Kaplan J D 2014 *Where Tori Fear to Tread: Hypermassive Neutron Star Remnants and Absolute Event Horizons or Topics in Computational General Relativity* Ph.D. thesis California Institute of Technology
- [38] Foucart F, Haas R, Duez M D, O’Connor E, Ott C D, Roberts L, Kidder L E, Lippuner J, Pfeiffer H P and Scheel M A 2015 *ArXiv e-prints* (Preprint [1510.06398](#))
- [39] Gold R, Bernuzzi S, Thierfelder M, Brüggmann B and Pretorius F 2012 *Phys. Rev. D* **86** 121501 (Preprint [1109.5128](#))
- [40] East W E and Pretorius F 2012 *Astrophys. J.* **760** L4 (Preprint [1208.5279](#))

- [41] Bauswein A and Janka H T 2012 *Phys. Rev. Lett.* **108** 011101 (*Preprint* 1106.1616)
- [42] Bauswein A and Janka H T 2012 *Phys. Rev. Lett.* **108** 011101 (*Preprint* 1106.1616)
- [43] Bauswein A, Janka H T, Hebeler K and Schwenk A 2012 *Phys. Rev. D* **86** 063001 (*Preprint* 1204.1888)
- [44] Hotokezaka K, Kiuchi K, Kyutoku K, Muranushi T, Sekiguchi Y, Shibata M and Taniguchi K 2013 *Phys. Rev. D* **88** 044026 (*Preprint* 1307.5888)
- [45] Bauswein A, Stergioulas N and Janka H T 2014 *Phys. Rev. D* **90** 023002 (*Preprint* 1403.5301)
- [46] Takami K, Rezzolla L and Baiotti L 2014 *Phys. Rev. Lett.* **113** 091104 (*Preprint* 1403.5672)
- [47] Clark J, Bauswein A, Cadonati L, Janka H T, Pankow C and Stergioulas N 2014 *Phys. Rev. D* **90** 062004 (*Preprint* 1406.5444)
- [48] Takami K, Rezzolla L and Baiotti L 2015 *Phys. Rev. D* **91** 064001 (*Preprint* 1412.3240)
- [49] Rosswog S 2013 *Nature* **500** 535–536
- [50] Rosswog S 2013 *Royal Society of London Philosophical Transactions Series A* **371** 20272 (*Preprint* 1210.6549)
- [51] Bauswein A, Goriely S and Janka H T 2013 *Astrophys. J.* **773** 78 (*Preprint* 1302.6530)
- [52] Tanaka M and Hotokezaka K 2013 *Astrophys. J.* **775** 113 (*Preprint* 1306.3742)
- [53] Hotokezaka K, Kyutoku K, Tanaka M, Kiuchi K, Sekiguchi Y, Shibata M and Wanajo S 2013 *Astrophys. J.* **778** L16 (*Preprint* 1310.1623)
- [54] Metzger B D and Fernández R 2014 *Mon. Not. R. Astron. Soc.* **441** 3444–3453 (*Preprint* 1402.4803)
- [55] Rosswog S, Korobkin O, Arcones A, Thielemann F K and Piran T 2014 *Mon. Not. R. Astron. Soc.* **439** 744–756 (*Preprint* 1307.2939)
- [56] Grossman D, Korobkin O, Rosswog S and Piran T 2014 *Mon. Not. R. Astron. Soc.* **439** 757–770 (*Preprint* 1307.2943)
- [57] Perego A, Rosswog S, Cabezón R M, Korobkin O, Käppeli R, Arcones A and Liebendörfer M 2014 *Mon. Not. R. Astron. Soc.* **443** 3134–3156 (*Preprint* 1405.6730)
- [58] Wanajo S, Sekiguchi Y, Nishimura N, Kiuchi K, Kyutoku K and Shibata M 2014 *Astrophys. J.* **789** L39 (*Preprint* 1402.7317)
- [59] Just O, Bauswein A, Pulpillo R A, Goriely S and Janka H T 2015 *Mon. Not. R. Astron. Soc.* **448** 541–567 (*Preprint* 1406.2687)
- [60] Goriely S, Bauswein A, Just O, Pllumbi E and Janka H T 2015 *Mon. Not. R. Astron. Soc.* **452** 3894–3904 (*Preprint* 1504.04377)
- [61] Dionysopoulou K, Alic D, Palenzuela C, Rezzolla L and Giacomazzo B 2013 *Phys. Rev. D* **88** 044020 (*Preprint* 1208.3487)
- [62] Palenzuela C 2013 *Mon. Not. R. Astron. Soc.* **431** 1853–1865 (*Preprint* 1212.0130)
- [63] Palenzuela C, Lehner L, Ponce M, Liebling S L, Anderson M, Neilsen D and Motl P 2013 *Phys. Rev. Lett.* **111** 061105 (*Preprint* 1301.7074)
- [64] Palenzuela C, Lehner L, Liebling S L, Ponce M, Anderson M, Neilsen D and Motl P 2013 *Phys. Rev. D* **88** 043011 (*Preprint* 1307.7372)
- [65] Dionysopoulou K, Alic D and Rezzolla L 2015 *Phys. Rev. D* **92** 084064 (*Preprint* 1502.02021)
- [66] Kiuchi K, Sekiguchi Y, Kyutoku K and Shibata M 2012 *Class. Quantum Grav.* **29** 124003 (*Preprint* 1206.0509)
- [67] Kiuchi K, Sekiguchi Y, Kyutoku K and Shibata M 2012 *Gravitational Waves and Neutrino Emissions from the Merger of Binary Neutron Stars Numerical Modeling of Space Plasma Slows (ASTRONUM 2011) (Astronomical Society of the Pacific Conference Series vol 459)* ed Pogorelov N V, Font J A, Audit E and Zank G P p 85
- [68] Galeazzi F, Kastaun W, Rezzolla L and Font J A 2013 *Phys. Rev. D* **88** 064009 (*Preprint* 1306.4953)
- [69] Palenzuela C, Liebling S L, Neilsen D, Lehner L, Caballero O L, O’Connor E and Anderson M 2015 *Phys. Rev. D* **92** 044045 (*Preprint* 1505.01607)
- [70] Shibata M, Kiuchi K, Sekiguchi Y and Suwa Y 2011 *Progress of Theoretical Physics* **125** 1255–1287 (*Preprint* 1104.3937)
- [71] Radice D, Abdikamalov E, Rezzolla L and Ott C D 2013 *Journal of Computational Physics* **242** 648–669 (*Preprint* 1209.1634)
- [72] Shibata M 1999 *Phys. Rev. D* **60** 104052 (*Preprint* gr-qc/9908027)
- [73] Shibata M 1999 *Progress of Theoretical Physics* **101** 1199–1233 (*Preprint* gr-qc/9905058)
- [74] Yamamoto T, Shibata M and Taniguchi K 2008 *Phys. Rev. D* **78** 064054 (*Preprint* 0806.4007)
- [75] Baiotti L, Hawke I, Montero P and Rezzolla L 2003 *A new three-dimensional general-relativistic hydrodynamics code Computational Astrophysics in Italy: Methods and Tools vol 1* ed Capuzzo-Dolcetta R (Trieste: MSAIt) p 210
- [76] Baiotti L 2004 *Numerical relativity simulations of non-vacuum spacetimes in three dimensions* Ph.D. thesis SISSA, International School for advanced studies

- [77] Baiotti L, Hawke I, Montero P J, Löffler F, Rezzolla L, Stergioulas N, Font J A and Seidel E 2005 *Phys. Rev. D* **71** 024035 (Preprint gr-qc/0403029)
- [78] Giacomazzo B and Rezzolla L 2007 *Classical and Quantum Gravity* **24** 235 (Preprint gr-qc/0701109)
- [79] Kellerman T, Baiotti L, Giacomazzo B and Rezzolla L 2008 *Class. Quantum Grav.* **25** 225007 (Preprint 0811.0938)
- [80] Kastaun W, Galeazzi F, Alic D, Rezzolla L and Font J A 2013 *Phys. Rev. D* **88** 021501 (Preprint 1301.7348)
- [81] Baiotti L, Shibata M and Yamamoto T 2010 *Phys. Rev. D* **82** 064015 (Preprint 1007.1754)
- [82] Anderson M, Hirschmann E W, Lehner L, Liebling S L, Motl P M, Neilsen D, Palenzuela C and Tohline J E 2008 *Phys. Rev. D* **77** 024006 (Preprint 0708.2720)
- [83] Anderson M, Hirschmann E W, Lehner L, Liebling S L, Motl P M, Neilsen D, Palenzuela C and Tohline J E 2008 *Phys. Rev. Lett.* **100** 191101 (Preprint 0801.4387)
- [84] Löffler F, Faber J, Bentivegna E, Bode T, Diener P, Haas R, Hinder I, Mundim B C, Ott C D, Schnetter E, Allen G, Campanelli M and Laguna P 2012 *Class. Quantum Grav.* **29** 115001 (Preprint 1111.3344)
- [85] Zilhão M and Löffler F 2013 *International Journal of Modern Physics A* **28** 1340014 (Preprint 1305.5299)
- [86] Moesta P, Mundim B, Faber J, Noble S, Bode T, Haas R, Loeffler F, Ott C, Reisswig C and Schnetter E 2013 General relativistic magneto-hydrodynamics with the Einstein Toolkit *APS Meeting Abstracts* p 10001
- [87] Thierfelder M, Bernuzzi S and Brüggmann B 2011 *Phys. Rev. D* **84** 044012 (Preprint 1104.4751)
- [88] Shibata M and Nakamura T 1995 *Phys. Rev. D* **52** 5428–5444
- [89] Baumgarte T W and Shapiro S L 1999 *Phys. Rev. D* **59** 024007 (Preprint gr-qc/9810065)
- [90] Duez M D *et al.* 2008 *Phys. Rev. D* **78** 104015 (Preprint 0809.0002)
- [91] Anderson M, Hirschmann E W, Liebling S L and Neilsen D 2006 *Classical and Quantum Gravity* **23** 6503–6524 (Preprint gr-qc/0605102)
- [92] Lindblom L, Scheel M A, Kidder L E, Owen R and Rinne O 2006 *Classical and Quantum Gravity* **23** 447 (Preprint gr-qc/0512093)
- [93] East W E, Pretorius F and Stephens B C 2012 *Phys. Rev. D* **85** 124010 (Preprint 1112.3094)
- [94] Garfinkle D 2002 *Phys. Rev. D* **65** 044029 (Preprint gr-qc/0110013)
- [95] Pretorius F 2005 *Classical and Quantum Gravity* **22** 425–451 (Preprint gr-qc/0407110)
- [96] Read J S, Baiotti L, Creighton J D E, Friedman J L, Giacomazzo B, Kyutoku K, Markakis C, Rezzolla L, Shibata M and Taniguchi K 2013 *Phys. Rev. D* **88** 044042 (Preprint 1306.4065)
- [97] Shibata M and Uryū K 2002 *Progress of Theoretical Physics* **107** 265–303 (Preprint gr-qc/0203037)
- [98] Shibata M, Taniguchi K and Uryū K 2003 *Phys. Rev. D* **68** 084020 (Preprint gr-qc/0310030)
- [99] Hotokezaka K, Kyutoku K, Okawa H and Shibata M 2015 *Phys. Rev. D* **91** 064060 (Preprint 1502.03457)
- [100] Punturo M *et al.* 2010 *Class. Quantum Grav.* **27** 084007
- [101] Andersson N, Ferrari V, Jones D I, Kokkotas K D, Krishnan B, Read J S, Rezzolla L and Zink B 2011 *General Relativity and Gravitation* **43** 409–436 (Preprint 0912.0384)
- [102] Shibata M, Taniguchi K and Uryū K 2005 *Phys. Rev. D* **71** 084021 (Preprint gr-qc/0503119)
- [103] Shibata M and Taniguchi K 2006 *Phys. Rev. D* **73** 064027 (Preprint astro-ph/0603145)
- [104] Baiotti L, Giacomazzo B and Rezzolla L 2008 *Phys. Rev. D* **78** 084033 (Preprint 0804.0594)
- [105] Osłowski S, Bulik T, Gondek-Rosińska D and Belczyński K 2011 *Mon. Not. R. Astron. Soc.* **413** 461–479 (Preprint 0903.3538)
- [106] Giacomazzo B, Rezzolla L and Baiotti L 2009 *Mon. Not. R. Astron. Soc.* **399** L164–L168
- [107] Kiuchi K, Sekiguchi Y, Shibata M and Taniguchi K 2009 *Phys. Rev. D* **80** 064037 (Preprint 0904.4551)
- [108] Akmal A, Pandharipande V R and Ravenhall D G 1998 *Phys. Rev. C* **58** 1804–1828 (Preprint arXiv:hep-ph/9804388)
- [109] Hotokezaka K, Kyutoku K, Okawa H, Shibata M and Kiuchi K 2011 *Phys. Rev. D* **83** 124008 (Preprint 1105.4370)
- [110] Read J S, Markakis C, Shibata M, Uryū K, Creighton J D E and Friedman J L 2009 *Phys. Rev. D* **79** 124033 (Preprint 0901.3258)
- [111] Antoniadis J, Freire P C C, Wex N, Tauris T M, Lynch R S, van Kerkwijk M H, Kramer M, Bassa C, Dhillon V S, Driebe T, Hessels J W T, Kaspi V M, Kondratiev V I, Langer N, Marsh T R, McLaughlin M A, Pennucci T T, Ransom S M, Stairs I H, van Leeuwen J, Verbiest J P W and Whelan D G 2013 *Science* **340** 1233232 (Preprint 1304.6875)
- [112] Demorest P B, Pennucci T, Ransom S M, Roberts M S E and Hessels J W T 2010 *Nature* **467** 1081–1083 (Preprint 1010.5788)
- [113] Bhowmick B, Bhattacharya M, Bhattacharyya A and Gangopadhyay G 2014 *Phys. Rev. C* **89** 065806 (Preprint 1403.0341)
- [114] Shen H, Toki H, Oyamatsu K and Sumiyoshi K 1998 *Progress of Theoretical Physics* **100** 1013–1031 (Preprint nucl-th/9806095)