

Large N matrix hyperspheres and the gauge-gravity correspondence

Mthokozisi Masuku, Mbavhalelo Mulokwe and João P. Rodrigues

*National Institute for Theoretical Physics,
School of Physics and Mandelstam Institute for Theoretical Physics,
University of the Witwatersrand, Johannesburg, Wits 2050, South Africa*
E-mail: Mthokozisi.Masuku@students.wits.ac.za,
Mbavhalelo.Mulokwe@students.wits.ac.za, Joao.Rodrigues@wits.ac.za

ABSTRACT: The large N dynamics of a subsector of $d = 0$ interacting complex multi matrix systems, which is naturally parametrized by a matrix valued radial coordinate, and which embodies the canonical AdS/CFT relationship between 't Hooft's coupling constant and radius, is obtained. Unlike the case of the single complex matrix, for two or more complex matrices a new repulsive logarithmic potential is present, and as a result the density of radial eigenvalues has support on an hyper annulus. For the single complex matrix, the integral over the angular degrees of freedom of the Yang-Mills interaction can be carried out exactly, and in the presence of an harmonic potential, the density of radial eigenvalues is shown to be of the Wigner type.

KEYWORDS: Matrix Models, $1/N$ Expansion, AdS-CFT Correspondence

ARXIV EPRINT: [1411.5786](https://arxiv.org/abs/1411.5786)

Contents

1	Introduction	1
2	Radial Sector	2
3	Large N configuration and hyperspheres	3
4	Angular degrees of freedom	6
5	Auxiliary two-matrix integral	7
6	Radial eigenvalues density in the Hoppe model	9
7	Summary	10

1 Introduction

The AdS/CFT correspondence [1–3] has provided new insights into the properties of several different gravitational objects from their corresponding description in terms of the large N limit of matrix valued (super) Yang-Mills theories. It is of great interest to understand how or if gravitational properties emerge from the large N limit of matrix theories, including in settings without supersymmetry or even conformal invariance. Of particular interest is the emergence of a semiclassical / geometric background in the large N limit of matrix theories at strong coupling.

Two defining relationships of the AdS/CFT correspondence in the canonical case are

$$R \sim \lambda^{\frac{1}{4}} \tag{1.1}$$

$$g_{\text{YM}}^2 \sim g_s \tag{1.2}$$

where R is the radius of both the AdS_5 and S^5 of the $AdS_5 \times S^5$ background, g_{YM} is the (super) Yang-Mills coupling, $\lambda = g_{\text{YM}}^2 N$ the 't Hooft coupling and g_s the Type IIB string coupling.

In this communication, we consider several properties associated with the large N limit of the integral

$$Z = \int [dX_I] e^{-S}, S = \frac{w^2}{2g_{\text{YM}}^2} \sum_{I=1}^6 \text{Tr} X_I^2 - \frac{1}{g_{\text{YM}}^2} \sum_{I \neq J}^6 \text{Tr}([X_I, X_J][X_I, X_J]) \tag{1.3}$$

which corresponds to the leading compactification (e.g., on $S^3 \times T^1$ or S^4) of the bosonic Higgs sector of $\mathcal{N} = 4$ SYM. The harmonic “frequency” w is a function of the curvature of

the manifold, and serves as an infrared regulator. g_{YM} “has been scaled out of the action”, so that the relationship (1.2) is apparent.

For this system, the relationship (1.1) now follows from dimensional analysis,¹ in that

$$[X_I] \sim \lambda^{\frac{1}{4}}. \tag{1.4}$$

If a suitable matrix “radial coordinate” can be identified, it will satisfy (1.1) simply by dimensional analysis, provided no other dimensional parameter is present. It would then be of great interest to obtain its large N dynamics.

For an even number of hermitian matrices (or an arbitrary number of complex matrices) a closed sub sector² dependent on a single matrix only, that has properties expected of such radial matrix, has indeed been identified [4, 5], and it is the purpose of this communication to discuss its large N dynamics.

In section 2, the radial sector, or the restriction of the system (1.3) to this closed subsector, is considered for an arbitrary number of complex matrices. The large N density of radial eigenvalues is obtained explicitly and shown to have a well defined strong coupling limit satisfying (1.1). A new feature present only in systems of strictly more than one complex matrix (as is relevant for $\mathcal{N} = 4$ SYM) is the appearance of a Penner [6–8] logarithmic potential that moves the eigenvalues away from the origin. In section 3, we integrate out the angular degrees of freedom in (1.3). This is carried out for a single complex matrix. In this case, the harmonic term is required, introducing another dimensionful parameter. By first mapping the system to a two matrix system, the large N limit of the radial eigenvalue distribution is obtained and shown to be of the Wigner type.

A review of the vast literature on matrix models in beyond of the scope of this brief communication. Aspects related to the emergence of geometries are reviewed in [9].

2 Radial Sector

We start by describing the closed subsector of systems of an even number of hermitian matrices which is naturally associated with a radial sector of the theory. First, we complexify by introducing complex matrices:

$$Z_1 = X_1 + iX_2 \quad , \quad Z_2 = X_3 + iX_4 \quad , \text{etc.}$$

and in general, for m complex matrices Z_A , $A = 1, \dots, m$, we consider the matrix,

$$\sum_{A=1}^m Z_A^\dagger Z_A. \tag{2.1}$$

This matrix is hermitian and positive definite, and its eigenvalues

$$\rho_i = r_i^2, \quad i = 1, \dots, N, \quad \rho_i \geq 0,$$

have a natural interpretation as matrix radial coordinates.

¹We extend to $d = 0$ the standard scalar dimensions $[X_I] = \frac{d-2}{2}$.

²In the sense of closure under Schwinger-Dyson equations.

The radially invariant sector of (1.3) is then:

$$S_R = \frac{w^2}{2g_{\text{YM}}^2} \text{Tr} \sum_{A=1}^m Z_A^\dagger Z_A + \frac{1}{2g_{\text{YM}}^2} \text{Tr} \left(\sum_{A=1}^m Z_A^\dagger Z_A \right)^2.$$

This sector has an enhanced $U(N)^{m+1}$ symmetry

$$Z_A \rightarrow V_A Z_A V^\dagger, \quad A = 1, \dots, m, \quad (2.2)$$

and depends only on the radial eigenvalues.

With a parametrization of the complex matrices $Z_A, A = 1, \dots, m$ in terms of a matrix valued radial matrix and $2m - 1$ unitary matrices,³ one can write

$$\int \prod_A \prod_{ij} dZ_A^\dagger{}_{ij} dZ_{Aij} = \int \prod_i d\rho_i \mathcal{J}(\rho_i) d[\text{Angular}]$$

The ‘‘angular’’ degrees of freedom can be integrated out in the radial sector.

$\mathcal{J}(\rho_i)$ has recently been obtained in closed form [5]. This results from the remarkable fact that correlators in this sector, with the enhanced symmetry (2.2), close under Schwinger-Dyson equations. The result is:

$$\begin{aligned} \mathcal{J}(\rho_i) &= C_m \prod_i d\rho_i \rho_i^{m-1} \prod_{i>j} \rho_i^{m-1} \rho_j^{m-1} (\rho_i - \rho_j)^2 \\ &= D_m \prod_i dr_i r_i^{2m-1} \prod_{i>j} r_i^{2m-2} r_j^{2m-2} (r_i^2 - r_j^2)^2 \\ &= C_m \prod_i d\rho_i \rho_i^{m-1} \Delta_{RM}^2(\rho_i) = D_m \prod_i dr_i r_i^{2m-1} \Delta_{RM}^2(r_i^2), \end{aligned} \quad (2.3)$$

where the antisymmetric product

$$\Delta_{RM}(\rho_i) \equiv \prod_{i>j} \rho_i^{\frac{m-1}{2}} \rho_j^{\frac{m-1}{2}} (\rho_i - \rho_j)$$

generalizes the well known Van der Monde determinant $\Delta = \prod_{i>j} (\rho_i - \rho_j)$, and C_m and D_m are numerical constants.

In the radial sector, we are then lead to the integral

$$Z = \int \prod_i d\rho_i \mathcal{J}(\rho_i) e^{-S_R(\rho_i)} = \int \prod_i d\rho_i e^{\ln \mathcal{J}(\rho_i) - S_R(\rho_i)} = \int \prod_i e^{-S_{\text{eff}}(\rho_i)}. \quad (2.4)$$

3 Large N configuration and hyperspheres

The large N configuration of (2.4) is given by the ‘‘saddle point’’ equation of $S_{\text{eff}}(\rho_i)$,

$$S_{\text{eff}}(\rho_i) = \frac{Nw^2}{2\lambda} \sum_i \rho_i + \frac{N}{2\lambda} \sum_i \rho_i^2 - N(m-1) \sum_i \ln \rho_i - \sum_{i \neq j} \ln |\rho_i - \rho_j| \quad (3.1)$$

³For an explicit such parametrizations in the $m = 1$ case, see [4].

and satisfies:

$$2 \sum_{j,j \neq i} \frac{1}{\rho_i - \rho_j} + \frac{N(m-1)}{\rho_i} = \frac{Nw^2}{2\lambda} + \frac{N}{\lambda} \rho_i \tag{3.2}$$

where $\lambda = g_{\text{YM}}^2 N$ is the 't Hooft's coupling. In terms of a density of eigenvalues $\Phi(\rho)$ normalized ($\Phi \rightarrow N\Phi$) so that $\int d\rho \Phi(\rho) = 1$,

$$\int_0^\infty \frac{d\rho' \Phi(\rho')}{\rho - \rho'} + \frac{(m-1)}{2\rho} = \frac{w^2}{4\lambda} + \frac{\rho}{2\lambda} \tag{3.3}$$

One observes that for $m \geq 2$, and in opposition to the single hermitean and single complex matrix case, there is, in addition to the repulsive logarithmic potential amongst the eigenvalues, a new single Penner [6–8] logarithmic potential which moves the eigenvalues away from the origin. The density of radial eigenvalues therefore is non-vanishing only between hyperspheres of radii $r_- = \sqrt{\rho_-}$ and $r_+ = \sqrt{\rho_+}$, i.e., within a “hyperannulus”. For $m = 1$, and in the strong coupling limit, the density of radial eigenvalues is non-vanishing within a hypersphere.

Inspection of either (3.2) or (3.3) establishes that the size of these hyperspheres in the strong coupling limit, i.e., when $w \rightarrow 0$, is set by the only parameter λ left in theory through a radius R satisfying

$$R^2 \sim \rho \sim \sqrt{\lambda}, \quad R \sim \lambda^{\frac{1}{4}} \tag{3.4}$$

Logarithmic Penner potentials [6–8] arise in several different contexts, such as in, for instance, the study of Gaussian ensembles of rectangular matrices [10–14].

They play a crucial role in the matrix model identified by Dijkgraaf and Vafa [15] as the “bridge” in the equivalence between partition functions of $d = 4$ $\mathcal{N} = 2$ quiver gauge theories and $d = 2$ conformal Toda field theories [16]. In all three cases, there is a Riemann surface involved: in the gauge theory it can be related to the Seiberg-Witten curve, in the Toda theory to the manifold in which the theory is defined, and in the quiver matrix model it is the spectral curve related to the resolvent of the model [17–19]. The positions of the poles of the derivate of the (sum of) Penner potentials correspond to insertions of vertex operators in the Toda/Liouville 2d theory.

The form of the density in (3.3) and values of ρ_{\pm} are obtained using standard methods [20], together with a careful treatment of the $\rho \rightarrow 0$ behaviour.

The solution is completely determined by the resolvent $G(z)$, defined on the complex plane:

$$G(z) = \int_{\rho_-}^{\rho_+} \frac{d\rho' \Phi(\rho')}{z - \rho'} = \frac{z}{2\lambda} - \frac{(m-1)}{2z} - \frac{q_0 + \frac{z}{\lambda}}{2z} \sqrt{(z - \rho_-)(z - \rho_+)}$$

For quiver matrix models, the singularities associated with the derivative of the Penner potentials have an interpretation as vertex insertions in Liouville theory, and remain poles of the resolvent. In our case, the Penner potential is thought of as one of the terms of the potential, and we follow Tan [21] in requiring the absence of a pole as $z \rightarrow 0$. This allows for a solution where ρ_- (and ρ_+) are positive for $m > 1$.

Absence of a pole as $z \rightarrow 0$ and the normalization of the density then require:

$$\begin{aligned}
 q_0 &= \frac{1}{2\lambda}(\rho_+ + \rho_-), & q_0\sqrt{\rho_+\rho_-} &= m - 1, \\
 -\frac{m-1}{2} + \frac{q_0}{4}(\rho_+ + \rho_-) + \frac{1}{16\lambda}(\rho_+ - \rho_-)^2 &= 1.
 \end{aligned}
 \tag{3.5}$$

With $s \equiv \rho_+ + \rho_-$ and $\Delta \equiv \rho_+ - \rho_-$ these can be solved to yield

$$\begin{aligned}
 s^2 &= \lambda \frac{4}{3}(m+1) \left[1 + \sqrt{1 + 3 \left(\frac{m-1}{m+1} \right)^2} \right] \\
 \Delta^2 &= s^2 - \frac{16\lambda^2(m-1)^2}{s^2}
 \end{aligned}
 \tag{3.6}$$

Using these expressions for s and Δ , one has, for instance:

$$\begin{aligned}
 \rho_- &= 0.416\lambda^{1/2}, & \rho_+ &= 3.10\lambda^{1/2}, & m &= 3 \\
 \rho_- &= 0.168\lambda^{1/2}, & \rho_+ &= 2.77\lambda^{1/2}, & m &= 2 \\
 \rho_- &= 0, & \rho_+ &= \frac{4}{3^{1/2}}\lambda^{1/2}, & m &= 1.
 \end{aligned}
 \tag{3.7}$$

In order to confirm the ρ prescription around the pole, particularly for the single complex matrix case ($m = 1$), while also providing a unified description for matrix systems with more than two complex matrices ($m \geq 2$), we extend the domain of definition of the density of eigenvalues to the real line. With $\rho = r^2$, $r > 0$, we define

$$2r\Phi(r^2) \equiv \phi(r) \equiv \phi(-r),
 \tag{3.8}$$

so that for an arbitrary function $f(r^2)$

$$\int_{-\infty}^{+\infty} dr f(r^2) \phi(r) = 2 \int_0^{+\infty} d\rho f(\rho) \Phi(\rho).
 \tag{3.9}$$

We remark that ($\rho = r^2$)

$$\begin{aligned}
 \int_0^{\infty} \frac{d\rho' \Phi(\rho')}{\rho - \rho'} &= \int_0^{\infty} \frac{2r' dr' \Phi(r'^2)}{r^2 - r'^2} = \frac{1}{2r} \int_0^{\infty} dr' \phi(r') \left(\frac{1}{r - r'} + \frac{1}{r + r'} \right) \\
 &= \frac{1}{2r} \int_{-\infty}^{\infty} \frac{dr' \phi(r')}{r - r'}
 \end{aligned}
 \tag{3.10}$$

As a result, (3.3) is equivalently written as

$$\int_{-\infty}^{\infty} \frac{dr' \phi(r')}{r - r'} = \frac{w^2 r}{2\lambda} + \frac{r^3}{\lambda} - \frac{(m-1)}{r}, \quad \int_{-\infty}^{+\infty} dr \phi(r) = 2.
 \tag{3.11}$$

The solution of (3.11) for $m > 2$ is symmetric, and it can be obtained following the analysis of [21, 22]. It is a two cut solution, with the cuts in the intervals $[-r_+, -r_-]$ and $[r_-, r_+]$, with $r_+ > r_- > 0$. One finds that $\rho_{\pm} = r_{\pm}^2$, with ρ_{\pm} given by equations (3.6) and that the densities are related by (3.8).

The solution of (3.11) when $m = 1$ is the standard symmetric single cut solution associated with a quartic interaction. It follows that equations (3.6) for the turning points indeed extend to $m = 1$ provided one sets $\rho_- = r_-^2 = 0$. Explicitly,

$$\pi\Phi(\rho) = \frac{1}{2\lambda\rho} \left(\rho + \frac{1}{2}(\rho_+ + \rho_-) \right) \sqrt{(\rho_+ - \rho)(\rho - \rho_-)}, \quad \rho_- \leq \rho \leq \rho_+$$

and

$$\pi\phi(r) = \frac{1}{\lambda r} \left(r^2 + \frac{1}{2}(r_+^2 + r_-^2) \right) \sqrt{(r_+^2 - r^2)(r^2 - r_-^2)}, \quad r_-^2 \leq r^2 \leq r_+^2,$$

with $\rho_{\pm} = r_{\pm}^2$ given by (3.6), (3.7).

4 Angular degrees of freedom

In the strong coupling limit, the scaling relationship (3.4) is unaffected by the angular degrees of freedom, as it follows from dimensional analysis.

For a single complex matrix ($m = 1$), the integration over the angular degrees of freedom in (1.3) can be carried out explicitly. Introducing matrix valued “polar coordinates”

$$X_1 + iX_2 = Z = RU, \quad Z^\dagger = U^\dagger R \tag{4.1}$$

with R hermitean and U unitary, R can be diagonalized as $R = V^\dagger r V$, with r a diagonal matrix and V unitary, we obtain [4, 23]

$$\int \prod_A \prod_{ij} dZ_{A^\dagger ij} dZ_{Aij} = C_{m=1} \int \prod_i d\rho_i \Delta^2(\rho_i) [dX][dS] \tag{4.2}$$

$$dX \equiv V dU U^\dagger V^\dagger, \quad dS \equiv dV V^\dagger$$

As a result of the properties of the de Haar measure, one has (the equality sign below is to be understood up to numerical constants),

$$\begin{aligned} Z &= \int [dX_I] \exp \left\{ -\frac{w^2}{2g_{\text{YM}}^2} \sum_{I=1}^2 \text{Tr} X_I^2 + \frac{1}{g_{\text{YM}}^2} \text{Tr}([X_1, X_2][X_1, X_2]) \right\} \\ &= \int [dZ^\dagger][dZ] \exp \left\{ -\frac{Nw^2}{2\lambda} \text{Tr} Z^\dagger Z - \frac{N}{2\lambda} (\text{Tr}((Z^\dagger Z)^2) - \text{Tr}(Z^{\dagger 2} Z^2)) \right\} \\ &= \int \prod_i d\rho_i \Delta^2(\rho_i) e^{-\frac{Nw^2}{2\lambda} \sum \rho_i - \frac{N}{2\lambda} \sum \rho_i^2} \int dU e^{\frac{N}{2\lambda} \text{Tr}(R^2 U^\dagger R^2 U)} \end{aligned} \tag{4.3}$$

$$= \int \prod_i d\rho_i \Delta^2(\rho_i) e^{-\frac{Nw^2}{2\lambda} \sum \rho_i - \frac{N}{2\lambda} \sum \rho_i^2} \exp \left\{ N^2 X(\rho_i, \rho_i, \frac{1}{2\lambda}) \right\} \tag{4.4}$$

The last integral is the well known Harish-Chandra- Itzykson-Zuber (HCIZ) integral [24], and it can be written in closed form in terms of a determinant. We recall that, in the large N limit, $X(\rho_i, \rho_i, \frac{1}{2\lambda})$ has an expansion in $1/\lambda$:

$$X \left(\rho_i, \rho_i, \frac{1}{2\lambda} \right) = \sum_{k=1}^{\infty} \frac{1}{k} \left(\frac{1}{2\lambda} \right)^k X_k(\rho_i, \rho_i), \tag{4.5}$$

where $X_k(\rho_i, \rho_i)$ is generically a function of the weights $1/N \sum_i \rho_i^n$, and is homogeneous of degree k in each of its variables. It follows that, in the strong coupling limit, should the integral be well defined, the size of the system remains determined by

$$R^2 \sim \rho \sim \sqrt{\lambda}, \quad R = \lambda^{\frac{1}{4}} \tag{4.6}$$

It is of particular interest to consider the leading $1/\lambda$ contribution in (4.5), as this is of the same order as the radially symmetric term in the action. With the leading term in the expansion (4.5) included, the effective action is now:

$$S_{\text{eff}}(\rho_i) = \frac{Nw^2}{2\lambda} \sum_i \rho_i + \frac{N}{2\lambda} \sum_i \rho_i^2 - \frac{1}{2\lambda} \left(\sum_i \rho_i \right)^2 - \sum_{i \neq j} \ln |\rho_i - \rho_j| \tag{4.7}$$

After rescaling and extending to the full real line, the saddle point equation now takes the form:

$$\int_{-\infty}^{\infty} \frac{dr' \phi(r')}{r - r'} = \frac{w^2 r}{2\lambda} + \frac{r^3}{\lambda} - \frac{rw_2}{\lambda}, \quad \int_{-\infty}^{+\infty} dr \phi(r) = 2, \quad \int_{-\infty}^{+\infty} dr r^2 \phi(r) = 2w_2 \tag{4.8}$$

The density of radial eigenvalues is readily obtained to be:

$$\begin{aligned} \pi\phi(r) &= \frac{r^2}{\lambda} \sqrt{r_+^2 - r^2}, \quad r_+ = 2\lambda^{\frac{1}{4}}, \quad -r_+ \leq r \leq r_+ \\ \pi\Phi(\rho) &= \frac{1}{2\lambda} \sqrt{\rho(\rho_+ - \rho)}, \quad \rho_+ = 4\lambda^{\frac{1}{2}}, \quad 0 \leq \rho \leq \rho_+ \end{aligned} \tag{4.9}$$

To move beyond a non-perturbative solution, we note that, as mentioned earlier, the integral can be written in closed form in terms of a determinant. Specifically,

$$Z = \text{Const.} \int \prod_i d\rho_i e^{-\left(\frac{Nw^2}{2\lambda} \sum_i \rho_i + \frac{N}{2\lambda} \sum_i \rho_i^2\right)} \det \left(e^{\frac{N}{2\lambda} \rho_i \rho_j} \right).$$

Note the cancellation of the Van der Monde determinant, i.e, the standard Coulombic-gas repulsion of the eigenvalues is absent. It turns out that this form of the partition function is not useful to obtain the large N distribution of eigenvalues. In order to facilitate this, we introduce an auxiliary two-matrix system in the next section.

5 Auxiliary two-matrix integral

In ρ -space,⁴ the partition function (4.3) can be written as

$$Z = \int [d\rho] \int dU e^{S_\rho} \tag{5.1}$$

$$= \int \prod_i d\rho_i \Delta^2(\rho) e^{-NV_\rho(\rho)} \int dU e^{\beta N \text{Tr} \rho U \rho U^\dagger}, \tag{5.2}$$

⁴In what follows, we will refer to the matrix integral expressed in terms of the matrices ρ and U as the matrix integral in ρ -space.

with

$$S_\rho = -NV_\rho(\rho) + \beta N \text{Tr} \rho U \rho U^\dagger. \quad (5.3)$$

Motivated by the appearance of a logarithmic potential for a larger number of matrices, we have in mind a potential of the form ($\epsilon > 0$):

$$V_\rho(\rho) = \frac{w^2}{2\lambda} \text{Tr} \rho + \frac{1}{2\lambda} \text{Tr} \rho^2 - \epsilon \text{Tr} \ln \rho. \quad (5.4)$$

For the saddle-point equation, we get

$$0 = -NV'_\rho(\rho) + \frac{\partial}{\partial \rho_i} \ln \Delta^2(\rho) + \beta N \langle U \rho U^\dagger + U^\dagger \rho U \rangle. \quad (5.5)$$

where $\langle . \rangle$ denotes the expectation value with respect to the angular degrees of freedom.

Let us now consider the auxiliary two (hermitean)-matrix integral defined via the action:

$$S = -N \text{Tr} V(A) - N \text{Tr} V(B) + N \text{Tr} A U B U^\dagger. \quad (5.6)$$

The partition function for this auxiliary two-matrix model reads

$$Z = \int dA \int dB \int dU e^{-N \text{Tr} V(A) - N \text{Tr} V(B) + N \text{Tr} A U B U^\dagger}, \quad (5.7)$$

where A and B are $N \times N$ hermitian matrices and dU is the Haar measure for the $U(N)$ unitary group.

In terms of the eigenvalues of the matrices A and B , which will be denoted by a_i and b_i ($i = 1, 2, \dots, N$), the partition function becomes

$$Z = \int \prod_i da_i \int \prod_i db_i \Delta^2(a) \Delta^2(b) \int dU e^{-N \sum_i V(a_i) - N \sum_i V(b_i) + \ln I(a,b)}, \quad (5.8)$$

where $\Delta^2(a)$ and $\Delta^2(b)$ are the Van der Monde determinants and $I(a,b)$ is the HCIZ integral:⁵

$$I(a,b) = \int dU e^{N \beta \text{Tr}(a U b U^\dagger)}. \quad (5.9)$$

At large N , the saddle-point equations are:

$$0 = -NV'(a_i) + \frac{\partial}{\partial a_i} \ln \Delta^2(a) + N \langle U b U^\dagger \rangle \quad (5.10)$$

$$0 = -NV'(b_i) + \frac{\partial}{\partial b_i} \ln \Delta^2(b) + N \langle U^\dagger a U \rangle. \quad (5.11)$$

Due to the exchange symmetry between A and B under the original integral (5.7), $a_i = b_i$ at the minimum. Adding (5.10) and (5.11), one obtains

$$0 = -NV'(a_i) + \frac{\partial}{\partial a_i} \ln \Delta^2(a) + \frac{N}{2} \langle U a U^\dagger + U^\dagger a U \rangle. \quad (5.12)$$

⁵Here, $a = \text{diag}(a_1, a_2, \dots, a_N)$ and $b = \text{diag}(b_1, b_2, \dots, b_N)$.

Comparison of (5.12) and (5.5) shows that the two saddle point equations agree if

$$2\beta = 1, \quad V'_\rho(\rho) = V'(A). \quad (5.13)$$

This is trivially achieved by rescaling $\rho \rightarrow \rho\lambda^{1/2}$. The action (5.3) becomes

$$S = -\frac{Nw^2}{2\lambda^{1/2}}\text{Tr}\rho - \frac{N}{2}\text{Tr}\rho^2 + N\epsilon\text{Tr}\ln\rho + \frac{N}{2}\text{Tr}\rho U\rho U^\dagger. \quad (5.14)$$

This action clearly satisfies the condition that $2\beta = 1$ - the other condition can be easily satisfied by setting $V'(A) = V'_\rho(\rho)$. Therefore, at the level of the saddle-point equations the matrix integral (in polar coordinates) with a Yang-Mills interaction is equivalent to an auxiliary matrix model with an action of the form

$$\begin{aligned} S = & -\frac{Nw^2}{2\lambda^{1/2}}\text{Tr}A - \frac{N}{2}\text{Tr}A^2 + N\epsilon\text{Tr}\ln A + N\text{Tr}AUBU^\dagger \\ & -\frac{Nw^2}{2\lambda^{1/2}}\text{Tr}B - \frac{N}{2}\text{Tr}B^2 + N\epsilon\text{Tr}\ln B. \end{aligned} \quad (5.15)$$

Methods to obtain the generating function $G(z)$, and hence the eigenvalue density, for these systems were developed in [25, 26] in the context of studies of the so called “induced QCD” [27, 28]. The techniques used are based on the Schwinger-Dyson equations of the system. Of particular interest is the limit $w \rightarrow 0$ while keeping ϵ small but finite, so that λ is the only dimensionful parameter, as considered in this communication until now. Indications are that in this limit, it may not be possible to obtain an eigenvalue density with support only on the positive real line in the presence of a pole at $z = 0$ [29].

In the following, we therefore set $\epsilon = 0$, and keep the w dependent harmonic potential. This is the Hoppe integral [30, 31].

6 Radial eigenvalues density in the Hoppe model

The action of the auxiliary system for the Hoppe integral in matrix valued “polar coordinates” (4.1) is then

$$\begin{aligned} S = & -\frac{Nw^2}{2\lambda^{1/2}}\text{Tr}A - \frac{N}{2}\text{Tr}A^2 + N\text{Tr}AUBU^\dagger \\ & -\frac{Nw^2}{2\lambda^{1/2}}\text{Tr}B - \frac{N}{2}\text{Tr}B^2. \\ = & \bar{w}\text{Tr}(A+B) - \frac{N}{2}\text{Tr}A^2 - \frac{N}{2}\text{Tr}B^2 + N\text{Tr}AUBU^\dagger \\ = & -2N\bar{w}^2\text{Tr}B - \frac{N}{2}\text{Tr}C^2 - N\bar{w}^2\text{Tr}C, \end{aligned} \quad (6.1)$$

where $C = A - B$ and $\bar{w}^2 = \frac{w^2}{2\lambda^{1/2}}$, and where the last equality is understood under the integral. Consequently, the partition function reads

$$\begin{aligned} Z = & \int dB \int dC e^{-2N\bar{w}^2\text{Tr}B - \frac{N}{2}\text{Tr}C^2 - N\bar{w}^2\text{Tr}C} \\ = & \text{Const.}(\bar{w}) \int \prod_i db_i e^{S_{\text{eff}}}, \end{aligned} \quad (6.2)$$

where the effective action describing the large N dynamics of the matrix B , and hence ρ , is given by

$$S_{\text{eff}} = \sum_{i \neq j} \ln |b_i - b_j| - 2N\bar{w}^2 \sum_i b_i. \quad (6.3)$$

The saddle-point equations are

$$\sum_{j, i \neq j} \frac{1}{b_i - b_j} = 2N\bar{w}^2. \quad (6.4)$$

In terms of the eigenvalue density $\Phi(b')$, the saddle-point equations read

$$\int_0^\infty \frac{db' \Phi(b')}{b - b'} = \bar{w}^2. \quad (6.5)$$

(Note that we have rescaled the eigenvalue density i.e. $\Phi(b) \rightarrow N\Phi(b)$.) Letting $b = r^2$ and, as explained before, extending to the full real line, the saddle-point equation can be written as

$$\int_{-\infty}^\infty \frac{dr' \phi(r')}{r - r'} = 2\bar{w}^2 r. \quad (6.6)$$

This is the well known integral equation for the Wigner distribution which, appropriately normalized, is:

$$\pi \phi(r) = 2\bar{w}^2 \sqrt{\frac{2}{\bar{w}^2} - r^2}, \quad 0 \leq r \leq \frac{\sqrt{2}}{\bar{w}}. \quad (6.7)$$

One can also solve the saddle-point equation in (6.5) directly, in which case one finds:

$$\pi \Phi(\rho) = \bar{w} \sqrt{\frac{2}{\rho} - \bar{w}^2}, \quad 0 \leq \rho \leq \frac{\sqrt{2}}{\bar{w}}. \quad (6.8)$$

In an approach based on commuting matrices, [32] have also obtained a radial Wigner distribution for the Hoppe model. However, given our definition of a radial coordinate, there is no approximation in our result.

7 Summary

Starting with the integral over systems of (an even number of) hermitean matrices interacting through a Yang-Mills potential, as is relevant to the Higgs sector of $\mathcal{N} = 4$ SYM, we identified a closed subsector that can naturally be interpreted as a radial subsector of the theory. For strictly more than two hermitean matrices, the measure introduces a logarithmic potential, a new result. The density of radial eigenvalues was obtained, and for strictly more than two hermitean matrices, was shown to have support between hyperspheres, and for two hermitean matrices, within a single hypersphere. For two hermitean matrices, the integral over the angular degrees of freedom was carried out explicitly, and in the presence of an harmonic potential, the density of radial eigenvalues was shown to be of the Wigner type, without approximation.

Acknowledgments

We would like to thank Robert de Mello Koch for interest in this project and for comments. This research is supported in part by the National Research Foundation of South Africa (Grant specific unique reference number UID 85974).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.M. Maldacena, *The Large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [[hep-th/9711200](#)] [[INSPIRE](#)].
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [[INSPIRE](#)].
- [3] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [[INSPIRE](#)].
- [4] M. Masuku and J.P. Rodrigues, *Laplacians in polar matrix coordinates and radial fermionization in higher dimensions*, *J. Math. Phys.* **52** (2011) 032302 [[arXiv:0911.2846](#)] [[INSPIRE](#)].
- [5] M. Masuku and J.P. Rodrigues, *How universal is the Wigner distribution?*, *J. Phys. A* **45** (2012) 085201 [[arXiv:1107.3681](#)] [[INSPIRE](#)].
- [6] R.C. Penner, *The moduli space of a punctured surface and perturbative series*, *Bull. Amer. Math. Soc.* **15** (1986) 73.
- [7] R.C. Penner, *Perturbative series and the moduli space of Riemann surfaces*, *J. Diff. Geom.* **27** (1988) 35 [[INSPIRE](#)].
- [8] J. Harer and D. Zagier, *The Euler characteristic of the moduli space of curves*, *Invent. Math.* **85** (1986) 457.
- [9] R. de Mello Koch and J. Murugan, *Emergent Spacetime*, [arXiv:0911.4817](#) [[INSPIRE](#)].
- [10] G.M. Cicuta, L. Molinari, E. Montaldi and F. Riva, *Large rectangular random matrices*, *J. Math. Phys.* **28** (1987) 1716 [[INSPIRE](#)].
- [11] A. Anderson, R.C. Myers and V. Periwal, *Complex random surfaces*, *Phys. Lett. B* **254** (1991) 89 [[INSPIRE](#)].
- [12] A. Anderson, R.C. Myers and V. Periwal, *Branched polymers from a double scaling limit of matrix models*, *Nucl. Phys. B* **360** (1991) 463 [[INSPIRE](#)].
- [13] R.C. Myers and V. Periwal, *Chiral noncritical strings*, *Nucl. Phys. B* **390** (1993) 716 [[hep-th/9210082](#)] [[INSPIRE](#)].
- [14] J. Feinberg and A. Zee, *Renormalizing rectangles and other topics in random matrix theory*, *J. Statist. Phys.* **87** (1997) 473 [[cond-mat/9609190](#)] [[INSPIRE](#)].
- [15] R. Dijkgraaf and C. Vafa, *Toda Theories, Matrix Models, Topological Strings and $N = 2$ Gauge Systems*, [arXiv:0909.2453](#) [[INSPIRE](#)].

- [16] L.F. Alday, D. Gaiotto and Y. Tachikawa, *Liouville Correlation Functions from Four-dimensional Gauge Theories*, *Lett. Math. Phys.* **91** (2010) 167 [[arXiv:0906.3219](#)] [[INSPIRE](#)].
- [17] H. Itoyama, K. Maruyoshi and T. Oota, *The Quiver Matrix Model and 2d-4d Conformal Connection*, *Prog. Theor. Phys.* **123** (2010) 957 [[arXiv:0911.4244](#)] [[INSPIRE](#)].
- [18] T. Eguchi and K. Maruyoshi, *Penner Type Matrix Model and Seiberg-Witten Theory*, *JHEP* **02** (2010) 022 [[arXiv:0911.4797](#)] [[INSPIRE](#)].
- [19] R. Schiappa and N. Wyllard, *An $A(r)$ threesome: matrix models, 2d CFTs and 4d $N = 2$ gauge theories*, *J. Math. Phys.* **51** (2010) 082304 [[arXiv:0911.5337](#)] [[INSPIRE](#)].
- [20] E. Brézin, C. Itzykson, G. Parisi and J.B. Zuber, *Planar Diagrams*, *Commun. Math. Phys.* **59** (1978) 35 [[INSPIRE](#)].
- [21] C.-I. Tan, *Generalized Penner models and multicritical behavior*, *Phys. Rev. D* **45** (1992) 2862 [[INSPIRE](#)].
- [22] R. de Mello Koch and J.P. Rodrigues, *The Collective field theory of a singular supersymmetric matrix model*, *Phys. Rev. D* **51** (1995) 5847 [[hep-th/9410012](#)] [[INSPIRE](#)].
- [23] M. Masuku, *Matrix Polar Coordinates*, MSc Thesis, Faculty of Science, University of the Witwatersrand (2009).
- [24] C. Itzykson and J.B. Zuber, *The Planar Approximation. 2.*, *J. Math. Phys.* **21** (1980) 411 [[INSPIRE](#)].
- [25] M.I. Dobroliubov, Yu. Makeenko and G.W. Semenoff, *Correlators of the Kazakov-Migdal model*, *Mod. Phys. Lett. A* **8** (1993) 2387 [[hep-th/9306037](#)] [[INSPIRE](#)].
- [26] Yu. Makeenko, *Some remarks about the two matrix Penner model and the Kazakov-Migdal model*, *Phys. Lett. B* **314** (1993) 197 [[hep-th/9306043](#)] [[INSPIRE](#)].
- [27] A.A. Migdal, *Exact solution of induced lattice gauge theory at large- N* , *Mod. Phys. Lett. A* **8** (1993) 359 [[hep-lat/9206002](#)] [[INSPIRE](#)].
- [28] V.A. Kazakov and A.A. Migdal, *Induced QCD at large- N* , *Nucl. Phys. B* **397** (1993) 214 [[hep-th/9206015](#)] [[INSPIRE](#)].
- [29] M. Mulokwe, *The Large- N Limit Of Matrix Models And AdS/CFT*, MSc Thesis, Faculty of Science, University of the Witwatersrand (2013).
- [30] J. Hoppe, *Quantum Theory of a Massless Relativistic Surface and a Two-Dimensional Bound State Problem*, Ph.D. Thesis, Department of Physics, Massachusetts Institute of Technology (1982).
- [31] V.A. Kazakov, I.K. Kostov and N.A. Nekrasov, *D particles, matrix integrals and KP hierarchy*, *Nucl. Phys. B* **557** (1999) 413 [[hep-th/9810035](#)] [[INSPIRE](#)].
- [32] D.E. Berenstein, M. Hanada and S.A. Hartnoll, *Multi-matrix models and emergent geometry*, *JHEP* **02** (2009) 010 [[arXiv:0805.4658](#)] [[INSPIRE](#)].