

UNIVERSITY OF SYDNEY

DOCTORAL THESIS

**Cosmological Implications of Quantum
Anomalies**

Author:
Neil D. Barrie

Supervisor:
Dr. Archil KOBAKHIDZE

*A thesis submitted in fulfilment of the requirements
for the degree of Doctor of Philosophy*

in the

University of Sydney High Energy Physics Group
School of Physics

November 2017

Declaration of Authorship

I, Neil David Barrie, declare that this thesis titled, ‘Cosmological Implications of Quantum Anomalies’ and the work presented in it are my own. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree at this University.
- Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated.
- Where I have consulted the published work of others, this is always clearly attributed.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work.
- I have acknowledged all main sources of help.
- Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Signed: Neil Barrie

Date: 25/07/2017

一期一会

UNIVERSITY OF SYDNEY

Abstract

Faculty of Science
School of Physics

Doctor of Philosophy

Cosmological Implications of Quantum Anomalies

by Neil D. BARRIE

The aim of this thesis is to investigate the possible implications of quantum anomalies in the early universe. We first consider a new class of natural inflation models based on scale invariance, imposed by the dilaton. In the classical limit, the general scalar potential necessarily contains a flat direction; this is lifted by quantum corrections. The effective potential is found to be linear in the inflaton field, yielding inflationary predictions consistent with observation.

A new mechanism forogenesis during inflation is presented, in which a new anomalous $U(1)_X$ gauge group is introduced. Anomaly terms source \mathcal{CP} and X violating processes during inflation, producing a non-zero CS number density that is distributed into baryonic and dark matter. The two $U(1)_X$ extensions considered in this general framework, gauged B and $B-L$ each containing an additional dark matter candidate, successfully reproduce the observed parameters.

We propose a reheating Baryogenesis scenario that utilises the Ratchet Mechanism. The model contains two scalars that interact via a derivative coupling; an inflaton consistent with the Starobinsky model, and a complex scalar baryon with a symmetric potential. The inflaton-scalar baryon system is found to act analogously to a forced pendulum, with driven motion near the end of reheating generating an η_B consistent with observation.

Finally, we argue that a lepton asymmetric $C\nu B$ develops gravitational instabilities related to the mixed gravity-lepton number anomaly. In the presence of this background, an effective CS term is induced which we investigate through two possible effects. Namely, birefringent propagation of gravitational waves, and the inducement of negative energy graviton modes in the high frequency regime. These lead to constraints on the allowed size of the lepton asymmetry.

These models demonstrate that a concerted approach in cosmology and particle physics is the way forward in exploring the mysteries of our universe.

Acknowledgements

First and foremost, I would like to thank my supervisor Archil Kobakhidze for introducing me to world of particle physics, and for giving me the tools to explore the field. I have thoroughly enjoyed my PhD studies, and have grown considerably as a researcher under your guidance. I look forward to continuing our work together into the future. I would also like to thank my Associate Supervisor Michael Schmidt for our many fruitful discussions over the course of my studies.

Thank you to all my collaborators for the opportunity to work with you, and for all the many insightful exchanges we have shared. I hope we shall continue to work together in the years to come. Special thanks to Kazuharu Bamba, Akio Sugamoto, Tatsu Takeuchi, and Kimiko Yamashita, for letting me join in their research pursuits. Also, a big thank you to Matthew Talia and Lei Wu for the many useful discussions over the course of our collaborations.

To the residents of Room 342, thank you for the many informative and humorous discussions. These last few years would not have been as enjoyable or as filled with laughter without you. Particular thanks go to my long time office mates - Adrian Manning, Carl Suster, Matthew Talia, Jason Yue.

I would also like to acknowledge Shennong for his amazing discovery 5000 years ago, which has made the last few years highly enjoyable and almost stress free.

To Sonia, thank you for always being so supportive, positive and understanding. You have helped me more than you know during the stressful, final half of my degree.

Finally, thank you to my family and friends for their ongoing support and help over the last few years, without which none of this could have been possible.

Contents

Declaration of Authorship	i
Abstract	iii
Acknowledgements	iv
Contents	v
1 Introduction	1
1.1 The Standard Model of Particle Physics	3
1.1.1 Formulation and Structure of the Standard Model	4
1.1.2 Symmetries in the Standard Model	7
1.1.3 Radiative Corrections and Quantum Anomaly Cancellation	9
1.2 The Standard Model of Cosmology and the Evolution of the Universe	12
1.2.1 The Friedmann-Robertson-Walker Metric	13
1.2.2 The History of the Universe	14
1.3 Mysteries to be Solved	16
1.3.1 The Inflationary Mechanism	16
1.3.2 The Origin of the Matter-Antimatter Asymmetry	20
1.3.3 Dark Matter	23
1.3.4 Neutrino Properties and the Cosmic Neutrino Background	25
1.3.5 Gravitational Waves	26
1.3.6 Hierarchy Problem in the Standard Model	26
1.4 Cosmological Implications of Quantum Anomalies	27
2 Scale Invariant Inflation	28
2.1 The Inflationary Epoch	29
2.2 Modelling the Inflationary Epoch	29
2.3 Scale Invariance, Weyl Transformations, and Non-minimal Couplings	33
2.4 Natural Inflation with Hidden Scale Invariance	34
2.4.1 Description of the Model	35
2.5 Quantum Corrected Potential in Curved Spacetime	37
2.6 Observational Signatures and Model Predictions	38
2.7 Conclusions and Future Prospects	40
3 An Asymmetric Universe from Inflation	41

3.1	The Matter-Antimatter Asymmetry	42
3.1.1	The Sakharov Conditions	42
3.1.2	Asymmetric Dark Matter	44
3.2	Topological Vacuum States and the Chern-Simons Number	44
3.2.1	Instanton and Sphaleron Transitions	45
3.3	A Model of Inflationary Cogenesis	47
3.3.1	Models with an Anomalous $U(1)_X$	48
3.4	Dynamics of an Anomalous Gauge Field During Inflation	51
3.5	Simultaneous Generation of Luminous and Dark Matter During Inflation	55
3.6	Replicating the Observed ρ_{DM}/ρ_B and η_B	57
3.7	Conclusions and Future Prospects	63
4	Baryogenesis During Reheating via the Ratchet Mechanism	64
4.1	The Reheating Epoch and Starobinsky Inflation	65
4.1.1	Starobinsky Inflation	66
4.2	Baryogenesis during Reheating via the Ratchet Mechanism	68
4.2.1	The Ratchet Mechanism	68
4.2.2	Description of the Model	68
4.3	Analytical Evaluation	71
4.3.1	Approximate Analytical Solution	74
4.3.2	Phase Locked States	76
4.4	The Generated Asymmetry Parameter	78
4.4.1	Conversion to Standard Model Particles	78
4.5	Conclusions and Future Prospects	79
5	Gravitational Waves and the Cosmic Neutrino Background	82
5.1	The Cosmic Neutrino Background	83
5.2	Gravitational Waves and the Graviton Propagator	85
5.2.1	Linearised Gravitational Waves	85
5.3	Graviton Polarisation Tensor in a Lepton Asymmetric $C\nu B$	86
5.4	Chern-Simons Modified Gravity and Observational Implications	90
5.4.1	Birefringent Propagation through an Asymmetric $C\nu B$	90
5.5	Induced Ghost-like Modes and Vacuum Decay	92
5.5.1	Photon Energy Spectrum from Induced Vacuum Decay	93
5.6	Conclusions and Future Prospects	96
6	Concluding Remarks and Outlook	98
A	Baryon and Lepton Number Anomalies in the Standard Model	101
A.1	Baryon Number Anomalies	101
A.2	Lepton Number Anomalies	102
A.3	Mixed Gauged Baryon and Lepton Number Anomalies	103
B	Further Details of Chapter 3 Calculations	104
B.1	F_+ Coefficients, Eq. (3.20)	104
B.2	F_- Coefficients, Eq. (3.21)	105

C Further Details of Chapter 5 Calculations	106
C.1 Dimensional Regularisation Integrals and Useful Relations	106
 Bibliography	 108

Chapter 1

Introduction

The Standard Models of Particle Physics and Cosmology have been highly successful at describing and reproducing the observed dynamics and properties of the Universe, but they are incomplete. Many mysteries regarding the workings of nature are yet to be resolved, for which new physics beyond the standard paradigms is required. Examples of these are the properties of neutrinos, the identity and origin of dark matter and dark energy, the origin of the matter-antimatter asymmetry, the inflationary mechanism, the quantum nature of gravity, the hierarchy problem, and more; each of which are indications that physics beyond the Standard Model exists. In the past few decades, many extensions to the Standard Model have been postulated in an attempt to explain and provide solutions to these problems. Many of these models have tried to solve the various problems of the Standard Model simultaneously. Any such extensions normally have many phenomenological implications, which can allow for the utilisation of a variety of tools to constrain the models. As of yet, none of these extensions have been accepted because they either are ruled out by experimental searches or current experiments are not sensitive enough to exclude their predictions.

It is no coincidence that the mysteries of the Standard Model of Particle Physics are predominantly associated with the very early universe and its evolution, rather this is the result of a strong intertwining of cosmological and particle dynamics. Despite the apparent divide between the cosmological and particle scales today, they are intimately connected, with potential discoveries in either field having large ramifications on the other. Therefore, to gain a deeper understanding of the primordial evolution of the universe it is imperative to understand the properties of the fundamental particles of nature, given that at very early times the microscopic dynamics of these particles directly dictated this evolution. The interconnectedness of these two fields is particularly relevant to modern day theoretical explorations given the current lack of strong evidence of beyond the Standard Model physics at the Large Hadron Collider LHC Run 2 and other terrestrial colliders. The ability for the LHC and near future terrestrial colliders

to discover new phenomena is limited by the energy scales that they can reach, while on the other hand cosmological observables have the potential to probe energies well beyond these experiments. Imprints and remnants of the dynamics of primordial high energy scale periods can potentially be seen in cosmological observables. An example of this is the Cosmic Microwave Background, which has illuminated many of the cosmological properties of our universe that we know today. Through considering a combination of observables from terrestrial collider searches and cosmological observables it may be possible to piece together the answers to many of the open questions of our universe.

This thesis will be structured, chronologically, as follows. In the remainder of this chapter we will briefly describe the Standard Models of Particle Physics and Cosmology before outlining the major questions facing these paradigms that will be addressed in this work, and the properties of anomalous symmetries in the Standard Model of Particle Physics. In Chapter 2, we will examine the inflationary epoch and a possible mechanism for inflation which also solves the hierarchy problem [1]. Chapters 3 and 4 present two possible models to explain the origin of the matter-antimatter asymmetry, the first acting during the inflationary epoch involving the introduction of a new gauge boson to the Standard Model [2, 3], and the other is a mechanism driven by the inflaton during reheating utilising the Ratchet Mechanism [4]. In Chapter 5, a novel way to utilise gravitational waves to illuminate the properties of the illusive Cosmic Neutrino Background will be considered [5]. Finally, in Chapter 6, we conclude with a discussion of the implications of each of these works and possible future paths for exploration. The focus of this thesis is on the cosmological implications of particle physics phenomena, but throughout my candidature we also conducted research into LHC Phenomenology that will not be discussed here [6-8].

1.1 The Standard Model of Particle Physics

The construction of the Standard Model of Particle Physics (SM) is one of the greatest achievements in modern science, the culmination of decades of theoretical and experimental endeavour by scientists from around the world with the goal of understanding the fundamental building blocks of nature; some of the foundational works and a review are listed [9–40]. The SM describes all of the known fundamental particles in nature and the interactions between them [41], apart from gravity which is still without a consistent quantum description. This includes all the fermions, comprising of quarks and leptons, and the gauge bosons which are the force carrying particles, as depicted in Figure 1.1. The SM is now complete, with the last piece of the puzzle being recently discovered, the Higgs boson. The Higgs particle was theoretically predicted decades ago to explain the origin of the masses of the SM particles [42–45], and was discovered at the Large Hadron Collider (LHC) [46] by the ATLAS [47, 48] and CMS [49, 50] collaborations.




FIGURE 1.1: The SM particle content, including masses, charges, and spins of each of the fermions and bosons.

1.1.1 Formulation and Structure of the Standard Model

The SM is a mathematical formulation that describes all the known fundamental particles of our universe, and their interactions which are mediated by the known forces - the strong, weak, and electromagnetic forces. It combines the half-integer spin fermions, including the leptons and quarks, and the integer spin gauge bosons, consisting of the photon, weak bosons, gluons, and Higgs particle. These are each described by fields, which are mathematical objects defined at every point in spacetime. It is built on the ideas of Quantum Field Theory, and the structure is fundamentally rooted in the ideas of symmetries and gauge invariance [51–58].

The SM is defined by the following direct product of groups $SU(3)_C \times SU(2)_L \times U(1)_Y$, with each group representing a gauge symmetry. The structure of the group determines the number and properties of the force carrying gauge bosons that couple to the associated charges. The SM is expressed in the form of a Lagrangian with all terms invariant under the gauge transformations described by these groups, which are local, spacetime dependent transformations. If the action is invariant under a given symmetry, then the physics derived from it will not change under such a transformation and the symmetry will have a corresponding conserved quantity. The SM gauge groups represent the following gauge bosons and forces,

- $SU(3)_C$: defines the strong force which is mediated by the gluons. These couple to fields that carry colour charge, which include the quarks and the gluons - known as Quantum Chromodynamics (QCD).
- $SU(2)_L$: the W^1 , W^2 and W^3 bosons mediate the weak force, with a linear combination of W^1 and W^2 forming the W^\pm , while W^3 makes up part of the Z boson and photon along with the hypercharge boson. Apart from self-couplings, the weak bosons couple only to the left-handed fermions, and as such maximally violate the \mathcal{C} and \mathcal{P} symmetry.
- $U(1)_Y$: the hypercharge boson is a linear combination of the Z boson, and the photon. The strength of the coupling to a given particle is defined by the hypercharge $Y = I_3 + Q$, where I_3 is the weak isospin and Q is the electric charge of the particle.

Each gauge group has a corresponding gauge field strength, $G_{\mu\nu}^a$, $W_{\mu\nu}^b$ and $B_{\mu\nu}$, with their respective gauge fields G_μ^a , W_μ^b and B_μ , which define the dynamics of the gauge bosons. The gauge field strength tensor for a non-Abelian gauge group is given by,

$$F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu + ig[A^\mu, A^\nu]. \quad (1.1)$$

The Higgs mechanism was proposed as a way for generating the masses of the SM particles, which involves spontaneous symmetry breaking [42–45, 59–62]. For an arbitrary scalar field

its potential may contain a minimum at which the field value is non-zero, if the scalar was to rest in this minima this would be known as a vacuum expectation value. This is made more interesting by the possibility that there are many degenerate such minima that are related by a continuous symmetry, which is broken by the scalar taking a vacuum expectation value. This is what the Higgs field undergoes in the Higgs mechanism. Applying the Higgs mechanism to the $SU(2)_L \times U(1)_Y$ portion of the SM leads to spontaneous symmetry breaking, that is, $SU(2)_L \times U(1)_Y$ is broken to a single $U(1)$ gauge group, which can be identified with the photon of electromagnetism. Given that the $U(1)$ symmetry is unbroken it represents a massless boson, the photon, and indicates that the other three bosons associated with the degrees of freedom of the original gauge group structure must now be massive. The photon is associated with generators from both groups of the unbroken electroweak part of the original SM gauge group structure, which contain linear combinations of the photon and Z boson generators that are resolved after spontaneous symmetry breaking. This illustrates the unification of the weak and electromagnetic forces. In the cosmological setting, this spontaneous symmetry breaking is related to the Electroweak Phase Transition (EWPT). Once the temperature gets below a certain critical point, the Higgs boson will take a non-zero vacuum expectation value, triggering spontaneous symmetry breaking in the SM when the temperature of the universe was about 100 GeV [63–66]. Above the EWPT, the weak gauge bosons are massless and the SM will be described by the unbroken gauge group structure.

Fermions	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	B	L
$Q_L^i = \begin{pmatrix} u \\ d \end{pmatrix}_L^i$	3	2	$\frac{1}{6}$	$\frac{1}{3}$	0
u_R^i	3	1	$\frac{2}{3}$	$\frac{1}{3}$	0
d_R^i	3	1	$-\frac{1}{3}$	$\frac{1}{3}$	0
$L^i = \begin{pmatrix} \nu \\ e \end{pmatrix}_L^i$	1	2	$-\frac{1}{2}$	0	1
e_R^i	1	1	-1	0	1

TABLE 1.1: The representations of the fermions, in reference to the SM gauge groups, including corresponding baryon and lepton numbers.

The fermions are described in singlet or doublet representations depending on whether they have a right or left chirality, respectively. Each generation of lepton and quark is represented by spinors, as defined by the Dirac equation. In Table 1.1, it is shown how the representation of each particle couples to each of the SM gauge groups $SU(3)_C$, $SU(2)_L$ and $U(1)_Y$.

The right-handed fermions are singlet fermions and do not couple to the weak bosons, unlike the left-handed fermion doublets. This defines the $SU(2)_L$ gauge symmetry as a chiral group. Only the quarks and gluons are charged under the $SU(3)_C$ group, with a triplet representation which corresponds to the three possible colours states. The $U(1)_Y$ gauge group of the SM describes a combination of the electromagnetic and weak Z gauge bosons. The coupling of this group to the fermions is defined as the hypercharge which is given by $Y = I^3 + Q$, where the $I^3 = \frac{1}{2}\tau^3$ is known as the weak isospin and is the third generator of the $SU(2)$ gauge group, giving the connection to the Z boson. The hypercharge of each fermion is given in Table 1.1.

The SM is defined by a Lagrangian density, which contains all the renormalisable terms that represent the possible field interactions, kinetic and mass terms. The Lagrangian describes the dynamics and evolution of each of the fundamental particles. The allowed terms in the Lagrangian are determined by the symmetries of the model, and how the fields transform under them, with the requirement that the action must be left unchanged under any transformations associated with the SM gauge symmetries, or assumed global symmetries. Imposing renormalisability requires the dimension of the operators in each term of the Lagrangian density to be ≤ 4 , such that the coefficients are dimensionless or have positive mass dimension. In any case, higher dimensional terms are generally suppressed at low energies by the mass scale of the coefficients. The SM Lagrangian density is given by,

$$\mathcal{L}_{SM} = \mathcal{L}_{Gauge} + \mathcal{L}_{Dirac} + \mathcal{L}_{Scalar} + \mathcal{L}_{Yukawa} , \quad (1.2)$$

where

$$\mathcal{L}_{Gauge} = -\frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} - \frac{1}{4}W_{\mu\nu}^b W^{b\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} , \quad (1.3)$$

describes the gauge fields and their related bosons,

$$\mathcal{L}_{Dirac} = i\bar{\psi}D_\mu\psi , \quad (1.4)$$

describes the free fermions, where ψ is any of the fermion representations within the theory,

$$\mathcal{L}_{Scalar} = (D_\mu\phi)^\dagger(D^\mu\phi) + m^2\phi^\dagger\phi - \frac{\lambda}{2}(\phi^\dagger\phi)^2 , \quad (1.5)$$

describes the Higgs kinetic term and potential,

$$\mathcal{L}_{Yukawa} = -y_{q_R}\bar{Q}_L\phi q_R - y_{e_R}\bar{L}\phi e_R + h.c. \quad (1.6)$$

describes each of the fermion interactions between the left-handed particles and their right-handed counterparts, via the Higgs boson. These terms generate the masses of the fermions, while the first term in Eq. (1.5) generate those for the W^\pm and Z bosons, once the Higgs

particle takes a non-zero vacuum expectation value. The covariant derivatives D_μ incorporate the interactions between the gauge fields and the SM fields.

It is possible the SM may not extend to high energies and rather that it is a low energy Effective Field Theory [67], which would indicate why it is unable to explain early universe phenomena. Meaning that it is valid below some energy scale, but breaks down above it, with new physics required; analogously to the divide between classical and quantum mechanics.

1.1.2 Symmetries in the Standard Model

The SM is built on the principles of gauge invariance through the description of the force carrying particles as vector bosons. This identification naturally leads to the requirement of gauge invariance associated with these vector fields, because without it the theory would be non-unitary and Lorentz violating, with the associated gauge bosons potentially having tachyonic degrees of freedom. These things would lead to an inconsistent and non-predictive theory. This means that the gauge symmetries are redundant, in that, the theory would not exist without their imposition. As such, these symmetries dictate what the allowed interactions are within the theory, or rather what terms can be contained within the Lagrangian. Gauge symmetries must also be protected from quantum anomalies - breakdowns in gauge invariance due to radiative corrections. They must be conserved at all orders of perturbation. Although, this is not necessarily required for global symmetries.

Classical global symmetries are characteristic symmetries of a theory at the tree level, and can be retained at all orders of perturbation. They can be present due to all the interactions allowed by the gauge symmetries respecting the symmetry accidentally, or they can be explicitly imposed on the theory. Each global symmetry in a theory has a corresponding charge which is conserved by the classical interactions. Although, global symmetries do not have to necessarily hold once quantum corrections are included. Symmetries broken by radiative corrections are known as anomalous symmetries. Gauge symmetries cannot be broken in this way because that would lead to a breakdown in the consistency of the theory through violations of gauge invariance.

Renormalisability requires that the mass dimension of each operator adds up to four or less, meaning that all coefficients must be dimensionless or have positive mass dimension. An accidental symmetry of the SM is one which is only present due to the requirement that the Lagrangian terms are renormalisable. Imposing this condition causes the removal of higher order terms from the Lagrangian that would have otherwise broken certain symmetries, subsequently retaining them as symmetries. Some examples of accidental symmetries of the SM Lagrangian are those associated with the global baryon number, and generational lepton numbers.

Global Baryon and Lepton Numbers

Baryon number is an accidental global symmetry of the SM. In the tree level SM, there are six possible $U(1)$ quark symmetries that correspond to each quark flavour, but are broken once the Cabbibo-Kobayashi-Maskawa (CKM) interactions are considered. This leads to there being only a single global $U(1)$ symmetry associated with the quarks, the total quark number, which can be identified with the baryon number symmetry of the SM. Each quark is assigned a baryon number charge of $\frac{1}{3}$ under this symmetry, as given in Table 1.1. The corresponding transformation is given by the following unitary transformations $q \rightarrow e^{i\frac{1}{3}\beta}q$, where β is a real number and q represents a quark field.

The generational lepton numbers are also accidental global symmetries of the SM. These are $U(1)$ symmetries given by the unitary transformations $l_i \rightarrow e^{i\alpha_i}l_i$, where α_i is a real number and i denotes the lepton generation. Although, the discovery of neutrino oscillations proved that these symmetries are violated. Instead the SM contains a single $U(1)$ associated with the total lepton number, as the oscillations still leave the total lepton number conserved [56]. Each of the leptons is assigned a lepton number equal to 1, as given in Table 1.1.

The corresponding Noether currents for the continuous global baryonic and leptonic symmetries are given by [68],

$$j_B^\mu \propto \bar{u}_i \gamma^\mu u^i + \bar{d}_i \gamma^\mu d^i \quad \text{and} \quad j_L^\mu \propto \bar{l}_i \gamma^\mu l^i + \bar{\nu}_i \gamma^\mu \nu^i, \quad (1.7)$$

where these lead to conservation of B and L at the tree level, $\partial_\mu j^\mu = 0$, and i denotes the generation of the fermion.

The $U(1)_B$ and $U(1)_L$ symmetries are present in the SM due to the imposition of renormalisability. There are non-renormalisable operators which can be written down that are consistent with the SM gauge symmetries, but explicitly break the baryon and lepton number symmetries. The lowest dimension operators that are examples of these are $\frac{QQQL}{\Lambda_B^2}$ and $\frac{LLHH}{\Lambda_L}$, respectively. These terms are strongly suppressed by their dimensional coefficients. For example, the first term gives a path for proton decay which, from experiment, has a decay time of $\tau_p > 10^{32-34}$ years, implying $\Lambda_B > 10^{15}$ GeV [69]. The high energy scales associated with these terms, Λ_B and Λ_L , leads to the suppression of these interactions at lower energies, but they could be indicative of new physics at those scales.

In this thesis, we are interested in symmetry violation by quantum anomalies, that is, induced by radiative corrections.

1.1.3 Radiative Corrections and Quantum Anomaly Cancellation

Anomalous symmetries are symmetries of the tree level SM that are violated once quantum corrections are included [70–72]. The SM requires radiative corrections of the tree level interactions so that precise predictions can be made, and so all the possible interactions can be determined and be taken into account. This can be done by perturbatively expanding the tree level vertices associated with interactions, in Feynman diagrams, into loops of a higher and higher order, where each subsequent expansion is suppressed relative to the previous one. These loop calculations give the quantum corrections to the corresponding interactions, and may even introduce new interactions that are not present at the tree level. Some examples of loop diagrams are presented in Figure 1.2, where in the cases presented, there is no tree level vertex with the same initial and final states, and hence such interactions are only possible with radiative corrections. These diagrams are also special, in that they may correspond to anomalous contributions to the associated baryon number symmetry, depending on the fermionic content of the theory.

Quantum anomalies can be induced for global symmetries, violating the conserved charges to which they correspond. The anomalies of a given symmetry are produced by the radiative corrections, the magnitude of which are found by considering what particles couple to the vertices of the expanded loop. This is determined by taking the trace of the vertex generators, doing so considers all the possible fermions that could traverse the loop, with the coefficient defining the magnitude. If zero, there is no anomaly present and the associated symmetries are preserved at that order of expansion. If non-zero, this gives you a correction to the Noether current of the corresponding global symmetry, which causes it to no longer be conserved, $\partial_\mu j^\mu \neq 0$. An example of this is shown in Figure 1.2, for the global baryon number current in the SM.

The gauge fields that couple to each vertex determine the possible fermions that can traverse the loop. For example, only left-handed fermions couple to the $SU(2)$ gauge field, so only q_L and l_L make up the possible constituents of the fermionic current in the corresponding loop corrections. There are similar constraints for $SU(3)$, that is, only the quarks Q_L , u_R and d_R are considered - while all SM fermions couple to the hypercharge gauge group $U(1)_Y$. Refer to Appendix A, for calculations of each of the 1-loop contributions for the baryon number and lepton number currents, as well as in the case that they are promoted to gauge symmetries.

Examples of anomalous symmetries in the SM are the global baryon and lepton number symmetries. In calculating the radiative corrections, the inclusion of a baryon number current means that only the quarks are considered in the fermionic loops. If one was to consider these as gauged symmetries, for example a gauged baryon number, then upon calculating the first order loop diagrams non-zero anomalies are found to be present for the $SU(2)^2 U(1)_B$ and $U(1)_Y^2 U(1)_B$ interactions. These vertex expansions are shown in Figure 1.2. This means that the baryon number symmetry has the following non-zero triangle anomalies:




FIGURE 1.2: The 1-loop anomalous radiative corrections involving a gauge boson associated with the $U(1)_B$ symmetry, (a) $SU(2)^2U(1)_B$, where the loop contains only left-handed quarks, and (b) $U(1)_Y^2U(1)_B$ where the loop contains only quarks.

For $SU(2)^2U(1)_B$, where all quarks contribute,

$$\mathcal{A}_2(SU(2)^2U(1)_B) = Tr[\tau^a \tau^b B] = \frac{n_g}{2} = \frac{3}{2}. \quad (1.8)$$

For $U(1)_Y^2U(1)_B$, where only left-handed quarks are considered,

$$\mathcal{A}_3(U(1)_Y^2U(1)_B) = Tr[YYB] = -\frac{n_g}{2} = -\frac{3}{2}. \quad (1.9)$$

where n_g is the number of quark generations, which is taken to be three.

It is found that the global baryon and lepton number symmetries have the same quantum anomalies when right handed neutrinos are included in the SM, but with $U(1)_L$ replacing $U(1)_B$ in Eq. (1.8) and Eq. (1.9). This indicates that $B - L$ is still a conserved global symmetry of the SM after quantum corrections are included.

The anomalous currents associated with the $U(1)_B$ and $U(1)_L$ currents source B and L violating processes, namely electroweak instanton transitions and sphaleron processes. These shall be discussed in Chapter 3, in the context of how this violation of B and L is utilised in Electroweak Baryogenesis, and how it can also be responsible for the washout of asymmetries generated early in the universes evolution.

The anomaly calculations of each of the quantum corrections for the $U(1)_B$ and $U(1)_L$ symmetries are given in Appendix A. These anomalies provide avenues for baryon and lepton number violation, as they are not strictly conserved quantities at the quantum level. If the $U(1)_B$ and $U(1)_L$ symmetries are introduced as local gauge symmetries these anomalies become gauge anomalies. This is a major issue for the validity of the theory and they must be cancelled, the methods of which we shall now discuss.

Gauge Anomaly Cancellation and the Green-Schwarz Mechanism

Quantum anomalies associated with a gauge symmetry must be cancelled to maintain the consistency of the theory. The reasons for this are two fold, firstly, they lead to a loss of unitarity of the S matrix and violation of Lorentz invariance, and secondly, they cause a breakdown in renormalisability. Each of which is required for a gauge invariant theory [73]. Such gauge anomalies can be cancelled in various ways through the inclusion of new degrees of freedom. Two examples of ways in which this can be done are the addition of new fermions, and the Green-Schwarz mechanism [74].

The inclusion of extra fermions can be used to cancel the anomalies induced by radiative corrections, as their inclusion leads to additional terms being present in the trace of the vertex generators. The quantum numbers of these new fermions can then be chosen such that the anomalies disappear. These can create new problems as they could mediate interactions which have not been observed at terrestrial colliders, or could form charged stable fields that could be detrimental to cosmological observables. Experimental investigations provide restrictions on the possible masses and couplings of any introduced particles, and the requirement of complete cancellation of the anomalies means that they can only carry certain quantum numbers.

The origin of the Green-Schwarz mechanism is in string theory applications, due to the appearance of unwanted gauge anomalies in such theories. To resolve this issue Green and Schwarz developed a method of restoring gauge invariance via the addition of new terms to the effective action [74], that lead to an effective field theory in which a new degree of freedom has been introduced to the theory. These new terms remove the anomalies by cancelling the non-invariance of the fermion measure in the action, leading to a gauge invariant theory that effectively describes the dynamics of the full theory at energies lower than a characteristic scale.

As an example of how this mechanism works, consider the inclusion of a new $U(1)_X$ symmetry to the SM, which has gauge anomalies. In this case, the Green-Schwarz terms added to the effective action contribute a new longitudinal degree of freedom to the introduced gauge field X_μ , which corresponds to the gauge field acquiring a mass. In doing so, we obtain an effective field theory, meaning that it can only sufficiently describe the interactions of the theory well below some new physics scale, which we shall denote f_X . This parametrises our ignorance of the full theory, which one could imagine contains heavy fermions that cancel the anomaly at scales well above f_X . For example, the full theory may involve the inclusion of leptoquarks at high energies, which can cancel the gauge anomalies present in the theory [75–77].

The new counter-terms added to the effective action are included in the Lagrangian density as follows,

$$\mathcal{L}_{counter} = -\frac{g_X^2}{16\pi^2} \mathcal{A} \theta X_{\mu\nu} \tilde{X}^{\mu\nu} , \quad (1.10)$$

where g_X is the coupling for the gauge field related to the anomaly, $X^{\mu\nu}$ is the field strength tensor, and \mathcal{A} is the associated anomaly coefficient. The θ component is pure gauge and is the longitudinal degree of freedom of the new gauge field X_μ . It is found that the tree level SM Lagrangian appears to not be gauge invariant once these counter-terms are added, although their variance cancels out against the gauge variation of the functional measure of fermion fields within the path integral quantization framework, leading to a gauge invariant theory. This term is reminiscent of the term for a massive gauge boson in the Stuekelberg formalism [78].

An example of mixed gauge anomalies is the result of promoting the gauged baryon number symmetry to a gauge symmetry, for which multiple anomalies are present. The following counter-terms must be included,

$$\mathcal{L}_{counter} = -\frac{3g_2^2}{32\pi^2}\theta(x)W_{\mu\nu}\tilde{W}^{\mu\nu} + \frac{3g_1^2}{32\pi^2}\theta(x)B_{\mu\nu}\tilde{B}^{\mu\nu} . \quad (1.11)$$

In this thesis, we shall be considering the cosmological implications of quantum anomalies such as these, and whether they can be utilised to explain various mysteries of the SM and SMC.

1.2 The Standard Model of Cosmology and the Evolution of the Universe

The Lambda Cold Dark Matter (Λ CDM) model is the current best model we have for describing the evolution of the universe from its beginning to the present day, and is considered the Standard Model of Cosmology (SMC) [79–82]. The SMC incorporates all known luminous matter, cold dark matter, and a cosmological constant to describe the dark energy density. The present day total energy content of the universe is split between these three main components - ordinary matter ($\sim 5\%$), dark matter ($\sim 27\%$), and dark energy ($\sim 68\%$). Observations of the Cosmic Microwave Background (CMB) in combination with other observational surveys provide data on the cosmological parameters of this model.

The measurement of the CMB has provided us a window into the early history of the universe [83–93]. Much of what we know about the universe comes from analysing the CMB and the temperature fluctuations that can be found within it. Knowledge of the properties of the universe prior to the formation of the CMB is difficult to glean due to the opaque nature of the primordial plasma prior to recombination. Despite this, the CMB can provide information from earlier times, such as inflation, allowing constraints to be placed on the properties of these prior epochs. In the case of inflation, this can be done through the spectral index and tensor to scalar ratio for which inflation makes predictions.




FIGURE 1.3: The Cosmic Microwave Background as measured by the PLANCK satellite [94].

The CMB is found to be highly isotropic and homogeneous on large scales, and as such, in the SMC the universe is considered to be static in conformal coordinates. Meaning that the universe evolves homogeneously and isotropically defined by a global scale factor $a(t)$. The gravitational metric associated with these evolutionary properties is known as the flat Friedmann-Robertson-Walker (FRW) universe, which is strongly supported by observations of the CMB. This is the basis of the understanding of the evolution of the universe, and shall be discussed below.

1.2.1 The Friedmann-Robertson-Walker Metric

One of the simplest models used to describe the evolution of the universe is the set of FRW cosmological models, which are a particular set of solutions of the Einstein equation that describe isotropic and homogeneous universes [95]. Isotropy means that they are spherically symmetric around a point in space and hence have no preferred direction, while homogeneity means that every point in space is equivalent, and hence there is no preferred position. This is supported by the remarkable uniformity of the CMB, observed by the WMAP [90] and PLANCK satellites [94]. Exact homogeneity would imply that there must also be no boundary or centre of the universe. For simplicity the universe is modelled as an isotropic and homogeneous spacetime.

This class of models includes three possible spatial geometries - closed, flat and open; each with constant curvature. A closed geometry is analogous to a sphere, which has constant positive curvature, flat to a plane, which has zero curvature, and open to a hyperbolic plane, which has constant negative curvature. For a general FRW universe, the metric in polar coordinates is given by,

$$ds^2 = dt^2 - a^2(t) \left(\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right), \quad (1.12)$$

where k defines the spatial curvature of the metric, and takes the values $k = 1, 0$, or -1 which correspond to closed, flat and open spaces, respectively. The scale factor $a(t)$ embodies the rate of expansion or compression of the universe.

From observation, we will be assuming that the universe is a simple flat FRW universe as the SMC considers, which has the following metric in Cartesian coordinates,

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2) , \quad (1.13)$$

or in conformal coordinates, which is defined by the following transformation of the time coordinate $dt = a(\tau)d\tau$,

$$ds^2 = a^2(t)(d\tau^2 - dx^2 + dy^2 + dz^2) , \quad (1.14)$$

and hence the metric can now be written as $g_{\mu\nu} = a(\tau)^2\eta_{\mu\nu}$, which is conformally flat. Therefore, this flat FRW metric is a suitable model for the spacetime of our universe and provides a simple framework in which to describe its evolution.

From the FRW metric can be derived the well-known Friedmann Equations, utilising the Einstein equation. They relate the nature of the expansion of the universe to the energy densities and curvature that is present, as described in Table 1.2. The Friedmann equations are as follows,

$$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi\rho}{3M_p^2} + \frac{\Lambda}{3} + \frac{k}{a^2} , \quad (1.15)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi}{3M_p^2}(\rho + 3p) + \frac{\Lambda}{3} . \quad (1.16)$$

where ρ and p are the mass density and pressure of the perfect fluid, respectively, k is the Gaussian curvature defined above, G is the gravitational constant, H is the Hubble rate [96], and Λ is the cosmological constant.

The evolution of the universe is defined by the properties of the scale factor, which is determined by the form of energy that is dominating. This is illustrated in the time evolution of the scale factors shown in Table 1.2, which are epoch dependent. In this table we use the dimensionless parameter w , known as the equation of state, to define each epoch. This parameter is related to the energy density by,

$$\rho \propto a^{-3(1+w)} . \quad (1.17)$$

1.2.2 The History of the Universe

The main epochs and events in the history of the universe, which we will be interested in, are as follows:

Inflation: Shortly after the Big Bang, or pre-inflationary state, the universe underwent a rapid period of expansion within a short period of time. This expansion is able to explain the homogeneity and flatness observed in the CMB, as well as the temperature fluctuations found

Type of Energy	w	$\rho(a)$	$a(t)$	$a(\tau)$
Matter Dominated	0	$\propto a^{-3}$	$\propto t^{2/3}$	$\propto \tau^2$
Radiation Dominated	$\frac{1}{3}$	$\propto a^{-4}$	$\propto t^{1/2}$	$\propto \tau$
Vacuum Dominated	-1	Constant	$\propto e^{Ht}$	$\propto \frac{-1}{H\tau}$

TABLE 1.2: Scale factors corresponding to each form of energy in a flat FRW universe.

in it. The exact nature of the mechanism leading to inflation is yet unknown, but it is thought to be caused by a particle known as the inflaton which dominates the energy density of the early universe. The induced period of inflation is a vacuum dominated era, during which the Hubble parameter is approximately constant and the scale factor is accelerating.

Reheating: After inflation the universe begins the transition to a radiation dominated epoch through a period known as reheating. The characteristics of the reheating epoch and the temperature of the resultant radiation dominated universe is determined by the properties of the inflationary mechanism and its interactions with the SM or mediator particles. The reheating temperature can be as high as $\sim 10^{15}$ GeV, for a high inflationary scale, and as low as ~ 1 MeV to not conflict with predictions from Big Bang Nucleosynthesis (BBN). The exact properties of reheating are complicated and are still not fully understood.

Baryogenesis: The universe has been observed to have an asymmetry between matter and antimatter, with minimal antimatter found to be present. The dynamics which lead to the origin of this asymmetry are generally assumed to occur at or above the electroweak scale, $T \sim 100$ GeV. This is highly dependent on the production mechanism, and could potentially happen as late as $T \sim 1$ MeV; once again, to not conflict with BBN.

Dark Matter Freeze-out: It is thought that the dark matter content of our universe was produced sometime during the radiation epoch. After being produced it would have thermally decoupled from the hot SM plasma once the temperature of it became low enough to suppress thermal production; assuming the dark matter interaction with the thermal plasma is small. As the interactions between the SM particles and dark matter are assumed to be fairly weak, this freeze out can occur fairly early on, meaning that the density is only affected by dilution due to the spatial expansion if the dark matter is stable relative to the age of the universe.

Electroweak Phase Transition: At $T \sim 100$ GeV the Higgs potential becomes such that the Higgs field takes a non-zero vacuum expectation value, leading to spontaneous symmetry breaking. As the temperature lowers, thermal fluctuations will cause certain regions to take a vacuum expectation value before others, leading to the formation of bubbles of true and false

vacuum. This phenomenon has been considered as the origin of the observed baryon asymmetry, with this mechanism being known as Electroweak Baryogenesis.

Neutrino Decoupling: As the neutrinos only couple via the weak interaction to the SM, they decouple from the plasma relatively early in the history of the universe, $T \sim 1$ MeV. The decoupled neutrinos make up the contents of the Cosmic Neutrino Background, which like the CMB, permeates the universe today.

Electron Positron Annihilation: At $T \sim 500$ keV the thermal production of $e^+ - e^-$ pairs freezes out, and the annihilation of these pairs, in the plasma, into photons will leave only the asymmetric part remaining. This annihilation increases the photon energy density of the universe.

Big Bang Nucleosynthesis: This is the period in which the nuclei observed in the universe today have their origin, through a series of well understood nuclear fusion reactions. This is well modelled by current nuclear physics models, and as such is very sensitive to new physics [97–107].

Recombination and Photon Decoupling: At a certain point in the evolution of the universe its temperature will be low enough that the electrons, and the nuclei generated in BBN, will begin to form atoms such as neutral hydrogen. As more nuclei form the universe becomes transparent to photons, which allows them to decouple from the plasma. This leads to the formation of the CMB, at the end of recombination, which we see today.

Dark Energy-Matter Equality: In more recent universal history, the dark energy density became larger than the matter energy density of the universe. This vacuum domination produces an accelerated expansion, which can be associated with a cosmological constant, and is the current state of the universe.

1.3 Mysteries to be Solved

Despite the successes of the SM and SMC in describing the nature and evolution of the universe, there are still many unsolved mysteries. In this thesis we hope to illuminate some of the following open problems.

1.3.1 The Inflationary Mechanism

The existence of an inflationary epoch was first postulated to solve unexplained observed phenomena and theoretical issues in cosmology [109–121]; namely, the flatness, horizon, and monopole problems [81, 122]. Since this time, significant amounts of evidence supporting a

Event	Time t	Redshift z	Temperature T
Inflation	10^{-34} s (?)	—	—
Baryogenesis	?	?	?
Dark matter freeze-out	?	?	?
Electroweak phase transition	20 ps	10^{15}	100 GeV
QCD phase transition	20 μ s	10^{12}	150 MeV
Neutrino decoupling	1 s	$6 \cdot 10^9$	1 MeV
Electron positron annihilation	6 s	$2 \cdot 10^9$	500 keV
Big Bang Nucleosynthesis	3 min	$4 \cdot 10^8$	100 keV
Matter-radiation equality	60 kyr	3400	0.75 eV
Recombination	260-380 kyr	1100 – 1400	0.26-0.33 eV
Photon decoupling	380 kyr	1000 – 1200	0.23-0.28 eV
Re-ionisation	100-400 Myr	11 – 30	2.4-7.0 meV
Dark energy-matter equality	9 Gyr	0.4	0.33 meV
Present	13.8 Gyr	0	0.24 meV

TABLE 1.3: The key events in cosmological history [108].

period of cosmic inflation has been obtained from a number of astrophysical observations [92], leading to it being a generally accepted part of the early universe evolution.

The flatness problem stems from the observed flatness of the universe, known to a very high precision, which appears to be fine-tuned. Rather than flatness being a special initial condition, inflation provides a natural way for such a lack of curvature to be the observed state of the universe. If one was to consider an initially curved spacetime, it would be approximately flat at very small scales. The rapid expansion of the universe produced by inflation will make these locally flat regions much larger, such that on large scales the universe will begin to appear flat. That is, the rapid spatial expansion quickly dilutes spatial curvature, pushing it towards zero. Hence the observable universe appearing flat is a natural consequence of inflation, if enough e-folds of expansion have occurred, resolving the problem.

A major mystery that inflation provides a solution for is the so called horizon problem. This is the observation that the universe is very homogeneous and appears to have been in thermal equilibrium, although if one is to trace back the evolution of the universe to the Big Bang it is found that all of today's causally disconnected regions would not have been in causal contact. This is because the universe has existed only for a finite period of time, so information can only have travelled a finite distance. This raises the question of why these causally disconnected regions are so well correlated. An inflationary epoch can provide the answer to this question by allowing the universe to have been in thermal equilibrium at very early times, prior to the onset of inflation. If the very early universe, just after the big bang, was in thermal equilibrium and then an inflationary epoch began, the resultant universe would be broken into many causally disconnected regions, after enough e-foldings, explaining what we observe today. This idea is depicted in Figure 1.4.




FIGURE 1.4: Illustration of the solution of the horizon problem provided by inflation [122].

A generic prediction of many Grand Unified Theories is the production of topological defects, such as monopoles, at phase transitions which occur as the universe cools from the very high temperatures at the end of reheating. The dynamics of the early universe would lead to the thermal production of a high density of monopoles. Unfortunately, it is found that the energy density of these monopoles in the universe would be too large to be consistent with current observations. This is known as the monopole problem [123–125]. Now if one is to postulate an inflationary epoch, this would lead to the significant dilution of the density of these relics,

removing them as a potential issue, assuming the reheating temperature is low enough to prevent their overproduction after inflation.

Another attractive property of the inflationary scenario is that it can generate the density perturbations observed via the inhomogeneities of the CMB. These density perturbations provided the seeds for large scale structure formation, and hence led to the formation of the galaxies we see today. As inflation proceeds, quantum fluctuations of the inflaton field, the scalar field which induces the inflationary epoch, are enlarged by the rapid spatial expansion, producing the temperature anisotropies we see in the CMB [126].

Once inflation ends the initial energy densities of matter have been diluted to negligible quantities, so the universe must be reheated. This is also taken into account in the inflationary mechanism, through the decay of the inflaton after inflation ends; this period is known as the reheating epoch. The product of the reheating epoch, is a radiation dominated universe with a thermalised plasma with characteristic temperature T_{rh} , after which the standard big bang cosmology proceeds.

Unfortunately, despite the many successes of the inflationary paradigm at explaining cosmological evolution, the exact mechanism for inflation is still not known. It is usually assumed that this epoch is the result of a scalar field, named the inflaton, which slowly rolls towards the minimum of its potential [122, 127–132]. The potential energy of this scalar field dominates the energy density of the universe, as long as the inflaton only rolls slowly in its potential. This almost constant potential energy, dominating at early times, leads to an effective vacuum energy dominated universe; a de Sitter space time. Such a vacuum energy dominated universe is characterised by exponential expansion, as discussed above. It should be noted that cosmological models other than inflation have been proposed to solve the issues described above, such as String gas and bounce cosmologies, with varying success [133–135].

Many inflaton candidates have been proposed since inflation was first postulated, with most involving the introduction of a scalar(s), but as of yet none have been experimentally verified [115, 126, 136–157]. Due to this, many attempts to constrain and test inflationary mechanisms have been considered [158–166]. One reason this is difficult is that with the current experimental sensitivity to inflationary observables, many models give degenerate solutions and hence are indistinguishable. This has led to recent work into identifying classes of inflationary models [167–170]. Another issue is that the inflaton itself is unlikely to be produced terrestrially due to it generically having a very high mass, well beyond the range of any current or future collider experiments. In Chapter 2, we will consider a new class of scale invariant inflationary scenarios which may have interesting implications for particle physics.

1.3.2 The Origin of the Matter-Antimatter Asymmetry

One of the major questions in modern physics is how the observed matter-antimatter asymmetry of the universe developed [171–184]. Antimatter was first predicted to be on equal footing with matter [185], but observational and experimental results suggest that this is incorrect. Terrestrial experimental investigations confirmed this in decays of K^0 ($d\bar{s}$) [186, 187], D^0 ($c\bar{u}$) [188] and B^0 ($s\bar{b}$) [189, 190] mesons, which have provided evidence for \mathcal{C} and \mathcal{CP} violation. As an asymmetry is observed, astronomical observations provide strong contradictory evidence to the equivalence of matter and antimatter [90, 94, 191, 192].

Observations indicate that the visible universe is dominated by matter and not antimatter. This baryon number density is determined by analysing the Baryon Acoustic Oscillations (BAO) measured from the CMB power spectrum, depicted in Figure 1.5, and the temperature fluctuations in the CMB [90, 92, 193]. BBN theory also accurately predicts the abundances of the primordial light elements and is sensitive to the size of the baryon asymmetry; providing constraints on it [104, 194–197]. The observed value of the matter-antimatter asymmetry of our Universe is determined from the combination of these measurements, and is parametrised by the baryon to entropy density ratio,

$$\eta_B = \frac{n_B - n_{\bar{B}}}{s} \simeq \frac{n_B}{s} \simeq 8.5 \cdot 10^{-11}, \quad (1.18)$$

where n_B ($n_{\bar{B}}$) is the baryon (antibaryon) number density, and s is the entropy density of the universe. It is possible that the universe may have begun with a net baryon number prior to inflation, but any baryon number density would be diluted away by the rapid expansion of inflation, hence it is assumed that it must be produced dynamically. The process of dynamical generation of the observed matter-antimatter asymmetry is known as Baryogenesis.

It has been proposed that the net baryon number of the universe is indeed zero and that many unmixed islands of matter and antimatter exist [198–200]. If separate sectors do exist it would be possible to observe annihilations at the boundaries between the regions. Electron-positron pairs involved in annihilation processes would produce high energy photons, which if in significant numbers can cause a skewing of the CMB spectrum. These processes can also heat the ambient plasma leading to an additional indirect spectral distortion. This would be observable as dilutions or perturbations in the CMB [201]. These gamma ray sources are not observed, and the CMB is found to be highly uniform, so to be consistent with observation the voids between these matter and antimatter sectors would be required to be large. However, the size of the possible voids between these sectors is also constrained by the CMB. Regions large enough to survive recombination would be observable in the CMB, but these have not been found; excluding the idea that there are sectors of matter and antimatter separated by voids within the observable universe. It is possible that we live in a matter island that is larger than the observable universe, but this is not supported experimentally [202–204].




FIGURE 1.5: The baryon acoustic oscillations observed in the CMB power spectrum, as measured by the PLANCK satellite [193].

Seeing as there is overwhelming evidence for the existence of the baryon asymmetry in the universe, we must consider how such an asymmetry could be produced.

The Sakharov Conditions

The Sakharov conditions [205], formulated by A.D. Sakharov in 1967, are the requirements for successful Baryogenesis in the early universe. The conditions are,

- **Baryon number violation** If immediately after the Big Bang there was zero net baryon number (B) and B is strictly conserved, then the net baryon number density of the universe would remain zero for all time. Therefore, B number violating processes are needed.
- **C and CP violation** The baryon number violating processes are required to violate the C and CP symmetries, such that either the matter or antimatter process is favoured.
- **A period of non-equilibrium** The processes that violate B , C and CP must occur in a period of non-equilibrium, so that the reverse reactions do not washout any generated asymmetry.

Any mechanism that wishes to produce a charge asymmetry in the early universe must satisfy these criteria, unless the theory violates CPT .

Baryogenesis in the Standard Model

One of the most studied mechanisms for the generation of the baryon asymmetry is Electroweak Baryogenesis. The reason for this is that the first model discovered in this class of Baryogenesis mechanisms was the SM itself, which contains all ingredients for satisfying the Sakharov conditions [206–217]. They are satisfied in the following ways:

- **Baryon number violation** The quantum anomalies associated with the global baryon number symmetry and the electroweak gauge group indicate that it is not a strictly conserved quantity within the SM. Through these anomalies, baryon number violation can occur via non-perturbative electroweak sphaleron transitions [72], which shall be discussed further in Chapter 3.
- **\mathcal{C} and \mathcal{CP} violation** The \mathcal{C} and \mathcal{P} symmetries are maximally violated in the SM by the chiral nature of the weak interactions. There is also \mathcal{CP} violation provided by a complex phase present in the CKM matrix.
- **A period of non-equilibrium** The electroweak phase transition (EWPT) is the push out-of-equilibrium utilised in SM Baryogenesis [65, 218, 219]. Once the universe cools, thermal fluctuations in the primordial plasma can lead to regions falling below the EWPT earlier than others; the EWPT occurs at a characteristic temperature $T \sim 100$ GeV. This leads to the formation of ‘bubbles’, or regions, of broken phase, where the Higgs boson has acquired a vacuum expectation value. The expansion of the walls of these bubbles creates the non-equilibrium conditions required for Baryogenesis. This process is depicted in Figure 1.6, where the \mathcal{CP} and B violating processes associated with the left handed fermions lead to the accumulation of a net baryon number in the interior of the expanding bubble.




FIGURE 1.6: A schematic of the Electroweak Baryogenesis mechanism [220].

Although the SM satisfies the Sakharov conditions, the resulting prediction for the baryon asymmetry generated is found to be $\eta_B^{sm} \simeq 10^{-18}$, approximately eight orders of magnitude lower than the observed value [221]. One reason for this discrepancy is that the strength of the \mathcal{CP} violation is too weak to produce a large enough asymmetry. Also, the Higgs mass is too high, $m_h \sim 125$ GeV, for the EWPT to be strongly first order, so the departure from equilibrium is too small [48, 217, 222, 223]. Therefore, the SM is not sufficient to explain the observed baryon asymmetry of the universe, and the existence of new physics is required.

Other Proposed Scenarios

Many models have been proposed to solve the issue of the baryon asymmetry of the universe (a selection of reviews are given here Ref. [224–235]), many of which are simple extensions of the SM [236–239]. Some of these models also try to resolve other unexplained physical phenomena alongside the generation of the baryon asymmetry, such as dark matter [240–242]. As of yet most of these models remain unproven because the related phenomena are beyond the reach of current experiments.

A very attractive paradigm for the generation of the matter-antimatter asymmetry is Leptogenesis [243–249]. In these models the asymmetry is initially generated in the neutrino sector, through extensions to the SM that also explain the origin of the neutrino masses. Once the asymmetry is generated the $B + L$ violating electroweak sphalerons redistribute some of the net lepton number into the baryonic sector, producing what we see today. The interest in this form of scenario is driven by the lack of a complete understanding of the neutrino sector, which could potentially be hiding key information to solving this and other mysteries in particle physics.

Extensions that have the baryon asymmetry of the universe generated before the inflationary epoch risk having the accumulated baryon number being completely diluted. The difficulty with producing the asymmetry during inflation [250–252] is the requirement that the rate of asymmetry production must be greater than the rate of dilution. If not, the generated asymmetry will be quickly diluted away. For this reason the majority of postulated mechanisms occur after reheating. In this work we will be considering two possible extensions of the SM that can explain the baryon asymmetry of the universe; one acting during inflation [2, 3] and the other during reheating [4].

1.3.3 Dark Matter

The existence of dark matter, an abundant form of non-luminous matter in the universe, was first proposed by Zwicky, in 1933, to explain the discrepancy between the expected and observed luminosity of galaxies, given their measured gravitational masses [253]. Further evidence for

dark matter was provided by the observation of anomalous galactic rotation curves [254–260], which to be resolved with theoretical predictions of the rotation profile required the existence of more luminous matter than could be seen [261, 262]. Since this time the existence of dark matter has been generally accepted as many more pieces of observational evidence have been gathered; these include gravitational lensing, the BAO, and its importance for cosmological evolution and structure formation, seeing as it is a key component of the Λ CDM cosmological model. Despite this, the true identity and nature of the constituents of dark matter are not known.

One of the most researched candidates for dark matter are Weakly Interacting Massive Particles (WIMPs) [263–265]. Originally, this model was very strongly favoured because of the so called WIMP miracle; a coincidence in which, if the dark matter candidate has a mass of 100 GeV and couples via the weak force to the SM, then its interaction cross section would be consistent with the observed dark matter abundance. The interaction cross section was also below the direct detection measurements at that time. The simplest WIMP miracle paradigm has now been ruled out by direct detection experiments such as those currently being undertaken at XENON [266] and LUX [267], which means other candidates must be explored. Many possibilities for the identity and properties of dark matter have been proposed [268–272], but as of yet none have been found. Neutrinos were once considered possible candidates for dark matter [273], but their masses and density are too low, and their relativistic nature early in the universe would have hindered structure formation. Non-particle explanations for dark matter, which constitute alternate gravity theories, have also been proposed, the most well-known being Modified Newtonian Dynamics [274], but these have been mostly unsuccessful in explaining all of the observational phenomena.

The dark matter particles are considered to be stable, to the extent that their lifetime is longer than the age of the universe, because if they decay too quickly they will fail to facilitate successful structure formation. It is also possible that there could be various stable particles present in non-zero densities in the universe, making up components of the dark matter density [275]. A current issue with the cold dark matter model is that simulations predict more satellite galaxies to be present than are observed, which could be a result of small structure formation suppression by dark matter which is warm, rather than cold as usually defined. If the dark matter carries more kinetic energy it can more easily escape gravitational potentials, and hence contribute to the washing out of small scale structures.

One of the most attractive paradigms for describing dark matter is asymmetric dark matter, that is there is a matter-antimatter asymmetry in the dark matter sector too, which may have a similar origin to that in the baryonic sector. This is motivated by the measurement of the

ratio of the energy densities of luminous and non-luminous matter,

$$\rho_{\text{DM}} \simeq 5.5\rho_B . \quad (1.19)$$

The similarity in these observed densities could indicate a connection between the dynamics and cosmological evolution of visible and dark matter. Many models attempting to explain this ratio have been proposed, and it is an active area of research [237–239, 276–299]. We will consider the possibility of an inflationary cogenesis scenario in Chapter 3.

1.3.4 Neutrino Properties and the Cosmic Neutrino Background

The true nature of neutrinos has been an intriguing mystery since they were first postulated due to their weakly interacting nature. There have been major advances in our understanding of the properties of neutrinos in the last couple of decades due to the ever improving sensitivity of experiments [300]. The Super-Kamiokande [301–306], Sudbury Neutrino Observatory (SNO) [307–309] and other experiments [310–312], provided the first experimental evidence of neutrino mass and furthered our knowledge of their interaction properties. More recent experiments have begun determining the neutrino mixing angles, and are getting closer to obtaining a measurement of the neutrino \mathcal{CP} violating phase.

One of the main questions regarding neutrinos at present is what are and is the origin of their masses; the mass hierarchy of the neutrino generations is also still unknown. Neutrinos can have either a Dirac mass term $m_\nu \bar{\nu}_L \nu_R$ or a Majorana mass term $m_\nu \nu_L^c \nu_L$, each of which have interesting phenomenological implications. Particularly the Majorana case, because of the lepton violating nature of the mass term, which can not only induce double beta decay, but can have implications for Baryogenesis. It is possible that there is also other neutrino species, such as sterile neutrinos which do not couple to any of the SM gauge fields, that can have interesting cosmological implications [313].

Another feature of neutrinos which makes them important for early universe cosmology, is that, much like photons and the production of the CMB, there is an analogous relic neutrino background; the Cosmic Neutrino Background ($C\nu\text{B}$). This background is produced much earlier than the CMB, and as such could have information encoded within that could unlock many of the current mysteries in particle physics and cosmology. The information encoded within the $C\nu\text{B}$ could help us gain a greater understanding of the neutrino sector, and also help illuminate the many open questions regarding the history of the universe. Unfortunately, the low temperature and density of this background today means it is unlikely we will ever be able to directly observe it. There is still hope that it could be indirectly observed, and this is what we shall be exploring in Chapter 5, through the possible effects the $C\nu\text{B}$ could have on gravitational waves.

1.3.5 Gravitational Waves

A determination of the fundamental description of gravity would be revolutionary in physics and have many wide reaching implications. The recent observation of gravitational waves by the LIGO collaboration [314] signals the beginning of the new era of gravitational wave astronomy, allowing the opening of a new window into the fundamental workings of gravity and the universe. To date, the LIGO collaboration has reported the measurement of gravitational waves from several binary black hole merger events [315, 316]. These observations have allowed constraints to be put on potential extensions to general relativity [317–320], and possibly in future, other areas of astrophysics through combined analyses, for example with ANTARES and IceCube [321]. This is a major step towards gaining a greater understanding of the workings of gravity. Future gravitational wave detectors, such as eLISA [322, 323], will also be able to differentiate the polarisations of incoming signals. This will provide information beyond the amplitude and waveform of the waves, and will enable a deeper analysis of the gravitational wave source as well as the fundamental workings of gravity itself. This could be achieved through identification of birefringent propagation effects, which could be smoking guns for certain extensions of General Relativity.

An exciting aspect of such searches is the possible implications for particle physics and early cosmological evolution. Unlike light, for which the early universe plasma was opaque prior to the CMB, it is possible that the gravitational wave remnants from events prior to the formation of the CMB could be observable by future detectors. These observations could provide constraints on early cosmology, and the particle dynamics at that time, that are not achievable with current measurement tools. An example of this shall be discussed in this work with reference to the Cosmic Neutrino Background [5], which could shed light on the true nature of neutrinos and also provide information concerning the origin of the matter-antimatter asymmetry.

1.3.6 Hierarchy Problem in the Standard Model

The hierarchy problem is associated with the unnaturalness of the difference between the apparent fundamental scales of nature, with regards to the SM and gravity, namely the weak and Planck scales [324–331]. This has led to much theoretical anxiety due to the level of fine tuning required to replicate observables if there are no new fundamental scales between the weak and Planck scales. This issue is illustrated by the predicted mass of the Higgs boson, for which the quantum corrections contain quadratic divergences dependent on the scale of new physics. The, recently discovered, Higgs mass has been found to be much lighter than the Planck scale, confirming this disparity in scales.

Supersymmetry has been one of the most studied beyond the SM theories because it can provide a potentially natural solution to the hierarchy problem [332]. The lack of discovery of the Supersymmetry partners has led to the consideration of increasingly complicated Supersymmetry models, which themselves contain issues with fine tuning.

Another solution to the hierarchy problem, which has gained increased interest in recent times, is the introduction of a conformal symmetry into the SM [333]. This model postulates a theory which has no fundamental scales, and that any observed scale is generated dynamically by the spontaneous breaking of the conformal symmetry. The Lagrangian of such a theory contains no dimensionful couplings, and as such has no mass terms. Seeing as the conformal symmetry removes fundamental scales it is a novel way of solving the hierarchy problem. Scale invariant theories have been studied extensively in the past, and have experienced a recent resurgence in interest, and have various interesting implications for the cosmological evolution of the universe [334–364]. In this work we shall consider the application of scale invariance to the inflationary epoch [1].

1.4 Cosmological Implications of Quantum Anomalies

In the following chapters we shall explore the implications of quantum anomalies in the evolutionary history of the universe. Firstly, in Chapter 2, we consider an inflationary mechanism within the setting of a scale invariant theory, providing both a mechanism for inflation and a solution to the hierarchy problem [1]. In Chapter 3, we formulate a generalised version of the inflationary mechanism for Baryogenesis we introduced in [2], which can now produce the correct baryon number density as well as dark matter; though the addition of an anomalous $U(1)_X$ and dark matter fermion to the SM [3]. Chapter 4, considers a new mechanism for Baryogenesis during the reheating epoch, in which we introduce a scalar inflaton and complex scalar baryon which are derivatively coupled [4]. In Chapter 5, we consider a new way to illuminate the properties of the neutrino sector through attempting to constrain the lepton asymmetry carried by the $C\nu B$ due to effects it can induce with respect to gravitational wave propagation and gravitational instabilities [5].

Chapter 2

Scale Invariant Inflation

The period of inflation in the early universe is a well-established paradigm in standard cosmology due to its success at solving various observational problems, as well as providing measurable predictions. It was first introduced to solve observational problems associated with the standard big bang cosmology, and is now a well-established theory with many models having been proposed and significant effort expended in the pursuit of experimental verification. Although it is generally agreed that there was an epoch of inflation prior to BBN, the exact mechanism that leads to the accelerated expansion is still unknown. A problem which makes this more difficult is the degeneracy of the predictions of many models, and the insufficient sensitivity in current measurements of inflationary scenario observables.

The most common approach to inducing this inflationary scenario is the introduction of a scalar field which acts as the inflaton, whose domination of the early universe's energy density, and traversal of its almost flat potential, leads to the exponential expansion. In this chapter we explore a new class of natural inflation mechanisms which exhibit scale invariance via the dilaton, and involve an arbitrary number of scalar fields that are non-minimally coupled to gravity. The scale invariance of the theory assures the flatness of the inflationary potential, which is then lifted by small quantum corrections that violate the conformal symmetry. The breaking of the associated scale symmetry can also provide an origin for the apparent hierarchy of scales observed in nature [1].

2.1 The Inflationary Epoch

Cosmic inflation is an attractive paradigm that resolves some outstanding puzzles of the standard hot Big Bang cosmology, such as the horizon, flatness, and monopole problems [109–121]. In addition, it provides a natural mechanism for the generation of nearly scale-invariant inhomogeneities through the quantum fluctuations of the inflaton field, that at later stages result in the observed large scale structure of the universe [126]. Observations of the CMB and large scale structure provide strong support for cosmic inflation [365].




FIGURE 2.1: The CMB as measured by the PLANCK satellite [92].

Many mechanisms for the inducement of the inflationary epoch have been proposed since the paradigm's inception. The majority of these ideas centre on the existence of a scalar field that is homogeneous and isotropic, and dominates the energy density of the universe, thus causing exponential expansion [115, 126, 136–156]. Given the rapid increase in the number of models proposed, work has recently gone into identifying classes of inflationary models [167–170]. This is in part due to the degeneracy of the simplest predictions of observable parameters between different models [158–166, 366].

There are also issues relating to the consistency of the inflationary theory, due to many models requiring the inflaton to take superplanckian values during its evolution [160, 367, 368]. Higher order operators, which are suppressed by the Planck scale, start to contribute significantly for large variations of the inflaton field during inflation. The effective field theory approximation, which favours $|\varphi| \ll M_p$, breaks down in such cases, and inflationary predictions may become unreliable [369, 370].

2.2 Modelling the Inflationary Epoch

The simplest mechanisms of inflation are proposed to be related to the evolution of a scalar field, known as the inflaton field [122, 127–132]. As an illustrative example, we consider a simple

version of this mechanism known as chaotic slow roll inflation, which is depicted in Figure 2.2. This model assumes that at the beginning of the universe the inflaton field is in an unstable vacuum state. Due to the quantum fluctuations of the inflaton field, it will eventually begin to roll down the potential well towards the true vacuum. As the inflaton field rolls slowly down the potential, inflation is occurring; this requires that the field is rolling slower than the rate of expansion. This ensures that as the inflaton rolls down the potential, the potential energy density remains approximately constant leading to a vacuum dominated epoch. Therefore, from the Friedman equations, the scale factor during the inflationary period, expressed in conformal coordinates, is,

$$a(\tau) \propto -\frac{1}{H\tau} , \quad (2.1)$$

where $\tau \in [-\infty, 0]$ is the conformal time, with $\tau \rightarrow 0$ as inflation proceeds. During this epoch the density of the other forms of energy quickly dilute away. At the end of inflation there is a period known as reheating, prior to the radiation epoch, in which the initial potential energy of the inflaton field is converted into SM particles, producing the matter energy densities we see today [371]. The properties of the reheating epoch will be discussed further in Chapter 4.




FIGURE 2.2: Illustration of the features of the chaotic slow roll inflationary mechanism [372].

We now want to describe the evolution of the inflaton field, so that we can determine the conditions required for inflation to occur, and the possible observable predictions we can obtain from such a mechanism. The inflaton is a scalar field described by the following Lagrangian,

$$\mathcal{L} = -\partial_\mu \phi \partial^\mu \phi - V(\phi) , \quad (2.2)$$

where ϕ is the inflaton field, and $V(\phi)$ is the inflationary potential, which could be of the following form,

$$V(\phi) = m^2 \phi^2 + \phi^4 \sum_{n=0}^{\infty} \lambda_n \left(\frac{\phi}{M_P} \right)^n . \quad (2.3)$$

We shall now consider the inflaton in a flat FRW universe. In this spacetime background the inflaton has the following equation of motion [373],

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0 , \quad (2.4)$$

where H is the Hubble parameter which defines the expansion rate, and $V'(\phi)$ is the first derivative with respect to ϕ of the general inflaton potential. This has the corresponding Friedmann equation,

$$H^2 = \frac{3}{M_p^2} \left(\frac{1}{2} \dot{\phi}^2 + V(\phi) \right) , \quad (2.5)$$

where $M_p = 1/\sqrt{8\pi G}$ is the reduced Planck mass. The inflaton field must slowly rolling down its potential for long enough such that the required amount of spatial expansion is achieved solving the known cosmological problems. In order for this to happen the following conditions need to be satisfied. Firstly, the inflaton's potential energy must dominate the kinetic energy of the field, such that its evolution is gradual, which means,

$$\dot{\phi}^2 \ll V(\phi) , \quad (2.6)$$

The other condition, is that the field's acceleration should be small, allowing a sufficient length of time for slow rolling and hence the inflationary setting to progress. That is,

$$\ddot{\phi} \ll |3H\dot{\phi}|, |V'(\phi)| . \quad (2.7)$$

These requirements allow the definition of the well-known slow-roll parameters, which must be much less than one to ensure the inflaton is slowly rolling and inflation is occurring. The slow roll parameters are as follows,

$$\epsilon = \frac{M_p^2}{2} \left(\frac{V'(\phi)}{V(\phi)} \right)^2 = \frac{1}{2M_p^2} \frac{\dot{\phi}^2}{H^2} , \quad (2.8)$$

$$\eta = M_p^2 \frac{V''(\phi)}{V(\phi)} = -\frac{\ddot{\phi}}{H\dot{\phi}} . \quad (2.9)$$

where these are the leading order slow roll parameters. The ability of a given scenario to successfully lead to an inflationary epoch requires the smallness of the slow-roll parameters, which is equivalent to flatness in the inflaton potential. These constraints ensure that we have an epoch of effective vacuum domination, acting like a cosmological constant. The smaller ϵ and η are, the longer the possible duration of inflation; with the end of inflation, and onset of reheating, occurring when these slow roll parameters are violated. The potential of the ϕ field can now be chosen to see whether it is consistent with an inflationary setting.

During the period of inflation, quantum fluctuations of the inflaton are inflated to produce the temperature anisotropies observed in the CMB today. The level of inhomogeneity produced in the inflationary epoch is related to the properties of the inflationary potential through the slow roll parameters. From the slow roll parameters the power spectrum of the observed scalar perturbations, P_s , the tensor-to-scalar ratio, r , and the spectral index n_s can be determined,

and compared with observation. These measurable model predictions are given by,

$$P_s = \frac{1}{24\pi^2 M_p^4} \frac{V(\phi_\star)}{\epsilon_\star}, \quad (2.10)$$

$$r = 16\epsilon_\star, \quad (2.11)$$

$$n_s = 1 - 6\epsilon_\star + 2\eta_\star. \quad (2.12)$$

All the quantities with subscript ‘ \star ’ in the above equations are evaluated at a field value $\phi = \phi_\star$ that corresponds to the field value after a number of e-folds of inflation, N_\star . The number of inflationary e-folds can be calculated approximately using the following relation,

$$N_\star \simeq \frac{1}{M_p} \int_0^{\varphi_\star} \frac{d\varphi}{\sqrt{2\epsilon}}. \quad (2.13)$$

In this analysis we have considered a single scalar field, but it is also possible to consider inflationary scenarios which contain multiple scalar fields, each of which could potentially contribute to the inflationary expansion [374–387]. In multifield inflation, the number of e-folds is a function of each of the scalar fields in the theory. One scalar may dominate and hence lead to an effective single field inflationary scenario, or you may get a mixture of the scalars contributing to produce a flat direction. Analogously, slow roll parameters and inflationary predictions of a similar form can be derived. In this chapter, we will be considering such a possibility.

The inflationary epoch has also been postulated to solve various other problems in particle physics, this shall be explored in Chapter 3 in relation to Baryogenesis and dark matter. These models can utilise the de Sitter nature of the inflationary setting, the associated reheating epoch, or the inflaton and its decay [388–391]. The possibility of using the reheating epoch for Baryogenesis will be considered in Chapter 4.

A class of natural inflation models has been suggested as a symmetry-motivated solution to the hierarchy problem in particle physics [137]. The inflaton in this class of models is a pseudo-Goldstone boson of a spontaneously broken anomalous global symmetry. The flatness of the inflaton’s potential is guaranteed by an approximate shift symmetry in the Lagrangian, meaning that the inflationary potential can easily satisfy the slow roll parameters, and hence can support an inflationary setting. However, the most recent observational results suggest that the simplest models of natural inflation are now disfavoured at 95% confidence level [365]. This type of model is of interest to the work considered in the rest of this chapter.

2.3 Scale Invariance, Weyl Transformations, and Non-minimal Couplings

Before considering the new inflationary mechanism that we will discuss in this chapter, we must first introduce the concept of scale invariance, within the context of a scalar Lagrangian including gravity, and the associated Weyl transformation. In some earlier works the existence of a symmetry associated with scale invariance was advocated as a possible explanation for the hierarchy problem without fine-tuning [328, 330]. This idea has received renewed attention in recent times due to the non-observation of Supersymmetry [392, 393]. The existence of a scale invariance symmetry in nature would have many ramifications for particle physics and cosmology. One of the interesting applications of the scale invariance symmetry is to the inflationary scenario, which has been considered more recently in the literature [354, 356, 358–360, 364, 394–396, 396–403].

A scale invariant theory is one where there are no physical scales, which is manifest in the classical action through the absence of dimensionful couplings. In this scenario any scales we observe must be dynamically generated, such as through a scalar taking a vacuum expectation value. This would be induced through quantum corrections, which break the scale invariance symmetry explicitly, as the scale invariance symmetry is typically anomalous.

An example of a scale invariant action that includes gravity is,

$$S = \int dx^4 \sqrt{-g} \left(\xi s^2 R - \frac{1}{2} \partial_\mu s \partial^\mu s - V(s) \right), \quad (2.14)$$

where we have a SM singlet scalar field s with a scale invariant potential $V(s)$, and ξ is the coupling of s to gravity.

Non-minimal coupling of a scalar field to gravity has been widely discussed in the literature [404, 405]. This idea can lead to many interesting implications for cosmology, particularly for inflation [406–409] and cosmological evolution [410]. The idea of a non-minimal coupling to gravity has been utilised to salvage the idea of Higgs inflation, reconciling the properties of the Higgs boson with the observational constraints of the associated inflationary scenario. Although such a solution requires the Higgs to have a large non-minimal coupling to gravity, namely $\xi \sim \mathcal{O}(10^5)$ [411].

The Einstein frame, that containing the usual Einstein-Hilbert term, can be transformed to the Jordan frame through the consideration of an extra scalar degree of freedom [412]. In the Jordan frame, the Einstein-Hilbert term is not present, and is replaced by a $\xi s^2 R$ term, as in the action in Eq. (2.14). The coefficient in front of this new term defines the nature of the scalar's coupling to gravity. In the case that $\xi = -1/12$, this is known as the conformal coupling, meaning that

the gravitational coupling is of the usual form. When making the Weyl transformation from this frame to the Einstein frame, this scalar is found to not have physical implications and is a fictitious degree of freedom. If instead $\xi \neq -1/12$, then this is known as a non-minimal coupling, and corresponds to an alteration to the gravitational-matter field interactions in the Einstein frame. The advantage of transforming to the Einstein frame from the Jordan frame, in our model, is the added simplicity of inflationary calculations in the Einstein frame.

We shall now exhibit the properties of the Weyl rescaling utilising the action given in Eq. (2.14). This action is expressed in the Jordan frame, in which there is no Einstein-Hilbert term. In order to obtain the Einstein frame we must make a conformal transformation, or Weyl rescaling, which is defined as,

$$g_{\mu\nu}^E = \Omega(s)^2 g_{\mu\nu} , \quad \text{where} \quad \Omega(s)^2 = \frac{\xi s^2}{\bar{M}_p^2} , \quad (2.15)$$

where $g_{\mu\nu}^E$ is the metric in the Einstein frame, and \bar{M}_p is the reduced Planck mass. In order for the action to remain invariant under the Weyl rescaling, the scalar and fermion fields must also be transformed as follows,

$$\sigma_E = \frac{\sigma}{\Omega(s)} \quad \text{and} \quad \psi_E = \frac{\psi}{\Omega(s)^{3/2}} , \quad (2.16)$$

while gauge vector fields are invariant under the transformation.

After undertaking this transformation the usual Einstein-Hilbert term is obtained, and the scalar potential is now,

$$V_E(s) = \frac{V_J(s)}{\Omega(s)^4} . \quad (2.17)$$

The scale invariance symmetry can be introduced into a theory through the introduction of a dilaton χ , which is the scalar field associated with the dilatation current. By scaling any coupling with dimension n by $\left(\frac{\chi}{f}\right)^n$ we can remove all the scales in a theory leaving only dimensionless couplings, where f is identified as the dilaton decay constant. The scales we see today would then be generated dynamically by the dilaton when it takes a vacuum expectation value, which breaks the scale invariance symmetry spontaneously. This is how the scale invariance symmetry shall be realised in our inflationary mechanism.

2.4 Natural Inflation with Hidden Scale Invariance

The main focus of this chapter is on a new class of natural inflation models that we have proposed, which are based on a hidden scale invariance. This is realised through the introduction of a spontaneously broken anomalous scaling symmetry with a corresponding pseudo-Goldstone boson, that we identify as the dilaton. We begin by considering a very generic scale-invariant

model, with an arbitrary number of scalar fields and all allowed scalar interaction operators in the scalar potential, with dimensionful couplings removed through appropriate rescaling by the dilaton. In this scenario there is found to always exist a direction in the field space that is absolutely flat in the classical limit. This bodes well for successful inflation given the ease with which the slow roll parameters can be satisfied by this model. Inflation proceeds along this direction, while the other fields in this parametrisation reside in their respective (meta)stable minima. Although, the flat direction must have a non-negligible slope, such that the inflation epoch is not eternal. Upon quantum corrections, the scale invariance of the model is broken, leading to the flat direction being lifted. Now we shall have a more detailed discussion of the workings of this model, and derive the observational predictions it produces.

2.4.1 Description of the Model

Consider a multifield inflationary scenario in which we have N scalar fields, $\{\phi_i\}$ (where $i = 1, 2, \dots, N$), each with a general non-minimal coupling to gravity, denoted ξ_i . Now take a Wilsonian effective field theory that describes the SM, or its extension, that is coupled to gravity at an ultraviolet scale we shall call Λ ,

$$S_\Lambda = \int dx^4 \sqrt{-g} \left[\left(\frac{M_p^2}{2} + \sum_{i=1}^N \xi_i(\Lambda) \phi_i^2 \right) R - \frac{1}{2} \sum_{i=1}^N \partial_\mu \phi_i \partial^\mu \phi_i - V(\phi_i) + \dots \right], \quad (2.18)$$

where $M_p \approx 2.4 \cdot 10^{18}$ GeV and we use the mostly positive signature for the metric tensor. Here we have displayed only the scalar sector, which includes the SM Higgs boson. The scalar potential $V(\phi_i)$ is a generic polynomial of the scalar fields $\{\phi_i\}$ respecting the relevant symmetries of the theory,

$$V(\phi_i) = \sum_{\{i_n\}} \lambda_{i_1, \dots, i_n}(\Lambda) \phi_{i_1} \dots \phi_{i_n}, \quad (2.19)$$

where $\lambda_{i_1, \dots, i_n}(\Lambda)$ is a coupling of mass dimension $(4 - n)$ defined at the Wilsonian cut-off Λ , while $\xi_i(\Lambda)$ is a dimensionless non-minimal coupling of the scalar field ϕ_i to gravity. The scale invariance is explicitly broken in Eq. (2.18) by the ultraviolet cut-off Λ , the Einstein-Hilbert term $\sim M_p^2 R$ and dimensionful couplings $\lambda_{i_1, \dots, i_n}$ ($n \neq 4$).

Now we propose that the underlying theory exhibits a scale invariance, which in the effective low-energy theory is implemented via a (non-linear) pseudo-Goldstone boson, that is the dilaton χ . A simple way to incorporate the dilaton field χ is to rescale all of the dimensionful parameters in Eq. (2.18) by the respective powers of χ/f , where f is the dilaton ‘‘decay constant’’. More

specifically, the following transformations are taken in the action given above,

$$\Lambda \rightarrow \Lambda \frac{\chi}{f} \equiv \lambda\chi, \quad M_p^2 \rightarrow M_p^2 \left(\frac{\chi}{f} \right)^2 \equiv \xi\chi^2, \quad (2.20)$$

$$\lambda_{i_1, \dots, i_n}(\Lambda) \rightarrow \lambda_{i_1, \dots, i_n}(\Lambda\chi/f) \left(\frac{\chi}{f} \right)^{4-n} \equiv \sigma_{i_1, \dots, i_n}(\lambda\chi)\chi^{4-n}. \quad (2.21)$$

Thus, instead of Eq. (2.18), we consider the transformed action,

$$S_{\lambda\chi} = \int dx^4 \sqrt{-g} \left[\left(\xi\chi^2 + \sum_{i=1}^N \xi_i(\lambda\chi)\phi_i^2 \right) R - \frac{1}{2} \partial_\mu \chi \partial^\mu \chi - \frac{1}{2} \sum_{i=1}^N \partial_\mu \phi_i \partial^\mu \phi_i - V(\phi_i, \chi) + \dots \right], \quad (2.22)$$

where the scalar potential is now given by,

$$V(\phi_i, \chi) = \sum_{\{i_n\}} \sigma_{i_1, \dots, i_n}(\lambda\chi) \chi^{(4-n)} \phi_{i_1} \dots \phi_{i_n}. \quad (2.23)$$

This action is manifestly scale invariant in the classical limit. As we shall see, this scale invariance is broken at the quantum level through the renormalisation group (RG) running of the dimensionless couplings, i.e., $\frac{\partial \sigma_{i_1, \dots, i_n}}{\partial \chi} \neq 0$, etc.

In the calculations that follow, it is convenient to use a ‘hyperspherical’ representation for the set of scalar fields $\{\phi_i, \chi\}$,

$$\begin{aligned} \phi_i &= \rho \cos(\theta_i) \prod_{k=1}^{i-1} \sin(\theta_k), \quad (i = 1, 2, \dots, N), \\ \chi &= \rho \prod_{k=1}^N \sin(\theta_k). \end{aligned} \quad (2.24)$$

Expressing the action in Eq. (2.22) through the ‘hyperspherical’ representation of the fields, we observe that the modulus field ρ factors out. That is, the first term in the action and the scalar potential presented in Eq. (2.22) can be written as $\sim \rho^2 \zeta(\theta_i) R$ and $\sim \rho^4 U(\theta_i)$, respectively, in which,

$$\zeta(\theta_i) = \xi(\lambda\chi) \prod_{k=1}^N \sin^2(\theta_k) + \sum_{i=1}^N \xi_i(\lambda\chi) \cos^2(\theta_i) \prod_{k=1}^{i-1} \sin^2(\theta_k), \quad (2.25)$$

$$U(\theta_i) = \prod_{k=1}^N \sin^{4-n}(\theta_k) \sum_{\{i_n\}} \sigma_{i_1, \dots, i_n}(\lambda\chi) \cos(\theta_{i_1}) \prod_{k=1}^{i_1-1} \sin(\theta_k) \dots \cos(\theta_{i_n}) \prod_{k=1}^{i_n-1} \sin(\theta_k) \quad (2.26)$$

We further assume that θ_i fields are relaxed in their stable or sufficiently long-lived (with lifetimes longer than the duration of the observable inflation) minima $\langle \theta_i \rangle = \theta_i^c$ at very early stages in the evolution of the universe. Hence, their dynamics are of no interest to us in what follows, and we can consider the following reduced form of Eq. (2.22),

$$\bar{S}_\rho = \int dx^4 \sqrt{-g} \left[\zeta(\rho) \rho^2 R - \frac{1}{2} \partial_\mu \rho \partial^\mu \rho - V(\rho) \right], \quad (2.27)$$

$$V(\rho) = \sigma(\rho) \rho^4, \quad (2.28)$$

where $\zeta \equiv \zeta(\theta_i^c)$ and $\sigma \equiv U(\theta_i^c)$. Hence, we arrive at an effective single-field model with a quartic potential and non-minimal coupling to gravity [413], but without the standard Einstein-Hilbert term. It also resembles the large field limit of the Higgs inflation model [411].

In order to reproduce the Einstein-Hilbert term in Eq. (2.27), the modulus field ρ has to develop a non-zero vacuum expectation value, $\langle \rho \rangle \equiv \rho_0$. If the vacuum configuration $\{\rho_0, \theta_i^c\}$ describes the current vacuum state of the universe, then $\rho_0 = \frac{M_p}{\sqrt{2\zeta(\rho_0)}}$ with $\zeta(\rho_0) \equiv \zeta_0 > 0$. Furthermore, the vacuum energy density, $\frac{\sigma(\rho_0)M_p^4}{4\zeta_0^2}$, in this case must be vanishingly small to satisfy the observational measurements of the dark energy density. That is, the scalar potential must be tuned so that $\sigma(\rho_0) \equiv \sigma_0 \sim 12\zeta_0^2 H_0^2 / M_p^2 \approx 0$, where H_0 is the present value of the Hubble parameter. However, inflation may end in a metastable state, which subsequently decays into the current vacuum state, alleviating this possible issue. Hence, we keep ρ_0 and σ_0 as free parameters.

The field ρ represents a flat direction of the potential in Eq. (2.28), which is constant for any value of ρ when considering the Einstein frame. Furthermore, for the special value of the coupling to gravity $\zeta_0 = -1/12$, the conformal coupling, ρ is a fictitious degree of freedom which is not manifest in the Einstein frame. In this case, the action in Eq. (2.27) in fact describes pure Einstein gravity with a cosmological constant.

2.5 Quantum Corrected Potential in Curved Spacetime

The ρ -dependence of the dimensionless couplings σ and ζ in Eq. (2.27) and Eq. (2.28) is determined by computing the quantum-corrected effective potential. At the classical level these couplings are constant and independent of ρ , and hence the action in Eq. (2.27) is classically scale invariant. In our analysis we wish to consider the 1-loop quantum corrections to the scalar potential of ρ , which are generated by the ρ^2 and ρ^4 interaction terms. In the case of these two interaction terms, the possible interactions induced via the radiative corrections will also be of the form ρ^2 and ρ^4 at 1-loop, but also with a logarithmic dependence on ρ analogous to

the Coleman Weinberg potential. These corrections can be interpreted as contributions to the running of the two dimensionless couplings σ and ζ .

Seeing as we wish to consider the inflationary setting we calculate the correction to the scalar potential in an FRW spacetime background. We use the closed form effective potential computed in Ref. [414] to obtain the 1-loop approximation of the couplings ζ and σ . The running of the couplings ζ and σ induced by the corrections is described by the β functions, β_σ and β_ζ , which encapsulate the dependence of the couplings on the energy scale. The loop corrected couplings are given by the following,

$$\zeta(\rho) = \zeta_0 + \frac{1}{2}\beta_\zeta \ln\left(\frac{\rho^2}{\rho_0^2}\right) \quad \text{and} \quad \sigma(\rho) = \sigma_0 + \frac{1}{2}\beta_\sigma \ln\left(\frac{\rho^2}{\rho_0^2}\right). \quad (2.29)$$

In our model the corresponding β functions are given by,

$$\frac{\beta_\zeta}{\zeta} = \frac{1}{4\pi^2} \frac{\sigma}{\zeta} (1 + 12\zeta) \quad \text{and} \quad \frac{\beta_\sigma}{\sigma} = \frac{9}{\pi^2} \sigma. \quad (2.30)$$

Thus, the the 1-loop approximations of the ζ and σ couplings are,

$$\zeta(\rho) = \zeta_0 + \frac{(12\zeta_0 + 1)\sigma_0}{16\pi^2} \ln\left(\frac{\rho^2}{\rho_0^2}\right), \quad (2.31)$$

$$\sigma(\rho) = \sigma_0 + \frac{9\sigma_0^2}{4\pi^2} \ln\left(\frac{\rho^2}{\rho_0^2}\right). \quad (2.32)$$

It is clear to see, that the classical scale invariance of the theory is broken by these radiative corrections, which is illustrated by the ρ dependence of the couplings found in the 1-loop corrections in Eq. (2.31) and Eq. (2.32). Note that $\sigma_0 \rightarrow 0$ is a conformal fixed-point of the theory, since the ρ dependence disappears in Eq. (2.31) and Eq. (2.32) in this limit. The conformal coupling $\zeta_0 = -1/12$ is also a fixed-point as $\zeta(\rho) = \zeta_0$. Hence, having σ small or ζ close to $-1/12$, near the respective fixed points, is natural in the technical sense. All these attractive features motivate us to consider scale invariance as an essential symmetry for natural inflation, with ρ being the inflaton field.

2.6 Observational Signatures and Model Predictions

To compute the inflationary observables we first wish to take the action in Eq. (2.27) and perform a Weyl rescaling to obtain the action in the Einstein frame. The rescaling is as follows,

$$g_{\mu\nu} \rightarrow \Omega^2 g_{\mu\nu}, \quad \Omega^2 = \frac{2\zeta\rho^2}{M_p^2}. \quad (2.33)$$

We also must make a field redefinition of the inflaton field ρ in order to obtain the canonical form of the kinetic term. The field redefinition is,

$$\rho = \rho_0 \exp\left(\frac{\sqrt{\tilde{\zeta}}}{M_p}\varphi\right), \quad (2.34)$$

where $\tilde{\zeta} = \frac{2\zeta}{1+12\zeta}$ with $\zeta > 0$ or $\zeta < -1/12$. Therefore, in the Einstein frame the action in Eq. (2.27) reads,

$$\bar{S}_\varphi = \int dx^4 \sqrt{-g} \left[\frac{M_p^2}{2} R - \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - V(\varphi) \right], \quad (2.35)$$

$$V(\rho(\varphi)) = \frac{M_p^4}{4} \frac{\sigma(\rho(\varphi))}{\zeta^2(\rho(\varphi))}. \quad (2.36)$$

In order to proceed with the actual calculations of the above observables, we substitute Eq. (2.31) and Eq. (2.32) into Eq. (2.36), and using Eq. (2.34) we can then express the effective potential in terms of the inflaton field φ in the Einstein frame. Next, let us consider the conformal limit where $\sigma_0 \rightarrow 0$ and $\zeta_0 \rightarrow -1/12$. The latter limit implies that ζ evolves slowly, $\zeta \approx \zeta_0$. If we assume further that $\sigma_0^2 \sqrt{\frac{2\zeta_0}{1+12\zeta_0}}$ approaches to some constant C , the potential in Eq. (2.36) can be well approximated by a potential which is linear in the inflaton field φ . It should be noted, a linear potential was obtained in another limit of the non-minimally coupling in Ref. [415]. Therefore, in our scenario the potential takes the following form,

$$V(\varphi) \approx \frac{162C}{\pi^2} M_p^3 \varphi. \quad (2.37)$$

This linear potential can now be used to compute the slow roll parameters and inflationary observables given in Eq. (2.8-2.13). These immediately imply that $\eta = 0$ and hence we find the relation for spectral index to be,

$$n_s \approx 1 - \frac{3}{8}r. \quad (2.38)$$

In terms of the number of observable e-folds N_\star , the model predictions are,

$$n_s - 1 \approx -0.025 \left(\frac{N_\star}{60}\right)^{-1}, \quad (2.39)$$

$$r = 0.0667 \left(\frac{N_\star}{60}\right)^{-1}. \quad (2.40)$$

If we take the observed value of the scalar perturbations - $P_s \simeq 10^{-9}$ - we require that $C \approx 5.5 \cdot 10^{-12} \left(\frac{N_\star}{60}\right)^{-3/2}$. The predictions found in Eq. (2.39) are in reasonable agreement with the

most recent analysis of the cosmological data [416], which suggest that,

$$n_s = 0.9669 \pm 0040 \quad (68\% \text{C.L.}) , \quad (2.41)$$

$$r_{0.01} < 0.0685 \quad (95\% \text{C.L.}) , \quad (2.42)$$

for the $\Lambda\text{CDM}+r$ model. As can also be seen in Figure 2.3, the predicted values fit well with observation. Further improvement of the accuracy of cosmological measurements will be critical for this scenario.




FIGURE 2.3: Constraints on Inflationary observables from the PLANCK satellite, including predictions of various inflationary potentials [365].

Note that, for large ($\xi \rightarrow \infty$) and small ($\xi \rightarrow 0$) non-minimal couplings, $n_s \gtrsim 1$, and thus the model is excluded by observation in these limiting cases.

2.7 Conclusions and Future Prospects

In this chapter, we have explored a new class of natural inflation models that exhibit hidden scale invariance, which is realised through the dilaton field. A very generic Wilsonian potential was considered, consisting of an arbitrary number of scalar fields, that was found to contain a flat direction in the classical limit, which was lifted by quantum corrections. Thus inflation can naturally, without fine-tuning, proceed as the inflaton field evolves along this direction. We find that in the conformal coupling limit, within the leading perturbative approximation, the generic model is reduced to a one-field model with a linear potential, $V(\varphi) \sim \varphi$, with the linear term being radiatively induced. Such a scenario leads to the specific predictions of the spectral index and the tensor-to-scalar ratio: $n_s - 1 \approx -0.025 \left(\frac{N_*}{60}\right)^{-1}$ and $r \approx 0.0667 \left(\frac{N_*}{60}\right)^{-1}$, respectively. These predictions are in reasonable agreement with observation, but more accurate cosmological measurements are required.

Chapter 3

An Asymmetric Universe from Inflation

It is generally considered that the generation of the observed matter-antimatter asymmetry must have occurred after the inflationary epoch, as otherwise it would have been diluted away by the rapid spacetime expansion. In order to produce a significant asymmetry during inflation, the production rate of baryonic charge must exceed its dilution rate. Despite this challenge, it has been found that inflationary dynamics may be able to support such a scenario. By utilising the observation that, if a large baryonic charge density is created due to small-scale quantum fluctuations, it will typically be stretched out over large scales due to inflation. In the last decade mechanisms have been proposed to explore this idea, but with varying success.

In this chapter, we argue that the inflationary setting can support the generation of both the matter-antimatter asymmetry and dark matter through the extension of the SM by an anomalous gauge symmetry [2, 3]. This is achieved through the addition of a general anomalous $U(1)_X$ and a dark matter fermion candidate ψ , carrying an X charge, to the SM. The associated anomaly terms source \mathcal{CP} and X charge violating processes, leading to the generation of a non-zero Chern-Simons number during inflation and subsequently a non-zero baryon number density, depending on the choice of X charge. This model is also motivated by, and able to explain, the observation that the dark matter energy density of the universe is of the same order of magnitude as that observed in the baryonic sector.

3.1 The Matter-Antimatter Asymmetry

The observed matter-antimatter asymmetry is one of the major mysteries of cosmology and particle physics. The SM predicts the existence of an asymmetry but it is many orders of magnitude smaller than that observed [214]. This asymmetry, whose energy density is believed to encompass approximately all the visible matter seen today, is quantified by the baryon asymmetry parameter,

$$\eta_B = \frac{n_B - n_{\bar{B}}}{s} \simeq \frac{n_B}{s} \simeq 8.5 \cdot 10^{-11} , \quad (3.1)$$

where n_B ($n_{\bar{B}}$) is the baryon (antibaryon) number density and s is the entropy density of the universe. The observed value of the baryon-to-entropy ratio of our Universe is determined from the observations of the temperature fluctuations in the CMB [90, 92, 193] and from BBN predictions of the light element abundances [104, 194–197].

The possibility of this asymmetry being an initial condition of the universe has been explored in the past, but has been found to be inconsistent with a successful inflationary epoch, other than in highly unnatural circumstances [417]. This is because the required initial energy density of the baryon asymmetry may dominate over the inflaton energy density, and hence inflation could not begin. This makes it very difficult to construct a model in which one can have an initial baryon number big enough to survive the dilution caused by inflation, and hence a dynamic generating mechanism during or after inflation is required. The compelling nature of this mystery has led to the proposal of many models for Baryogenesis; for some reviews see Refs. [180, 217, 224, 226, 228, 418–422]. A possible connection between the inflationary setting and the dynamical generation of the observed baryon asymmetry has been considered in the past, but due to the large dilution associated with inflation it is difficult to accumulate a large enough asymmetry. Due to this, most mechanisms of Baryogenesis are assumed to have occurred after inflation; during the reheating epoch [423] or the radiation epoch, prior to BBN. In this chapter, we will consider an unorthodox mechanism which acts during inflation, utilising an anomalous gauge symmetry extension to the SM.

In order to explain the observed baryon asymmetry of the universe, any proposed mechanism must satisfy a set of criteria known as the Sakharov conditions [205].

3.1.1 The Sakharov Conditions

The Sakharov conditions [205], formulated by A.D. Sakharov in 1967, are the requirements for successful Baryogenesis in the early universe, assuming \mathcal{CPT} is conserved. They are as follows:

- **B violation** If immediately after the Big Bang there was zero net baryon number and B is strictly conserved, then the net baryon number density of the universe would remain

zero. Furthermore, if B is strictly conserved and there was a large initial net B this would be almost completely diluted by the end of the inflationary epoch. This is in conflict with observational evidence, hence requiring the existence of B violating interactions or mechanisms [424]. Examples of possible B violating processes are perturbative proton decay [425, 426], and non-perturbative sphaleron transitions [217].

- **\mathcal{C} and \mathcal{CP} violation** The above baryon number violating processes are also required to violate the \mathcal{C} and \mathcal{CP} symmetries. Consider that there are interactions that do violate baryon number e.g. $X \rightarrow a + b$, with the antimatter equivalent $\bar{X} \rightarrow \bar{a} + \bar{b}$. If there is no \mathcal{C} and \mathcal{CP} violation then $\Gamma(X \rightarrow a + b) = \Gamma(\bar{X} \rightarrow \bar{a} + \bar{b})$, where Γ is the decay rate, and similarly for the reverse reactions. This means that the matter and antimatter decays will add and subtract baryon number at the same rate, leading to no net change. Therefore, \mathcal{C} and \mathcal{CP} violation is required for a net baryon number to be generated.
- **A period of non-equilibrium** The processes that violate B , \mathcal{C} and \mathcal{CP} must occur in a period of non-equilibrium. In thermal equilibrium, the corresponding forward and reverse reactions occur at the same rate, for both matter and antimatter processes. This means that even if \mathcal{C} and \mathcal{CP} are violated, the B violating reverse and forward reaction rates of each of the matter and antimatter processes will cancel out. The expected production of B in thermal equilibrium, at a temperature $T = \frac{1}{\beta}$, is given by,

$$\begin{aligned} \langle B \rangle_T &= \text{Tr}(e^{-\beta H} B) = \text{Tr}((\mathcal{CPT})(\mathcal{CPT})^{-1} e^{-\beta H} B) \\ &= \text{Tr}(e^{-\beta H} (\mathcal{CPT})^{-1} B (\mathcal{CPT})) \\ &= -\text{Tr}(e^{-\beta H} B) = -\langle B \rangle_T, \end{aligned} \quad (3.2)$$

where \mathcal{CPT} is the composition of the three discrete transformations \mathcal{C} , \mathcal{P} , and \mathcal{T} [217]. The \mathcal{CPT} operator commutes with H and anti-commutes with B . This implies that in thermal equilibrium the average baryon number production is zero. Hence a period of non-equilibrium is a requirement for a net baryon number to be produced. If the forward reactions were to become favoured, and the processes satisfy $\Gamma(X \rightarrow a + b) \neq \Gamma(\bar{X} \rightarrow \bar{a} + \bar{b})$, meaning \mathcal{C} and \mathcal{CP} violation, the baryon number each reaction produces will not cancel, with their reverse reactions or each other. Therefore, the inequality of the reaction rates for the forward processes will lead to an abundance of baryons over antibaryons, or vice-versa, in non-equilibrium conditions.

In fact, these criteria apply to not only the generation of a baryon asymmetry, but also any other particle asymmetry present in the universe. This is of interest with respect to dark matter, which may constitute an asymmetry produced in the dark matter sector. Such a possibility has interesting cosmological and phenomenological implications, and is an active area of research.

3.1.2 Asymmetric Dark Matter

The idea of a common origin of luminous and dark matter traces back to the 90's [199, 200, 277], but has received renewed interest in recent years; see a review in [427] and references therein. The major motivation to this hypothesis comes from the observation that the present-day mass density of dark matter is of the same order of magnitude as the density of visible matter [92],

$$\rho_{\text{DM}} \simeq 5.5\rho_B . \quad (3.3)$$

The similarity in these observed densities is perhaps an indication of a strong connection between the physics and cosmological evolution of visible and dark matter. Hence, within this picture an asymmetry of similar size may be expected to be generated among dark matter particles and antiparticles. In the model considered in this chapter, visible and dark matter are connected by a common anomalous gauged $U(1)_X$, which we introduce in addition to the gauge group of the SM. It is proposed that this anomalous gauge symmetry is responsible for the observed particle asymmetries, with the ratio of dark to luminous mass density related to how the X charge asymmetry is distributed between these two sectors.

3.2 Topological Vacuum States and the Chern-Simons Number

A concept that will be important in theogenesis scenario, we consider in this chapter, is the Chern-Simons (CS) number. The CS number is an integer related to the winding number, which quantifies the topological non-triviality of a given vacuum state. The topological nature of the vacuum state arises due to non-trivial boundary conditions in the associated gauge theory. Full derivatives normally integrate to zero in the action integral because of the assumed boundary conditions. Although, due to the internal structure of the gauge theories involved this may not always be the case. These gauge field configurations can lead to non-trivial topology at the boundaries which are dependent on the choice of gauge, analogous to the boundary conditions in a kink solution. The vacua are each denoted with winding numbers that define certain homotopy classes, or gauge classes. It is not possible to smoothly deform an element of one homotopy class into an element of another class due to topological obstruction [53, 54, 428].

These topologically distinct vacua are degenerate, and are separated by potential barriers, analogous to a set of degenerate potential wells. Each are denoted by a CS number, which is proportional to the winding number, as depicted in Figure 3.1. As these vacuum states are related to the boundary conditions of the full derivative terms, they do not contribute to perturbation theory, but they can provide interesting non-perturbative effects through transitions between them, which we shall discuss below.

The CS number of a given gauge field configuration, in terms of the associated gauge field A_μ^a , is defined as [429],

$$n_{CS} := n_g \frac{g^2}{32\pi^2} \int d^3x \epsilon^{ijk} \text{Tr}(A_i \partial_j A_k + \frac{2ig}{3} A_i A_j A_k), \quad (3.4)$$

where $A_\mu = A_\mu^a T^a$, with T^a the generators of the applicable gauge group, and g is the corresponding coupling constant.

Such topologically non-equivalent vacua are of particular relevance to the $SU(2)$ gauge group in the SM, where at infinity there is a set of mappings to the number of windings around the S^3 symmetry of the internal $SU(2)$ space. The number of windings is related to the CS number. In the SM, the $SU(2)$ theory has anomalies associated with the B and L global symmetries. This topological structure is related to these anomalies, and as we shall see, transitions between the vacua violate the associated global charges.




FIGURE 3.1: A sphaleron transition between degenerate vacua with different CS numbers.

3.2.1 Instanton and Sphaleron Transitions

An instanton is a semi-classical solution of the gauge field equations of motion in Euclidean space, that describes a topologically non-trivial vacuum gauge field configuration [430–437]. The instanton transitions are non-perturbative processes, meaning that they cannot be described in the Lagrangian by a renormalisable operator, and as such the interaction cannot be represented by a Feynman diagram. However, an effective operator, or vertex, can be constructed. In the SM, instanton solutions are present for the weak and strong gauge groups ($SU(2)$ and $SU(3)$), and have anomalous global currents through which such transitions can be mediated.

In EWB scenarios, the source of B violation comes from so called sphaleron transitions. These are vacuum to vacuum transitions induced by thermal excitations that result in a change in the CS number, which is depicted in Figure 3.1. The $SU(2)$ gauge field vacuum configurations lead

to the manifestation of these topological vacua, and as such the transitions between vacuum states are mediated via the B and L global anomalous currents. Each transition event results in $\Delta B = \Delta L = n_g$, meaning that $B - L$ is conserved; n_g is the number of generations. An example of such a process is given in Figure 3.2.

As described above, instanton quantum tunnelling processes or thermal excitations can change the topological number of a vacuum state. During such a transition, the fermion fields must also evolve as the gauge field configuration changes between topologically distinct vacuum states. This leads to the generation of particles depending on the associated anomalous symmetries.




FIGURE 3.2: An example of a sphaleron transition, three antileptons are converted into nine quarks, conserving $B - L$.

Seeing as sphalerons correspond to thermal excitations over the potential barriers, rather than quantum tunnelling, they are highly suppressed at low temperatures, and are only important at temperatures above the characteristic energy of the potential barrier. In the case of the electroweak vacuum, the sphaleron processes become important at temperatures above the EWPT ($T \sim 100$ GeV). The thermal rate of sphaleron processes in the broken phase, below the temperature of the EWPT, is [81],

$$\Gamma_{sph}(T) = \mu \left(\frac{M_W}{\alpha_W T} \right)^3 M_W^4 e^{-\frac{E_{sph}(T)}{T}}, \quad (3.5)$$

where M_W is the mass of the W boson, T is the temperature, μ is a constant, $E_{sph}(T)$ is the sphaleron energy, and $\alpha_W = \frac{g_2^2}{4\pi} \simeq \frac{1}{29}$ with g_2 being the gauge coupling of $SU(2)$.

This indicates why these baryon and lepton number violating processes are not observed today. An estimate of the rate of sphaleron transitions per unit volume, in which κ is a dimensionless constant, is given by,

$$\Gamma_{sph}(T) \simeq \kappa \alpha_W^4 T^4. \quad (3.6)$$

This rate is strongly dependent on the temperature of the state, where when the temperatures are between the EWPT ($T \sim 100$ GeV) and 10^{12} GeV the sphalerons are in thermal equilibrium [438, 439]. Therefore, these processes are of interest when investigating early universe particle dynamics, due to the high temperatures that may be present.

Although not integral to our scenario, electroweak sphaleron transitions need to be taken into account. They can act in thermal equilibrium after reheating, leading to B and L redistribution. Therefore, depending on the reheating temperature, the $B - L$ number contained within the X charge asymmetry generated by our model will be redistributed between the SM fermionic degrees of freedom by equilibrium sphaleron transitions during the radiation era. The quantity of the initial $B - L$ number density distributed into baryon number is given by the following [440],

$$n_B = \frac{28}{79} n_{B-L} , \quad (3.7)$$

where this relation is applicable when the temperature of the primordial plasma is above $T \sim 100$ GeV.

The time variation of the vacuum state in an inflationary setting can lead to the accumulation of CS number in the presence of the X charge anomalies. In the scenario we propose in this work, the accumulated topological charge in the vacuum will thus correspond to a net X charge.

3.3 A Model of Inflationary Cogenesis

In what follows, we consider an application of these ideas in an inflationary setting to construct a new mechanism for generating both luminous and dark matter during cosmic inflation. It has been suggested that inflation may play an even more prominent role in cosmology than solving the mysteries discussed in Chapter 1, but by also generating the observed matter-antimatter asymmetry in the universe [2, 250–252, 441, 442]. Namely, in [2] we argued that a successful Baryogenesis scenario can be realised during inflation within models containing anomalous gauge symmetries [73]. One of the first uses of anomalies for the generation of the baryon asymmetry was first utilised in a model of inflationary leptogenesis [251], in which a lepton asymmetry is produced during inflation due to the gravitational birefringence induced through the gravitational lepton number anomaly coupled to an introduced pseudoscalar field.

In the model we consider in this chapter, ordinary and dark matter both carry charges associated with an anomalous $U(1)_X$ group. Anomaly terms in the model Lagrangian source \mathcal{CP} and $U(1)_X$ charge violating processes during inflation, producing corresponding non-zero CS numbers which are subsequently reprocessed into baryon and dark matter densities. Other recent works have considered relating the generation of the luminous matter-antimatter asymmetry with an asymmetry in the dark sector within a gauged $U(1)_X$ extension of the SM [443–449].

In the early universe, when the expansion rate is faster than processes with fermion chirality flip, the gauged anomalies may effectively appear within the SM [450]. Indeed, it has been argued that anomalous production of the right-handed electron number is possible through the hypercharge anomaly in the SM [451]. An inflationary version of the above scenario is discussed in [252]. The anomalous gauge theory we consider can also be viewed as an effective low-energy theory, which admits a fundamental completion free of gauge anomalies. The obvious candidates for such an anomalous gauge theory are gauged baryon and lepton numbers, or any linear combination thereof.

As with any \mathcal{CPT} invariant model attempting to explain the dynamical generation of a charge asymmetry, it must satisfy the Sakharov's conditions as discussed above, and ours does so as follows. The anomalies present upon introduction of the $U(1)_X$ gauge boson provide our X charge violation. On top of this, the gauge invariance of the $U(1)_X$ requires a pseudoscalar field, that describes the longitudinal polarization of the X boson, to couple to these anomalies. In the cosmological setting these interactions spontaneously violate \mathcal{CP} invariance, and the inflationary epoch provides the push out-of-equilibrium that is required for the accumulation of the asymmetry. This will be discussed in more detail below. Note that this mechanism differs from the one presented in [452], in which a non-zero CS number in the hypercharge field is generated during inflation. In their case, the conversion of this to baryon number happens at the electroweak scale through altering the right-handed electron chemical potential, while also requiring a strongly first order electroweak phase transition which is not achieved within the SM.

3.3.1 Models with an Anomalous $U(1)_X$

In this general mechanism, we consider an extension of the SM that is based on the $SU(3) \times SU(2) \times U(1)_Y \times U(1)_X$ gauge group and contains an additional fermion(s) that shall act as a dark matter candidate. The introduction of a scalar to play the role of the inflaton is also required, but the detailed dynamics of the inflationary epoch is not important for our analysis. The new $U(1)_X$ gauge symmetry is assumed to be anomalous, and hence the corresponding gauge boson will be necessarily massive with gauge invariance realised non-linearly. The longitudinal degree of freedom of this $U(1)_X$ gauge field is then described by a scalar field $\theta(x)$, which allows anomaly cancellation through the introduction of appropriate counter-terms [73], through the Green-Schwarz mechanism as discussed in Chapter 1. This theory can be viewed as a low energy limit of an anomaly-free theory, either within ordinary QFT or string theory. In the presence

of a cubic anomaly $U(1)_X^3$, the additional Lagrangian terms important to our analysis are,

$$\begin{aligned} \frac{1}{\sqrt{-g}}\mathcal{L}_X = & -\frac{1}{4}g^{\mu\alpha}g^{\nu\beta}X_{\mu\nu}X_{\alpha\beta} + \frac{1}{2}f_X^2g^{\mu\nu}(g_XX_\mu - \partial_\mu\theta)(g_XX_\nu - \partial_\nu\theta) \\ & - \mathcal{A}_1\frac{g_X^2}{16\pi^2}\theta(x)X_{\mu\nu}\tilde{X}^{\mu\nu}, \end{aligned} \quad (3.8)$$

where $X_{\mu\nu}$ denotes the field strength of the $U(1)_X$ gauge boson with corresponding coupling constant $g_X = m_X/f_X$, f_X is a parameter that defines the mass of the $U(1)_X$ boson (m_X), and $\tilde{X}^{\mu\nu} = \frac{1}{2\sqrt{-g}}\epsilon^{\mu\nu\rho\sigma}X_{\rho\sigma}$ is the dual field strength, in which $\epsilon^{\mu\nu\rho\sigma}$ is the Levi-Civita tensor. We have omitted fermion interactions and the charged current j_X terms. The final term in Eq. (3.8) is responsible for maintaining gauge invariance of the full quantum theory description under $U(1)_X$ transformations, as discussed above. In the proceeding analysis any associated gravitational anomaly is ignored as its contribution is considered to be negligible with respect to the other anomalous contributions.

We shall consider two example applications of this general model forogenesis. Namely, we consider the extensions involving an anomalous gauged $B-L$ and B [75–77, 453–460]. The examples we discuss shall also contain a fermionic field(s) ψ , which carries a chiral charge under the anomalous gauge symmetry and is sterile under the SM gauge symmetry. The charges of each of the fermions under these additional gauge symmetries are given in Table 3.1. The mass m_ψ is an extra parameter which can be directly introduced within the non-linear realisation of the anomalous gauge symmetry. Typically, such a mass is also generated radiatively within the low-energy effective theory, reflecting a more conventional mechanism for mass generation within an ultraviolet anomaly-free completion.

The stability of the dark matter candidate ψ is ensured as the Lagrangian in Eq. (3.8) has no interaction vertices which allow violation of the global ψ number locally. Violation of X charge, and hence also the dark X charge, only occurs in this model through the non-perturbative generation of CS number in an expanding spacetime. This production is only of significance during inflation, as contributions during subsequent matter and radiation dominated epochs are negligible due to minimal \mathcal{CP} violation and push out-of-equilibrium. This means that after the inflationary epoch ends, the dark matter density to entropy density ratio is fixed, assuming no significant additional sources of entropy.

Case 1: $U(1)_{B-L}$ and a sterile fermion ψ

In the SM, an additional $U(1)_{B-L}$ gauge symmetry is anomalous unless three right handed neutrinos are introduced. The associated anomalies are trace and cubic: $\mathcal{A}_0(U(1)_{B-L}) = -3$ and $\mathcal{A}_1(U(1)_{B-L}^3) = -3$. We introduce N_ψ new right-handed (for definiteness) Weyl fermions ψ , some of which act as dark matter candidates in our model. For simplicity we assume that they

Fermion Field	$SU(3)$	$SU(2)$	$U(1)_Y$	Case 1: $U(1)_{B-L}$	Case 2: $U(1)_B$
$Q_L^i = \begin{pmatrix} u \\ d \end{pmatrix}_L^i$	3	2	1/6	1/3	1/3
u_R^i	3	1	2/3	1/3	1/3
d_R^i	3	1	-1/3	1/3	1/3
$L^i = \begin{pmatrix} \nu \\ e \end{pmatrix}_L^i$	1	2	-1/2	-1	0
e_R^i	1	1	-1	-1	0
ψ	1	1	0	q_ψ	q_ψ

TABLE 3.1: The representations of the SM fermions and dark fermion ψ in reference to the gauge symmetries of the theory.

carry the same $B - L$ charge q_ψ and interact only via exchange of the $B - L$ gauge boson. The addition of these states alters the $B - L$ anomalies as follows: $\mathcal{A}_0 := \mathcal{A}_0(U(1)_{B-L}) = -N_\psi q_\psi - 3$ and $\mathcal{A}_1 := \mathcal{A}_1(U(1)_{B-L}^3) = -N_\psi q_\psi^3 - 3$. In this case, the dark matter fermion does not introduce any new anomalies. We will ignore the gravitational anomaly \mathcal{A}_0 in our analysis, but it should be noted that taking $q_\psi = -3$ or -1 and $N_\psi = 1$ or 3 , respectively, eliminates \mathcal{A}_0 . Obviously, $N_\psi = 3$, $q_\psi = -1$ removes all anomalies, so we are not interested in such a charge assignment in this paper.

The addition of the $B - L$ gauge symmetry and dark matter candidate to the SM leads to a Lagrangian density of the same form given for the general case presented in Eq. (3.8).

Case 2: $U(1)_B$ and a sterile baryon ψ

Gauging the baryon number symmetry of the SM results in the inclusion of two mixed anomalies involving the weak and hypercharge gauge groups: $\mathcal{A}_2(SU(2)^2 \times U(1)_B) = 3/2$ and $\mathcal{A}_3(U(1)_Y^2 \times U(1)_B) = -3/2$. The addition of a new sterile state ψ leaves these mixed anomalies unchanged, but introduces two new unmixed anomalies: $\mathcal{A}_0 := \mathcal{A}_0(U(1)_B) = -N_\psi q_\psi$ and $\mathcal{A}_1 := \mathcal{A}_1(U(1)_B^3) = -N_\psi q_\psi^3$. Hence, there are four anomalies, each of which will contribute to baryonic charge generation during the inflationary epoch, but only two of which will include generation of fermions in the dark matter sector, namely, \mathcal{A}_0 and \mathcal{A}_1 .

The presence of additional mixed anomalies means that extra anomaly cancelling terms are required with respect to the gauged $B - L$ case considered above, that is,

$$\begin{aligned} \frac{1}{\sqrt{-g}} \mathcal{L}_X = & -\frac{1}{4} g^{\mu\alpha} g^{\nu\beta} X_{\mu\nu} X_{\alpha\beta} + \frac{1}{2} f_X^2 g^{\mu\nu} (g_X X_\mu - \partial_\mu \theta) (g_X X_\nu - \partial_\nu \theta) \\ & - \mathcal{A}_1 \frac{g_X^2 \theta(x)}{16\pi^2} X_{\mu\nu} \tilde{X}^{\mu\nu} - \mathcal{A}_2 \frac{g_1^2 \theta(x)}{16\pi^2} B_{\mu\nu} \tilde{B}^{\mu\nu} - \mathcal{A}_3 \frac{g_2^2 \theta(x)}{16\pi^2} W_{\mu\nu}^a \tilde{W}^{a\mu\nu}, \end{aligned} \quad (3.9)$$

where $B_{\mu\nu}$ and $W_{\mu\nu}$ denote the hypercharge and weak field strengths respectively, with corresponding coupling constants g_1 and g_2 .

3.4 Dynamics of an Anomalous Gauge Field During Inflation

For a model to successfully produce a charge asymmetry in the early universe it must satisfy the well-known Sakharov conditions [205]. We will now discuss the framework of our new mechanism for cogenesis and how it satisfies these criteria in more detail.

Firstly, we wish to describe the universe using the Friedmann-Robertson-Walker metric tensor, which represents a homogeneous, isotropic and spatially flat cosmological spacetime. In conformal coordinates the metric can be expressed as: $g_{\mu\nu} = a^2(\tau)\eta_{\mu\nu}$. During inflation the scale factor $a(\tau)$ is given by the following,

$$a(\tau) = -1/H_{\text{inf}}\tau, \quad (3.10)$$

where H_{inf} is the expansion rate during inflation ($H_{\text{inf}} \cong \text{constant}$). The conformal time is given in the range $\tau \in [-\frac{1}{H}, -\frac{1}{H}e^{-N_{\text{inf}}}]$ during inflation, where N_{inf} is the number of e-folds during the inflationary epoch, such that the scale factor is $a(\tau_0) = 1$ at the beginning of inflation.

To allow analytical treatment, the analysis that follows requires certain simplifying assumptions. For the θ field we only consider a classical homogeneous background configuration, $\theta(\tau, \vec{x}) = \theta(\tau)$, and ignore quantum fluctuations over it. We take $g_X \ll 1$ such that the $\theta(x)$ and $X_\mu(x)$ fields essentially decouple from each other. This also implies that the $U(1)_X$ boson is light relative to the scale f_X , $m_X/f_X \ll 1$, and hence we will not be interested in its dynamics during inflation. With these assumptions the Lagrangian in Eq. (3.8) becomes,

$$\begin{aligned} \mathcal{L}_X = & -\frac{1}{4}\eta^{\mu\alpha}\eta^{\nu\beta}X_{\mu\nu}X_{\alpha\beta} + \frac{1}{2}a(\tau)^2\eta^{\mu\nu}(m_X X_\mu - \partial_\mu\phi(\tau))(m_X X_\nu - \partial_\nu\phi(\tau)) \\ & - \mathcal{A}_1 \frac{g_X^2\phi(\tau)}{32\pi^2 f_X} \epsilon^{\mu\nu\alpha\beta} X_{\mu\nu} X_{\alpha\beta}, \end{aligned} \quad (3.11)$$

where $\phi(\tau) \equiv f_X\theta(\tau)$. From this Lagrangian follows the equation of motion for $\phi(\tau)$,

$$(a^2\phi')' = 0, \quad (3.12)$$

where $\phi' \equiv d\phi/d\tau$ and we have ignored any terms quadratic in X_μ . Solving for $\phi'(\tau)$ we obtain,

$$\phi'(\tau) = \frac{a^2(\tau_0)\phi'_0}{a^2(\tau)}, \quad (3.13)$$

where ϕ'_0 is an integration constant associated with the ‘field velocity’ at the start of inflation, which is defined at $\tau = \tau_0$, where $a(\tau_0) = 1$. An upper limit on the value of ϕ'_0 is provided by the requirement that the initial energy density of the ϕ field be less than that of the inflaton field. This upper limit is thus $\phi'_0 \lesssim H_{\text{inf}}M_p$, where $M_p = 1/\sqrt{G}$ is the Planck mass. Substituting Eq.

(3.13) into the linearised equation of motion for the X_μ gauge field gives,

$$\left(\partial_\tau^2 - \vec{\nabla}^2 + \left(\frac{a(\tau_0)m_X}{H_{\text{inf}}\tau} \right)^2 \right) X^i + \kappa_X \tau^2 \epsilon^{ijk} \partial_j X_k = 0, \quad (3.14)$$

where

$$\kappa_X = |\mathcal{A}_1| \frac{a^2(\tau_0)g_X^2\phi'_0 H_{\text{inf}}^2}{4\pi^2 f_X}, \quad (3.15)$$

and the gauge $X_0 = \partial_i X_i = 0$ has been chosen. The source of \mathcal{CP} violation in our model is apparent in Eq. (3.14) where the two terms have opposite \mathcal{P} , and hence, \mathcal{CP} transformations.

In the discussion that follows we treat the $U(1)_X$ gauge boson as a massless particle, as we have assumed $m_X \ll H_{\text{inf}}$. To then quantize this model we promote the X gauge boson fields to operators and assume that the boson has two possible circular polarisation states,

$$X_i = \int \frac{d^3\vec{k}}{(2\pi)^{3/2}} \sum_\alpha \left[G_\alpha(\tau, k) \epsilon_{i\alpha} \hat{a}_\alpha e^{i\vec{k}\cdot\vec{x}} + G_\alpha^*(\tau, k) \epsilon_{i\alpha}^* \hat{a}_\alpha^\dagger e^{-i\vec{k}\cdot\vec{x}} \right], \quad (3.16)$$

where $\vec{\epsilon}_\pm$ denotes the two possible helicity states of the $U(1)_X$ gauge boson ($\vec{\epsilon}_+^* = \vec{\epsilon}_-$) and the creation, $\hat{a}_\alpha^\dagger(\vec{k})$, and annihilation, $\hat{a}_\alpha(\vec{k})$, operators satisfy the canonical commutation relations,

$$\left[\hat{a}_\alpha(\vec{k}), \hat{a}_\beta^\dagger(\vec{k}') \right] = \delta_{\alpha\beta} \delta^3(\vec{k} - \vec{k}'), \quad (3.17)$$

and

$$\hat{a}_\alpha^a(\vec{k})|0\rangle_\tau = 0, \quad (3.18)$$

where $|0\rangle_\tau$ is an instantaneous vacuum state at time τ .

The mode functions in Eq. (3.16) are described by the following equations, from Eq. (3.14),

$$G_\pm'' + \left(k^2 + \frac{\lambda^2}{\tau^2} \mp \kappa_X \tau^2 k \right) G_\pm = 0, \quad (3.19)$$

where $\lambda = \frac{m_X}{H_{\text{inf}}}$, which is assumed to be small as stated above.

Solving for the mode functions G_\pm in Eq. (3.19) gives,

$$G_+(\tau, k) = 2^{\frac{1+\nu}{2}} e^{-z} 2\tau^{\frac{1}{2}+\nu} \left[C_1 U \left(\frac{1+\nu}{2} - \frac{\Omega_k}{4}, 1+\nu, z \right) + C_2 M \left(\frac{1+\nu}{2} - \frac{\Omega_k}{4}, 1+\nu, z \right) \right] \quad (3.20)$$

and

$$G_-(\tau, k) = 2^{\frac{1+\nu}{2}} e^{z} \tau^{\frac{1}{2}+\nu} \left[C_3 U \left(\frac{1+\nu}{2} - \frac{i\Omega_k}{4}, 1+\nu, \frac{z}{i} \right) + C_4 M \left(\frac{1+\nu}{2} - \frac{i\Omega_k}{4}, 1+\nu, \frac{z}{i} \right) \right] \quad (3.21)$$

where $z = \frac{k^2 \tau^2}{\Omega_k}$, $\Omega_k = \sqrt{\frac{k^3}{\kappa_X}}$, $\nu = \frac{1}{2} \sqrt{1 - 4\lambda^2} \sim \frac{1}{2} - \lambda^2$, $U(a, b, z)$ is a confluent hypergeometric function of the second kind, and $M(a, b, z)$ is a confluent hypergeometric function of the first kind (Kummer Function).

In the limit $|\tau| \rightarrow 0$ (or $k^2 + \frac{\lambda^2}{\tau^2} \gg \kappa_X \tau^2 k$), \mathcal{CP} -invariant wave modes are obtained. These are described by,

$$X_i = \int \frac{d^3 \vec{k}}{(2\pi)^{3/2}} \sum_{\alpha} \left[F_{\alpha}(\tau, k) \epsilon_{i\alpha} \hat{b}_{\alpha} e^{i\vec{k} \cdot \vec{x}} + F_{\alpha}^*(\tau, k) \epsilon_{i\alpha}^* \hat{b}_{\alpha}^{\dagger} e^{-i\vec{k} \cdot \vec{x}} \right], \quad (3.22)$$

where the wave mode functions F_{\pm} are found to be,

$$F_{+}(\tau, k) = \frac{\sqrt{\pi\tau}}{2} H_{\nu}^{(2)}(k\tau) e^{-i\frac{\pi}{2}(\frac{1}{2} + \nu)} \quad \text{and} \quad F_{-}(\tau, k) = \frac{\sqrt{\pi\tau}}{2} H_{\nu}^{(1)}(k\tau) e^{i\frac{\pi}{2}(\frac{1}{2} + \nu)}. \quad (3.23)$$

By matching the modes in Eq. (3.20) and (3.21) to those in Eq. (3.23) and using the known Wronskian normalisation we can determine the coefficients C_{1-4} . For more details on this calculation and the form of the coefficients see Appendix B.

Particle Creation during Inflation and Bogolyubov Transformations

In an expanding spacetime it is difficult to define a time independent vacuum state because the Hamiltonian becomes time dependent. In a flat spacetime, the vacuum is defined with reference to plane wave solutions, such that excitations from the vacuum state correspond to plane waves. In an expanding spacetime background the form of the annihilation and creation operators will change with time, and thus the canonical vacuum state can only be defined at any given instant in time. This means that the Hamiltonian's instantaneous energy eigenstates will not be the same at all times, and hence the vacuum state will not be as well. That is, the vacuum state at one time will correspond to an excited state of the vacuum at another time, which means that on comparing the vacua, a relative particle number with reference to one another may be found. This is how the expansion of the universe during inflation can lead to particle production [461–464]. It is possible to relate the particle number of two vacuum states by a so called Bogolyubov transformation. This type of transformation allows the determination of the particle content of an evolving vacuum state with respect to another vacuum state.

If we define two different annihilation operators \hat{a} and \hat{b} we in turn define the corresponding a and b vacua; $|0_a\rangle$ and $|0_b\rangle$, respectively. It is possible to express the b -vacuum as a superposition of a particle states and vice versa. The a -vacuum will contain no a particles, but may have a b -particle density. A Bogolyubov transformation performed between the a -vacuum and b -vacuum

can allow the calculation of this relative particle density, and is defined as [373],

$$\hat{b}_\gamma = \alpha_\gamma \hat{a}_\gamma + \beta_\gamma^* \hat{a}_\gamma^\dagger \quad \text{and} \quad \hat{b}_\gamma^\dagger = \alpha_\gamma^* \hat{a}_\gamma^\dagger + \beta_\gamma \hat{a}_\gamma, \quad (3.24)$$

where α and β are k dependent complex numbers. In order to determine α and β the modes corresponding to each vacuum state must be matched at some point in their evolution. This is done as follows,

$$v^*(t_0) = \alpha_\gamma u_\gamma^*(t_0) + \beta_\gamma u_\gamma(t_0), \quad (3.25)$$

$$v^{*'}(t_0) = \alpha_\gamma u_\gamma^{*'}(t_0) + \beta_\gamma u_\gamma'(t_0), \quad (3.26)$$

where v and u are the modes defined for the a and b vacuum states, respectively.

In our model, a Bogolyubov transformation will be utilised to match the evolving inflationary vacuum and the vacuum state for super horizon modes. This is because, at scales smaller than the Hubble rate the effects of spacetime curvature become negligible, and solutions must converge to plane waves. Rather than matching the vacua to calculate the net particle density accumulated, we shall determine the accumulated CS number density that is induced by the X charge violating anomalous interactions during the inflationary epoch.

In our scenario, we compare the birefringent and \mathcal{CP} -invariant modes to derive the Bogolyubov coefficients relating the two sets of creation and annihilation operators, $\{\hat{a}_\alpha^a, \hat{a}_\alpha^{a\dagger}\}$ and $\{\hat{b}_\alpha^a, \hat{b}_\alpha^{a\dagger}\}$, in Eq. (3.16) and Eq. (3.22). The Bogolyubov transformations in this case are defined by,

$$\hat{b}_\alpha^a(\vec{k}) = \alpha_\alpha a_\alpha^{a\dagger}(\vec{k}) + \beta_\alpha^* \hat{a}_\alpha^a(\vec{k}), \quad (3.27)$$

$$\hat{b}_\alpha^{a\dagger}(\vec{k}) = \alpha_\alpha^* a_\alpha^a(\vec{k}) + \beta_\alpha \hat{a}_\alpha^{a\dagger}(\vec{k}). \quad (3.28)$$

The relevant Bogolyubov coefficients are found to be,

$$\alpha_\pm = 1 - \frac{1}{2^{1-\nu}} \left(1 \pm \frac{i\lambda^2}{(k\tau)^{1-2\lambda^2}} \left(1 - \frac{\pi(k\tau)^{1-2\lambda^2}}{2^\nu} \right) \mp \frac{i2^{1-\nu}(k\tau)^{\lambda^2}}{\sqrt{k}} e^{\mp i\pi\lambda^2/2} G_{\pm}^{\prime*} \Big|_{\frac{k\tau^2}{k}, k|\tau| \rightarrow 0} \right), \quad (3.29)$$

and,

$$\beta_\pm = \frac{e^{\mp i\pi\lambda^2}}{2^{1-\nu}} \left(1 \mp \frac{i\lambda^2}{(k\tau)^{1-2\lambda^2}} \left(1 - \frac{\pi(k\tau)^{1-2\lambda^2}}{2^\nu} \right) \pm \frac{i2^{1-\nu}(k\tau)^{\lambda^2}}{\sqrt{k}} e^{\pm i\pi\lambda^2/2} G_{\mp}^{\prime*} \Big|_{\frac{k\tau^2}{k}, k|\tau| \rightarrow 0} \right), \quad (3.30)$$

where we have considered the superhorizon modes ($k|\tau| \approx 0$).

3.5 Simultaneous Generation of Luminous and Dark Matter During Inflation

Now that we have determined the dynamics of the X_μ gauge field we can calculate the general X charge density generated during inflation. It is known that the anomalous non-conservation of the X charge current is given by,

$$\partial_\mu (\sqrt{-g} j_X^\mu) = \mathcal{A}_1 \frac{g_X^2}{32\pi^2} \epsilon^{\mu\nu\rho\sigma} X_{\mu\nu} X_{\rho\sigma} \equiv \mathcal{A}_1 \frac{g_X^2}{8\pi^2} \partial_\mu (\sqrt{-g} K^\mu) , \quad (3.31)$$

where $K^\mu = \frac{1}{2\sqrt{-g}} \epsilon^{\mu\nu\rho\sigma} X_{\nu\rho} X_\sigma$ is a topological current. This implies that the net X charge density $n_X = n_x - n_{\bar{x}} \equiv a^{-1}(\tau) \langle 0 | j_X^0 | 0 \rangle$ is related to the CS number density of the $U(1)_X$ gauge boson by the following equation,

$$n_X = |\mathcal{A}_1| \frac{g_X^2}{8\pi^2} a(\tau_{\text{end}}) n_{CS} , \quad (3.32)$$

where $\tau = \tau_{\text{end}}$ is the conformal time at the end of inflation, and $n_X(\tau_0) = n_{CS}(\tau_0) = 0$ at the start of inflation. The form of the CS number is given below, in which we wish to consider only large scale superhorizon modes ($k|\tau| \simeq 0$),

$$\begin{aligned} n_{CS} &= \frac{1}{a^4(\tau_{\text{end}})} \epsilon^{ijk} \lim_{k|\tau| \rightarrow 0} \langle 0 | X_i \partial_j X_k | 0 \rangle \\ &\simeq \frac{1}{4\pi^2 a^4(\tau_{\text{end}})} \int_\mu^\Lambda k dk \left[|G'_+|^2_{\frac{\kappa\tau^2}{k}, k|\tau| \rightarrow 0} - |G'_-|^2_{\frac{\kappa\tau^2}{k}, k|\tau| \rightarrow 0} \right] - \mathcal{O}(\lambda^2) , \end{aligned} \quad (3.33)$$

where we ignore small terms with quadratic or higher orders of λ . Note that the upper limit in the integral in Eq. (3.33) simply cuts out sub-horizon modes for which our approximate calculations are not applicable. The ultraviolet modes do not give a significant contribution anyway, since they act as \mathcal{CP} -invariant planewaves, which expectantly lead to a cancellation between the positive and negative frequency modes. The dominant contribution to n_{CS} is given by infrared modes, and in fact the integral is divergent. This divergence is reminiscent of the well-known infrared divergence of de Sitter-invariant two-point functions, which possibly signals that the pure de Sitter approximation of the inflationary phase becomes inadequate in our case. There is no commonly accepted prescription for regularization of these types of divergences in the literature and we simply introduce an infrared cut-off μ .

We assume that the only non-negligible source of entropy density is the process of reheating after inflation, for which the entropy density produced is $s \simeq \frac{2\pi^2}{45} g_* T_{\text{rh}}^3$, where T_{rh} is the associated reheating temperature and $g_*(T_{\text{rh}}) \simeq 106.75$. Upon taking a first order expansion around $\Omega_k = 0$ in Eq. (3.33), we obtain the following expression for the X charge asymmetry parameter

generated by the unmixed anomaly,

$$\begin{aligned} \eta_X = \frac{n_X}{s} &\approx |\mathcal{A}_1| \frac{30g_X^2}{\pi^{10}g_*} \Gamma\left(\frac{3}{4}\right)^4 e^{-3N_e} \left(\frac{\kappa_X}{\mu T_{\text{rh}}^2}\right)^{\frac{3}{2}} \\ &\approx 8.5 \cdot 10^{-11} |\mathcal{A}_1|^{5/2} \left(\frac{m_X}{10^{12} \text{ GeV}}\right)^5 \left(\frac{\phi'_0}{10^{32} \text{ GeV}^2}\right)^{\frac{3}{2}} \left(\frac{H}{10^{14} \text{ GeV}}\right) \\ &\times \left(\frac{T_{\text{rh}}}{2 \cdot 10^{11} \text{ GeV}}\right)^{-2} \left(\frac{f_X}{10^{14} \text{ GeV}}\right)^{-\frac{13}{2}} \left(\frac{\mu}{10^{-42} \text{ GeV}}\right)^{-\frac{3}{2}}, \end{aligned} \quad (3.34)$$

where N_e denotes the minimum number of e-folds required to solve the horizon and flatness problems, and includes the additional dilution that occurs if the reheating period is not instantaneous. The number of e-folds that contribute to the dilution of n_X is,

$$N_e = N_{\text{inf}} + N_{\text{rh}} \simeq 27.5 + \frac{2}{3} \ln\left(\frac{H_{\text{inf}} M_p}{(1 \text{ GeV})^2}\right) - \frac{1}{3} \ln\left(\frac{T_{\text{rh}}}{1 \text{ GeV}}\right), \quad (3.35)$$

where

$$N_{\text{inf}} \simeq 34 + \ln\left(\frac{T_{\text{rh}}}{100 \text{ GeV}}\right) \quad \text{and} \quad N_{\text{rh}} \simeq \frac{2}{3} \ln\left(\frac{H_{\text{inf}} M_p}{T_{\text{rh}}^2}\right) - 1.89. \quad (3.36)$$

In Eq. (3.34), it can be seen that taking the parameter κ_X to be large increases the asymmetry. Also, the infrared cut-off μ must be sufficiently small. In what follows we consider two possible cut-offs - the minimal box cut-off [465], $\mu = H_0 \approx 10^{-42} \text{ GeV}$, which accounts for all the modes that are within the present Hubble horizon, and $\mu = H_{\text{BBN}} \approx 10^{-25} \text{ GeV}$ which includes all of the modes within the Hubble horizon at the beginning of the period of BBN; assuming $T_{\text{BBN}} \sim 1 \text{ MeV}$.

A similar relation to Eq. (3.34) can be derived for the mixed anomalies. In particular, these can be present in the case of gauged baryon number $U(1)_B$, as considered in [2]. In the case of the electroweak and hypercharge mixed anomalies the extra contribution to the total X charge asymmetry is,

$$\eta_X^{\text{mixed}} = \frac{n_X^{\text{mixed}}}{s} \approx (|\mathcal{A}_2|^{5/2} g_1^5 + 3|\mathcal{A}_3|^{5/2} g_2^5) \frac{15}{4\pi^{13}g_*} \Gamma\left(\frac{3}{4}\right)^4 e^{-3N_e} \left(\frac{\kappa}{\mu T_{\text{rh}}^2}\right)^{\frac{3}{2}}, \quad (3.37)$$

where $\kappa = \frac{\phi'_0 H_{\text{inf}}^2}{f_X}$, and we have assumed that they have the same IR cut-off μ .

In the derivation of the asymmetry parameters above, Eq. (3.34) and Eq. (3.37), we have assumed that the only non-negligible contribution to the generated charge asymmetry is produced during the inflationary epoch. The conditions for the mechanism considered here may still be active during the radiation epoch, but the overall effect will be negligible as the push out-of-equilibrium is considered to be too small in later epochs; hence the total X charge is assumed to be conserved once inflation ends. One exception to this is the possibility of sphaleron redistribution which will violate both the SM B and L charges equally. The mutual dilution

of the charge and the entropy densities, after reheating, ensures there is no further dilution of the asymmetry parameter. No additional washout processes have been considered in the above derivation.

In the following section we utilise the known properties of sphaleron transitions to determine the distribution of X charges amongst fermionic species after the EWPT, if the reheating temperature is greater than the critical temperature ($T_c \sim 100$ GeV). How the sphaleron processes redistribute the X charge is dependent on the specific model being considered - the type of charge gauged, the associated anomalies, and the properties of the new fermion(s) introduced.

3.6 Replicating the Observed ρ_{DM}/ρ_B and η_B

Now that the X charge asymmetry parameter has been calculated we can determine under what conditions this model will replicate simultaneously the observed dark to luminous matter mass density ratio and the baryon asymmetry parameter. The generated X charge density can be decomposed into SM and dark matter components as follows,

$$n_X = n_X^{\text{SM}} + n_D , \quad (3.38)$$

The SM component will have an associated $B - L$ charge which will be reprocessed by the action of sphaleron transitions, before or at the EWPT, into a known fermionic distribution. The dark matter candidate considered here will be unaffected by the sphaleron transitions as it is assumed here to be a singlet under the electroweak interactions, although this does not have to be the case.

After the EWPT the $B - L$ charge will be distributed between B and L charges as follows; $(B - L)_{\text{SM}} = \frac{79}{28}B$ and $(B - L)_{\text{SM}} = -\frac{79}{51}L$. We require that the resultant SM baryon number asymmetry is consistent with that which is observed, given in Eq. (3.1). The baryon asymmetry parameter will be given by the following relation,

$$\eta_B = \epsilon(\eta_X^{\text{mixed-SM}} + \eta_X^{\text{unmixed-SM}}) , \quad (3.39)$$

where ϵ is a step function defined by,

$$\epsilon := \epsilon(T_{\text{rh}}) = \begin{cases} \frac{28}{79} & T_{\text{rh}} > T_c \\ 1 & T_{\text{rh}} < T_c \end{cases} , \quad (3.40)$$

Henceforth we will assume that the mixed anomalies only contribute to the SM sector, $\eta_X^{\text{mixed}} = \eta_X^{\text{mixed-SM}}$, as our dark matter candidate is sterile under the SM gauge groups.

It is assumed that the X charge density generated is initially uniformly distributed between each of the applicable fermion degrees of freedom, that is,

$$\eta_X^{SM} = \eta_X^{\text{mixed}} + \frac{\sum_i N_{SM}^i |q_{SM}^i|}{\sum_i N_{SM}^i |q_{SM}^i| + \sum_i N_D^i |q_D^i|} \eta_X^{\text{unmixed}} , \quad (3.41)$$

$$\eta_D = \frac{\sum_i N_D^i |q_D^i|}{\sum_i N_{SM}^i |q_{SM}^i| + \sum_i N_D^i |q_D^i|} \eta_X^{\text{unmixed}} , \quad (3.42)$$

where the index i corresponds to the particle species, N_i is the corresponding number of degrees of freedom, and q_i is the associated X charge. Therefore, the baryon asymmetry parameter defined above is given by,

$$\eta_B = \epsilon \eta_X^{SM} = \epsilon \left(\eta_X^{\text{mixed}} + \frac{\sum_i N_{SM}^i |q_{SM}^i|}{\sum_i N_{SM}^i |q_{SM}^i| + \sum_i N_D^i |q_D^i|} \eta_X^{\text{unmixed}} \right) . \quad (3.43)$$

The dark matter to luminous matter mass density ratio is given by,

$$\frac{\rho_D}{\rho_B} = \frac{m_\psi}{m_B} \frac{q_B \eta_D}{q_\psi \eta_B} = \frac{\eta_D}{q_\psi \eta_B} \left(\frac{m_\psi}{1 \text{ GeV}} \right) , \quad (3.44)$$

where we have assumed $m_B = 1 \text{ GeV}$ and $q_B = 1$. In the following analysis we will assume that this ratio is fixed to the observed value given in Eq. (3.3). Hence upon considering parameters that give the correct η_B we will also obtain the observed relic dark matter abundance. Now we wish to consider this framework in the two scenarios introduced earlier; namely, a gauged $B-L$ and a gauged B number, each including a single sterile fermion charged under the given group.

Case 1: $U(1)_{B-L}$ and a sterile fermion

In this scenario we must sum over all the SM fermions, $\sum_i N_{SM}^i |q_{SM}^i| = 21$, assuming no RH neutrinos have been added. Only the unmixed cubic anomaly contributes to the $B-L$ charge generation. Using these facts and Eq. (3.44) we derive the following dark matter to luminous matter mass density ratio,

$$\frac{\rho_D}{\rho_B} \approx \frac{1}{21\epsilon} \left(\frac{m_\psi}{1 \text{ GeV}} \right) , \quad (3.45)$$

where we have chosen $N_\psi = 1$.

In the following analysis we wish to consider our mechanism as the only source of both the dark matter and baryon asymmetry in the universe, and as such require that both Eq. (3.1) and Eq. (3.3) are satisfied. This immediately leads to the following prediction for the dark matter mass in this scenario,

$$m_\psi \approx 116\epsilon \text{ GeV} . \quad (3.46)$$

Hence the dark matter candidate ψ must have a mass $m_\psi \approx 41$ GeV, or $m_\psi \approx 116$ GeV; for $T_{\text{rh}} > T_c$ and $T_{\text{rh}} < T_c$ respectively. For consistency with the initial assumption that $m_\psi \ll H_{\text{inf}}$, the inflationary Hubble rate must be greater than $\sim 10^3$ GeV. Interestingly, the fixing of the ratio ρ_D/ρ_B sets the mass of the associated dark fermion, and is independent of the $B-L$ charge of the dark matter candidate. If the mass of the dark matter candidate is lower than 116ϵ GeV, then it cannot be the only component of the dark matter energy density of the universe and an additional component to the dark matter must be introduced, but we do not consider that scenario here. It should be noted that this mass relation assumes the correct baryon asymmetry parameter is generated, and hence the other parameters of the model are constrained, which we consider now.

The required replication of the observed baryon asymmetry and dark to luminous mass density ratio results in the following condition on the model parameters, for $\mu = H_0$,

$$\eta_B \approx \epsilon |\mathcal{A}_1|^{5/2} \frac{101}{21 + |q_\psi|} \frac{1}{\pi^{13} g_*} \frac{m_X^5}{f_X^5} e^{-3N_e} \left(\frac{\kappa}{\mu T_{\text{rh}}^2} \right)^{3/2} \quad (3.47)$$

$$\approx 3.5 \times 10^{-18} \text{ GeV}^{-1/2} \epsilon \frac{|\mathcal{A}_1|^{5/2}}{21 + |q_\psi|} \frac{m_X^5}{f_X^5} \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \left(\frac{\phi'_0}{f_X} \right)^{3/2}. \quad (3.48)$$

It is found that this can satisfy Eq. (3.1) for a wide range of parameter values. To see this we shall consider an example. Let us first assume ϕ'_0 takes its maximal value, $H_{\text{inf}} M_p$, to ensure maximal \mathcal{CP} violation. We will also identify the scale f_X with the inflationary Hubble rate, $\sim H_{\text{inf}}$, and the $B-L$ charge of the dark matter fermion to be $q_\psi = -1$. Under these assumptions we derive the following relation on the model parameters from Eq. (3.1) and Eq. (3.48),

$$\eta_B \approx 7 \times 10^9 \text{ GeV} \epsilon g_X^5 \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \Rightarrow \epsilon g_X^5 \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \approx 10^{-20} \text{ GeV}^{-1}, \quad (3.49)$$

which reduces to:

$$g_X \sim 10^{-4} \text{ GeV}^{-1/5} \frac{T_{\text{rh}}^{2/5}}{H_{\text{inf}}^{1/5}} \quad \text{or} \quad m_X \sim 10^{-4} \text{ GeV}^{-1/5} T_{\text{rh}}^{2/5} H_{\text{inf}}^{4/5}. \quad (3.50)$$

In this example, the satisfaction of this relation guarantees the correct baryon asymmetry and hence relic dark matter density. This relation leads to interesting constraints on the allowed parameter space. We find that for instantaneous reheating, $H_{\text{inf}} \sim \frac{(T_{\text{rh}}^{\text{max}})^2}{M_p}$, we require $g_X \sim 0.4$. This is likely larger than that assumed in the model ($g_X \ll 1$), and means that this relation can be satisfied for almost all allowed reheating temperatures for a given inflationary Hubble rate. Hence, the reheating epoch must not be instantaneous and the reheating temperature cannot conflict with BBN constraints ($T_{\text{rh}} > \mathcal{O}(1)$ MeV). From the mass of the dark matter candidate and constraints on the inflationary potential we require that the inflationary Hubble

rate is in the range $10^{14} \text{ GeV} > H_{\text{inf}} > 10^3 \text{ GeV}$. Taking this into account, we see that the size of the coupling g_X that can satisfy this relation lies in the range $10^{-8} \lesssim g_X \lesssim 0.4$, with the lowest value corresponding to the input parameters $H_{\text{inf}} = 10^{14} \text{ GeV}$ and $T_{\text{rh}} = \mathcal{O}(1) \text{ MeV}$. Interestingly, the mass of the gauge boson can be as low as $m_X \sim \mathcal{O}(1) \text{ MeV}$, when taking $H_{\text{inf}} = 10^3 \text{ GeV}$ and $T_{\text{rh}} = \mathcal{O}(1) \text{ MeV}$, leading to a coupling of order 10^{-6} .

It could be possible to probe experimentally the areas of parameter space in which the X boson mass is lowest, although this is made difficult by the associated tiny coupling constant. The generically small couplings would also make it difficult to detect the predicted dark matter candidate.

If we now consider the IR cut-off to be the Hubble rate at the beginning of BBN, $H_{\text{BBN}} \approx 10^{-25} \text{ GeV}$, the condition becomes,

$$\eta_B \approx \epsilon |\mathcal{A}_1|^{5/2} \frac{101}{21 + |q_\psi|} \frac{1}{\pi^{13} g^*} \frac{m_X^5}{f_X^5} e^{-3N_e} \left(\frac{\kappa}{\mu T_{\text{rh}}^2} \right)^{\frac{3}{2}} \quad (3.51)$$

$$\approx 10^{-43} \text{ GeV}^{-1/2} \epsilon \frac{|\mathcal{A}_1|^{5/2}}{21 + |q_\psi|} \frac{m_X^5}{f_X^5} \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \left(\frac{\phi'_0}{f_X} \right)^{\frac{3}{2}}. \quad (3.52)$$

This choice of cut-off provides a significantly more constrained result, but can still be satisfied with an appropriate choice of parameters. Considering the same example model discussed above,

$$\eta_B \approx 2 \times 10^{-16} \text{ GeV} \epsilon g_X^5 \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \Rightarrow \epsilon g_X^5 \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \approx 4 \cdot 10^5 \text{ GeV}^{-1}, \quad (3.53)$$

which reduces to,

$$g_X \sim 10 \text{ GeV}^{-1/5} \frac{T_{\text{rh}}^{2/5}}{H_{\text{inf}}^{1/5}} \quad \text{or} \quad m_X \sim 10 \text{ GeV}^{-1/5} T_{\text{rh}}^{2/5} H_{\text{inf}}^{4/5}. \quad (3.54)$$

Considering instantaneous reheating requires a coupling of $g_X \sim 400$, this means that an extended reheating epoch is needed for consistency with our assumption that $g_X \ll 1$. The size of the couplings that can satisfy this relation lie in the range $10^{-3} \lesssim g_X \ll 1$, where the lower bound is for $H_{\text{inf}} = 10^{14} \text{ GeV}$ and $T_{\text{rh}} = \mathcal{O}(1) \text{ MeV}$. This allowed window is much smaller than found in the previous case. The minimum mass of the boson is much higher with $m_X \sim 160 \text{ GeV}$, when $H_{\text{inf}} = 10^3 \text{ GeV}$ and $T_{\text{rh}} = \mathcal{O}(1) \text{ MeV}$. Due to the higher couplings in this case, the model is likely much easier to constrain with experiments in the lower m_X regime, hence a very large H_{inf} and low T_{rh} would be required.

Case 2: $U(1)_B$ and a sterile baryon

If we now consider a gauged baryon number extension to the SM we must sum over all of the baryonic degrees of freedom, $\sum_i N_{SM}^i |q_{SM}^i| = 12$. In this scenario the contributions of the mixed anomalies $SU(2)^2 \times U(1)_B$ and $U(1)_Y^2 \times U(1)_B$ must be included, which generate a net charge only in the form of luminous matter. Hence, we find that the dark matter to luminous matter mass density ratio is given by,

$$\frac{\rho_D}{\rho_B} = \frac{1}{\epsilon} \frac{N_\psi}{12 + N_\psi |q_\psi|} \frac{|\mathcal{A}_1|^{5/2}}{\frac{12|\mathcal{A}_1|^{5/2}}{N_\psi |q_\psi| + 12} \frac{m_X^5}{f_X^5} + |\mathcal{A}_2|^{5/2} g_1^5 + 3|\mathcal{A}_3|^{5/2} g_2^5} \frac{m_X^5}{f_X^5} \left(\frac{m_\psi}{1 \text{ GeV}} \right) \quad (3.55)$$

$$\approx \frac{1}{\epsilon} \frac{|q_\psi|^{15/2}}{12|q_\psi|^{15/2} \frac{m_X^5}{f_X^5} + |q_\psi| + 12} \frac{m_X^5}{f_X^5} \left(\frac{m_\psi}{1 \text{ GeV}} \right), \quad (3.56)$$

where $m_B = 1 \text{ GeV}$ and $q_B = 1$ have been set. In the second line we have taken $g_1^2 \simeq \frac{4\pi}{60}$ and $g_2^2 \simeq \frac{4\pi}{29}$, and used the anomaly values given above: $\mathcal{A}_2 = 3/2$ and $\mathcal{A}_3 = -3/2$. Upon rearranging, and requiring Eq. (3.3), we find the following expression for the mass of the dark matter candidate,

$$m_\psi \approx \epsilon \frac{f_X^5}{m_X^5} \frac{11(12|q_\psi|^{15/2} \frac{m_X^5}{f_X^5} + |q_\psi| + 12)}{2|q_\psi|^{15/2}} \text{ GeV}, \quad (3.57)$$

which simplifies to,

$$m_\psi \approx \frac{70}{\epsilon g_X^5} \text{ GeV}, \quad (3.58)$$

when taking $|q_\psi| = 1$ and $g_X \ll 1$. Therefore, in this scenario we require that $H_{\text{inf}} \gtrsim 10^7 \text{ GeV}$, which implies that the mass of the dark matter candidate lies in the range $10^6 \text{ GeV} \lesssim m_\psi \ll H_{\text{inf}}$. The required dark matter mass is greater than that found in the previous case due to the additional contributions to the luminous sector from the mixed anomalies, and the interplay of this with the mass density ratio.

The associated constraint imposed by the observed baryon asymmetry is given by,

$$\eta_B \approx \epsilon \frac{\mathcal{A}}{\pi^{13} g_*} e^{-3N_e} \left(\frac{\kappa}{\mu T_{\text{rh}}^2} \right)^{\frac{3}{2}} \quad (3.59)$$

$$\approx 3 \times 10^{-19} \text{ GeV}^{-1/2} \epsilon \mathcal{A} \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \left(\frac{\phi'_0}{f_X} \right)^{\frac{3}{2}}, \quad (3.60)$$

where $\mathcal{A} = \left(\frac{12|q_\psi|^{15/2}}{|q_\psi| + 12} \frac{m_X^5}{f_X^5} + |\mathcal{A}_2|^{5/2} g_1^5 + 3|\mathcal{A}_3|^{5/2} g_2^5 \right)$. As we assume $g_X^2 \ll 1$ this reduces to $\mathcal{A} \sim 1$, except when considering very large q_ψ , which we do not here. Similar to the $U(1)_{B-L}$ scenario, we assume ϕ'_0 takes its maximal value, $H_{\text{inf}} M_p$, and identify the scale f_X with the inflationary Hubble rate, H_{inf} . Here the B charge of the dark matter fermion is chosen to be $q_\psi = 1$. Under these assumptions, we derive the following relation on the model parameters

from Eq. (3.1) and Eq. (3.48),

$$\eta_B \approx 4 \times 10^8 \text{ GeV} \epsilon \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \rightarrow \epsilon \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \approx 2 \cdot 10^{-19} \text{ GeV}^{-1} \rightarrow H_{\text{inf}} \approx \frac{2 \cdot 10^{-19} \text{ GeV}^{-1}}{\epsilon} T_{\text{rh}}^2. \quad (3.61)$$

Interestingly, this relation is only dependent on the inflationary Hubble rate and the reheating temperature. Upon further inspection it appears to be almost consistent with instantaneous reheating, or a very efficient reheating process, hence in this scenario a close to maximal reheating temperature is required for any given inflationary Hubble rate.

From the analysis of the allowed dark matter mass range above, we find that the inflationary rate must be in the window $10^{14} \text{ GeV} \gtrsim H_{\text{inf}} \gtrsim 10^7 \text{ GeV}$. The corresponding allowed range of reheating temperatures is then $10^{16} \text{ GeV} \gtrsim T_{\text{rh}} \gtrsim 7 \cdot 10^{12} \text{ GeV}$. The Eq. (3.58) and allowed mass range also give us a lower bound on the allowed coupling, which we find is $g_X^2 \sim 10^{-5}$, although this can be relaxed for larger dark matter charge values. These constraints also imply that the mass of the new X gauge boson must be at least greater than $\sim 50 \text{ TeV}$, which suppresses the ability to detect the new boson and dark matter fermion. These values of the gauge coupling and boson mass range are well out of the range of any current collider experiments [466].

The constraint obtained when taking the IR cut-off to be $\mu = H_{\text{BBN}} \approx 10^{-25} \text{ GeV}$ is,

$$\eta_B \approx 10^{-44} \text{ GeV}^{-1/2} \epsilon \mathcal{A} \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \left(\frac{\phi'_0}{f_X} \right)^{\frac{3}{2}}. \quad (3.62)$$

Considering the same assumptions as for the previous cut-off we find,

$$\eta_B \approx 10^{-17} \text{ GeV} \epsilon \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \rightarrow \epsilon \frac{H_{\text{inf}}}{T_{\text{rh}}^2} \approx 6 \cdot 10^6 \text{ GeV}^{-1} \rightarrow H_{\text{inf}} \approx \frac{6 \cdot 10^6 \text{ GeV}^{-1}}{\epsilon} T_{\text{rh}}^2. \quad (3.63)$$

The constraints on the mass of the dark matter particle and coupling are the same for this case, but now the relation between the inflationary Hubble rate and reheating temperature has changed. Now we require an extended reheating epoch, with a maximum reheating temperature of $T_{\text{rh}} \sim 4 \text{ TeV}$ for $H_{\text{inf}} \sim 10^{14} \text{ GeV}$ and minimum of $T_{\text{rh}} \sim 1 \text{ GeV}$ when $H_{\text{inf}} \sim 10^7 \text{ GeV}$. This change in the allowed reheating temperatures is the only alteration caused by the use of the other cut-off.

In summary, the ability to detect the introduced gauge boson appears likely to be difficult due to the generic requirement of a high mass, except in special cases, and small coupling. The dark matter candidate would be more easily detectable in the case of the gauged $B - L$ compared to the gauged B scenario due to it having a significantly smaller mass, which is of order $10 - 100 \text{ GeV}$. Although as the dark matter candidate can only interact with the SM particles via the $U(1)_X$ gauge boson, the ability to detect it is highly dependent on the gauge boson mass and

coupling, for which there is a wide range of possibilities. It may be possible to obtain constraints on the parameter space from Z' searches [41, 77, 466–477].

3.7 Conclusions and Future Prospects

In this chapter, we have investigated a model for simultaneous generation of luminous and dark matter during the inflationary epoch, achieved through the introduction of an anomalous gauge interaction and sterile fermion to the SM. It has been found that this scenario for cogenesis can successfully reproduce observations for the two possible cases considered - a gauged B and a gauged $B - L$ charge.

In each scenario, we considered the parameter spaces that correctly predict both the dark matter to baryonic matter density ratio and baryon asymmetry. Interestingly, for the $U(1)_{B-L}$ extension we obtain a prediction for the mass of the dark matter candidate which is independent of the other choice of parameters, assuming a given relation between g_X , T_{rh} and H_{inf} . While in the $U(1)_B$ case, we find the model replicates the observed η_B as long as a relation between the reheating temperature and inflationary Hubble rate is adhered to, and the coupling satisfies $g_X^2 > 10^{-5}$.

The general mechanism for cogenesis developed here can be applied to more complex models involving other or extra anomalous gauge symmetries, as well as additional sterile or non-sterile fermionic states. It is possible that these additions could lead to a loosening of the constraints on the allowed parameters imposed by the observed matter-antimatter asymmetry, through extra contributions to the generation of luminous matter or dark matter. This would likely lead to altered requirements on the mass of the dark matter candidate.

Our general framework could also provide a mechanism for magnetogenesis, through the production of universal hypermagnetic fields via the $U_Y(1)$ CS term [478]. One could craft a model that not only generates this but also the particle asymmetries, providing an extra route for constraining the allowed parameter space.

Further to this, the study of the associated collider phenomenology of these models is of interest. Although, this is made difficult by the possibility of very small couplings and high masses. Despite this, there are certain areas of the parameter space that can already be constrained by results from collider experiments.

Chapter 4

Baryogenesis During Reheating via the Ratchet Mechanism

Another interesting and relatively unorthodox approach to generating the baryon asymmetry of the universe is via the inflaton during the reheating epoch. Unlike the dilution problems associated with inflationary cogenesis considered above, the difficulties with a reheating scenario are related to the high level of uncertainty and complexity of the dynamics associated with the reheating phase. The exact nature of the reheating epoch is mostly unknown, but it is a period dominated by the inflaton dynamics, and as such is strongly related to the properties of the inflaton and inflationary potential. To try to alleviate this issue and for simplicity, we will consider a Starobinsky inflationary scenario, which converges to the usual $\mu^2\Phi^2$ potential during reheating.

We propose a new scenario for Baryogenesis during the reheating epoch that utilises the Ratchet mechanism, a model inspired by molecular motors in biological systems, and their ability to generate directed motion. This is achieved through the correlated behaviour between the inflaton and a complex scalar baryon. If the inflaton and the scalar baryon couple via a derivative coupling, the behaviour of the scalar baryon phase θ is found to be analogous to that of the forced pendulum, potentially producing a non-vanishing value of $\dot{\theta}$ which is necessary to generate a non-zero baryon number density [4].

4.1 The Reheating Epoch and Starobinsky Inflation

A consequence of the rapid expansion that occurs during inflation is that the energy densities of SM particles are diluted to negligible quantities, so in order to explain the observed matter energy densities today they must be produced dynamically some time after inflation ends. This phenomena is known as reheating, and involves the conversion of the energy density stored in the inflationary potential into matter, through the decay of the inflaton [371, 479–488]. This process begins once the slow roll conditions are violated, that is, when the inflationary potential is no longer flat enough to support the inflationary scenario. In the canonical model, the almost homogeneous inflaton field oscillates coherently in its potential, being damped by Hubble expansion and a frictional term associated with the decay of the inflaton into matter particles and radiation. These processes are occurring in an out-of-equilibrium setting that continues until the decay rate becomes the dominant source of damping. For a $m^2\phi^2$ like inflaton potential during reheating, the coherent oscillations of the inflaton field induce an approximate matter dominated epoch [482]. Once the decay rate becomes the main source of damping, the universe will be dominated by relativistic particles produced from the distribution of the inflaton energy density into SM particles, and possibly other exotic species, and will begin to thermalise.

Once reheating is completed the universe thermalises and enters a radiation dominated epoch, with an initial temperature, also known as the reheating temperature, which is related to the initial energy in the inflationary potential and the decay rate of the inflaton. Faster inflaton decay leads to a more efficient transfer of energy from the inflationary sector to the radiation sector, and hence produces a higher reheating temperature. The exact details of the reheating epoch are complicated and mostly unknown, and as such it is an active area of scientific exploration, with possible probes of its properties considered [489–491]. We can constrain the maximal and minimal reheating temperatures using the maximum allowed inflationary scale and successful BBN, respectively, although this still permits a large range of reheating temperatures [492]. The uncertainty associated with this epoch lends itself to being an interesting source of answers to the mysteries surrounding cosmology and particle physics, particularly Baryogenesis [493], which we shall discuss in this chapter.

It is possible to determine a naive upper limit on the reheating temperature for a given inflationary mechanism by considering the Friedmann equations, and the approximate energy density contained within the inflaton potential,

$$\rho_{\text{inf}} \simeq U(\Phi_i) \simeq 3M_p^2 H_{\text{inf}}^2, \quad (4.1)$$

where Φ_i is the initial value of the inflaton, $U(\Phi)$ is the inflaton potential, and $M_p = 1/\sqrt{8\pi G}$ is the reduced Planck mass. This energy density can then be equated to that associated with a radiation epoch with characteristic temperature T_{rh} , assuming efficient thermalisation [128],

which is given by,

$$\rho_{\text{rh}} = \frac{\pi^2}{30} g_* T_{\text{rh}}^4 . \quad (4.2)$$

This gives a maximum reheating temperature of,

$$T_{\text{rh}} \simeq \left(\frac{90}{\pi^2 g_*} \right)^{1/4} \sqrt{M_p H_{\text{inf}}} . \quad (4.3)$$

For this temperature to be achieved, one must assume instantaneous and lossless decay of the inflaton into SM particles. The inflaton decay can be parametrised in the inflaton's equations of motion through the introduction of the decay width Γ , acting as an additional damping term,

$$\ddot{\Phi} + 3H\dot{\Phi} + \Gamma\dot{\Phi} + V'(\Phi) = 0 . \quad (4.4)$$

The value taken by Γ is dependent on the fields that the inflaton decays into and the strength of the associated interactions. The physical implications of the magnitude of Γ is that once $H \sim \Gamma$, the inflaton decays will dominate and reheating will end, leading to a radiation epoch with the characteristic temperature dictated by the properties of Γ . The length of the reheating epoch will be approximately Γ^{-1} . The associated reheating energy density can be found using the Born approximation,

$$\rho_{\text{rh}} \simeq 3\Gamma^2 M_p^2 , \quad (4.5)$$

which upon comparing to the energy density at the end of reheating, we find,

$$T_{\text{rh}} \simeq \left(\frac{90}{\pi^2 g_*} \right)^{1/4} \sqrt{\Gamma M_p} . \quad (4.6)$$

The number of e-folds of expansion that occur during reheating are,

$$N_{\text{rh}} = \frac{1}{3} \ln \left(\frac{\rho_{\text{inf}}}{\rho_{\text{rh}}} \right) \simeq \frac{2}{3} \ln \left(\frac{H_{\text{inf}}}{\Gamma} \right) . \quad (4.7)$$

It should be noted that the T_{rh} given in Eq. (4.6) is the maximum temperature of the radiation dominated universe, but it may not necessarily be the hottest temperature achieved after inflation has ended, rather temperatures during reheating could be higher. This can have interesting implications for phase transitions and production of relics.

4.1.1 Starobinsky Inflation

Many models for the mechanism of inflation have been proposed, each of which must produce observational predictions consistent with the constraints depicted in Figure 2.3. One model

which is in good agreement with these constraints is the Starobinsky, or R^2 , type models [110, 126, 494]. Given this agreement with the data, we shall consider this as the inflationary scenario in our investigation.

The Starobinsky action is the following,

$$S = \int dx^4 \sqrt{-g} \frac{M_p^2}{2} \left(R - \frac{R^2}{6\mu^2} \right), \quad (4.8)$$

where in the Einstein frame of the scalar version we have the following inflationary potential,

$$U(\Phi) = \frac{3\mu^2 M_p^2}{4} \left(1 - e^{-\sqrt{2/3}\Phi/M_p} \right)^2, \quad (4.9)$$

where $\mu = (1.3 \times 10^{-5})M_p$ is the inflaton mass [495]. From this potential the slow roll parameters and corresponding predictions can be found. Using the tools presented in Chapter 2, the Starobinsky model produces the following inflationary predictions,

$$n_s \simeq 1 - \frac{2}{N} \quad \text{and} \quad r \simeq \frac{12}{N^2}, \quad (4.10)$$

where N is the number of inflationary e-folds.

The reheating period in the Starobinsky model approaches the usual $\frac{1}{2}\mu^2\Phi^2$, leading to this epoch being characterized as an approximate matter dominated epoch [482, 495, 496]. This can be seen in the expansion of the inflationary potential around Φ ,

$$U(\Phi) = \frac{1}{2}\mu^2\Phi^2 - \frac{\mu^2}{\sqrt{6}M_p}\Phi^3 + \dots \quad (4.11)$$

where the first term dominates for $\Phi < M_p$. From numerical calculations it is found that in the Starobinsky model the value the inflaton field takes at the beginning of the reheating epoch is [496],

$$\Phi_i = \Phi(t_i) = 0.62 M_p, \quad (4.12)$$

with a corresponding Hubble parameter of,

$$H_i = H(t_i) = 6.2 \times 10^{12} \text{ GeV}. \quad (4.13)$$

In the proceeding analysis, we will assume that the inflaton velocity $\dot{\Phi}(t_i)$ at this point in time is approximately zero for simplicity.

4.2 Baryogenesis during Reheating via the Ratchet Mechanism

Baryogenesis during reheating has been an area of interest in the literature for some time, given the complicated and uncertain nature of the epoch. In what follows we shall outline a new mechanism for Baryogenesis in which baryon number generation is driven by the oscillations of the inflaton field. As with any successful model of Baryogenesis that conserves \mathcal{CPT} , the Sakharov conditions must be satisfied in order to reproduce the observed value of the baryon asymmetry parameter,

$$\eta_B \simeq 8.5 \cdot 10^{-11} . \quad (4.14)$$

Of the many Baryogenesis models that have been proposed, there is a class that generates the baryon asymmetry via the coherent time-evolution of a complex scalar field that carries a baryon number charge [175, 231, 481, 493, 497–500]. In this work we consider a similar scenario in the setting of reheating, with a new scalar baryon coupled to the coherently oscillating inflaton field.

4.2.1 The Ratchet Mechanism

The new Baryogenesis mechanism we propose here acts during the reheating epoch, and is inspired by the ratchet models that describe molecular motors in biological systems [501, 502]. An example of this is the directed motion of myosin molecules along actin filaments which is achieved through cyclic chemical reactions that act as a driving force. The model we propose here utilises an analogous ratchet mechanism, in which the reflection symmetry of a scalar potential is broken by its interaction with the inflaton, with the inflaton's coherent oscillations providing a driving force. This driving force is supplied by the oscillation of the inflaton in its potential during reheating, while the position of the motor is embodied in the phase θ of the complex scalar baryon field. These two scalar fields interact via a derivative coupling which violates \mathcal{CP} , breaking the reflection symmetry of the scalar baryon's potential. The form of the interaction between the two scalars is similar to that in the Baryogenesis mechanism considered in Ref. [503], although in their case \mathcal{CPT} violation is required.

4.2.2 Description of the Model

We construct a model consisting of two scalar fields - a real scalar field Φ that we identify as the inflaton, and a complex scalar baryon ϕ . In the ensuing analysis we assume that the dynamics during reheating are dominated by these two scalars and only consider interactions of the inflaton with SM fields via an effective friction term. Our model is described by the

following action,

$$S = \int d^4x \sqrt{-g} \left[g_{\mu\nu} \partial^\mu \phi^* \partial^\nu \phi - V_0(\phi, \phi^*) + \frac{1}{2} g_{\mu\nu} \partial^\mu \Phi \partial^\nu \Phi - U(\Phi) + \frac{i}{\Lambda} g_{\mu\nu} (\phi^* \overleftrightarrow{\partial}^\mu \phi) \partial^\nu \Phi \right], \quad (4.15)$$

where $U(\Phi)$ is the inflationary potential, which we take to be the Starobinsky potential given in Eq. (4.9). The scalar baryon potential, $V_0(\phi, \phi^*)$, is defined as,

$$V_0(\phi, \phi^*) = \lambda \phi^* \phi (\phi - \phi^*) (\phi^* - \phi) + \dots, \quad (4.16)$$

where the ellipses denote terms that depend only on the product $\phi^* \phi$, which are not relevant to the dynamics in this mechanism.

The coupling between the inflaton and the scalar baryon is introduced as, $\mathcal{L}_{int} = -\frac{1}{\Lambda} j_B^\mu \partial_\mu \Phi = -\frac{\phi_r^2}{\Lambda} \partial^\mu \theta \partial_\mu \Phi$. This is a dimension five operator, and as such is suppressed by the mass scale Λ which is the energy cut-off at which the effective description of this scalar coupling breaks down. This interaction term also violates \mathcal{C} and \mathcal{CP} , which is a necessary ingredient for successful Baryogenesis.

Upon observation it can be seen that if $\lambda = 0$, the action will be invariant under the global $U(1)$ symmetry defined by the transformation $(\phi, \phi^*) \rightarrow (e^{i\alpha} \phi, e^{-i\alpha} \phi^*)$, where α is a constant. We identify the charge associated with this $U(1)$ symmetry as the baryon number B , where the complex scalar field ϕ is a baryon to which we assign unit baryonic charge. From the action we can calculate the corresponding baryon number current, or Noether current for the baryon symmetry, which is found to be,

$$j_B^\mu = i(\phi \partial^\mu \phi^* - \phi^* \partial^\mu \phi). \quad (4.17)$$

We now consider the following polar coordinate parametrisation of the ϕ field,

$$\phi = \frac{1}{\sqrt{2}} \phi_r e^{i\theta}. \quad (4.18)$$

Under the global baryon number transformation, the phase θ transforms as $\theta \rightarrow \theta + \alpha$, while ϕ_r is invariant. In this parametrisation the baryon number density, which corresponds to the time component of Eq. (4.17), is given by,

$$n_B = j^0 = \phi_r^2 \dot{\theta}. \quad (4.19)$$

This implies that within the framework of our mechanism we must produce a non-zero $\dot{\theta}$ to have a net baryon asymmetry generated. Rewriting the scalar baryon potential in terms of this reparametrisation, ϕ_r and θ , we obtain the following,

$$V(\phi_r, \theta) = V_0(\phi, \phi^*) = \lambda \phi_r^4 \sin^2 \theta + \dots \quad (4.20)$$

where the ellipses now denote terms that depend only on ϕ_r . In the rest of our analysis, we assume that the terms that only depend on ϕ_r in $V(\phi_r, \theta)$ are such that they keep ϕ_r approximately fixed to a constant non-zero value, and that only the dynamics of the phase θ need be considered. The charge conjugation symmetry \mathcal{C} is given by $\mathcal{C} : \phi \rightarrow \phi^*$, or $\theta \rightarrow -\theta$, so \mathcal{C} is conserved in this potential. B invariance is related to the translational invariance of the phase θ , which is clearly violated by this potential, assuming $\lambda \neq 0$.

Therefore, in the new parametrisation of the scalar baryon the action takes the form,

$$S = \int d^4x \sqrt{-g} \left[\frac{\phi_r^2}{2} g_{\mu\nu} \partial^\mu \theta \partial^\nu \theta - \lambda \phi_r^4 \sin^2 \theta + \frac{1}{2} g_{\mu\nu} \partial^\mu \Phi \partial^\nu \Phi - U(\Phi) - \frac{\phi_r^2}{\Lambda} g_{\mu\nu} \partial^\mu \theta \partial^\nu \Phi \right]. \quad (4.21)$$

Seeing as we wish to consider the cosmological setting of reheating we take the gravitational metric $g_{\mu\nu}$ to be the flat FRW metric,

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = dt^2 - a^2(t) dx^2. \quad (4.22)$$

Given this isotropic and homogeneous background, we extend this assumption to the properties of the scalar baryon and inflaton, for which spatial variation will be ignored in our analysis. In this background, the action of our model reads,

$$S = \int dt a(t)^3 \left[\frac{\phi_r^2}{2} \dot{\theta}^2 - \lambda \phi_r^4 \sin^2 \theta + \frac{1}{2} \dot{\Phi}^2 - U(\Phi) - \frac{\phi_r^2}{\Lambda} \dot{\theta} \dot{\Phi} \right]. \quad (4.23)$$

Therefore, in our model, the Sakharov conditions are satisfied in the following ways. Firstly, B violation is achieved by the scalar baryon potential $V(\phi_r, \theta)$. Secondly, the derivative coupling changes sign under both \mathcal{C} and \mathcal{CP} . We have ignored the spatial dependence of the scalar fields, so the parity symmetry \mathcal{P} is always conserved by this derivative coupling term. The required push out-of-equilibrium will be provided by the reheating epoch, induced by the coherent oscillation of the inflaton field. During reheating the derivative coupling between the scalar baryon and inflaton can lead to a kind of resonance effect when the respective potentials are of a similar order. As we shall see, when this is the case we can observe directed motion in the scalar baryon phase, and hence a non-zero baryon number density.

Now, we can carry out the calculations required to determine the baryon number density generated in this framework. In our analysis we will take the initial phase of the scalar baryon to be zero, placing the scalar baryon initially in the minimum of its potential. To ensure there is no initial bias between matter and antimatter we assume that the initial phase velocity $\dot{\theta}$ is zero.

4.3 Analytical Evaluation

We shall now find an analytical solution for the scalar baryon phase equation of motion so that we can determine the region of parameter space where we obtain driven motion, and hence have a non-zero baryon number density. Firstly, we find the equations of motion for θ and Φ , using the action presented in Eq. (4.23),

$$(\ddot{\Phi} + 3H\dot{\Phi}) + \left(\Gamma\dot{\Phi} + \frac{dU(\Phi)}{d\Phi} \right) - \frac{\phi_r^2}{\Lambda}(\ddot{\theta} + 3H\dot{\theta}) = 0, \quad (4.24)$$

$$(\ddot{\theta} + 3H\dot{\theta}) - \frac{1}{\Lambda}(\ddot{\Phi} + 3H\dot{\Phi}) = 0, \quad (4.25)$$

where $\Gamma\dot{\Phi}$ is the inflaton friction term, added in by hand, which encapsulates the decay of the inflaton into SM or mediator particles during reheating, and H is the Hubble parameter. After some rearrangement, the above equations read,

$$\left(1 - \frac{\phi_r^2}{\Lambda^2} \right) (\ddot{\Phi} + 3H\dot{\Phi}) + \left(\Gamma\dot{\Phi} + \frac{dU(\Phi)}{d\Phi} \right) + \frac{\lambda\phi_r^4}{\Lambda} \sin(2\theta) = 0, \quad (4.26)$$

$$\left(1 - \frac{\phi_r^2}{\Lambda^2} \right) (\ddot{\theta} + 3H\dot{\theta}) + \frac{1}{\Lambda} \left(\Gamma\dot{\Phi} + \frac{dU(\Phi)}{d\Phi} \right) + \lambda\phi_r^2 \sin(2\theta) = 0. \quad (4.27)$$

The $\left(1 - \frac{\phi_r^2}{\Lambda^2} \right)$ factor can be absorbed through rescaling of the parameters Γ , μ^2 in $U(\Phi)$, and λ , as follows,

$$\tilde{\Gamma} \equiv \frac{\Gamma}{1 - \phi_r^2/\Lambda^2}, \quad \tilde{\mu}^2 \equiv \frac{\mu^2}{1 - \phi_r^2/\Lambda^2}, \quad \tilde{\lambda} \equiv \frac{\lambda}{1 - \phi_r^2/\Lambda^2}, \quad (4.28)$$

and hence the equations simplify to,

$$(\ddot{\Phi} + 3H\dot{\Phi}) + \left(\tilde{\Gamma}\dot{\Phi} + \frac{d\tilde{U}(\Phi)}{d\Phi} \right) + \frac{\tilde{\lambda}\phi_r^4}{\Lambda} \sin(2\theta) = 0, \quad (4.29)$$

$$(\ddot{\theta} + 3H\dot{\theta}) + \frac{1}{\Lambda} \left(\tilde{\Gamma}\dot{\Phi} + \frac{d\tilde{U}(\Phi)}{d\Phi} \right) + \tilde{\lambda}\phi_r^2 \sin(2\theta) = 0. \quad (4.30)$$

During reheating in the Starobinsky inflationary model, the oscillation of the massive inflaton gives rise to an approximate matter dominated epoch, during which the Hubble parameter H

is approximately given by,

$$H = \frac{\dot{a}}{a} \approx \frac{2}{3t} . \quad (4.31)$$

We shall also assume that during reheating the Starobinsky potential can be expressed approximately as,

$$\tilde{U}(\Phi) \approx \frac{1}{2} \tilde{\mu}^2 \Phi^2 \quad \rightarrow \quad \frac{d\tilde{U}(\Phi)}{d\Phi} \approx \tilde{\mu}^2 \Phi , \quad (4.32)$$

as found in the expansion given in Eq. (4.11). Note that early on in the reheating epoch the full potential should be used. This approximation is valid towards the end of the epoch, when the inflaton is oscillating near the potential minimum. The equations of motion are now,

$$\left(\ddot{\Phi} + \frac{2}{t} \dot{\Phi} \right) + \left(\tilde{\Gamma} \dot{\Phi} + \tilde{\mu}^2 \Phi \right) + \frac{\tilde{\lambda} \phi_r^4}{\Lambda} \sin(2\theta) = 0 , \quad (4.33)$$

$$\left(\ddot{\theta} + \frac{2}{t} \dot{\theta} \right) + \frac{1}{\Lambda} \left(\tilde{\Gamma} \dot{\Phi} + \tilde{\mu}^2 \Phi \right) + \tilde{\lambda} \phi_r^2 \sin(2\theta) = 0 . \quad (4.34)$$

In this form we can now begin solving these equations.

Behaviour of the Inflaton

We wish for the inflaton's motion to be unaffected by the dynamics of θ . This ensures that the properties of the reheating epoch are retained and the approximation given in Eq. (4.31) remains valid. To do so, we assume that the $\sin(2\theta)$ term in Eq. (4.33) can be neglected - the condition for this assumption shall be provided below. The equation of motion of the inflaton Φ then becomes,

$$\ddot{\Phi} + \left(\frac{2}{t} + \tilde{\Gamma} \right) \dot{\Phi} + \tilde{\mu}^2 \Phi = 0 . \quad (4.35)$$

This equation can be easily solved in the case when $\tilde{\Gamma} \ll \tilde{\mu}$, which is a valid assumption in our scenario. The approximate solution to this equation is,

$$\Phi(t) = \Phi_i \left(\frac{t_i}{t} \right) e^{-\tilde{\Gamma}(t-t_i)/2} \cos[\tilde{\mu}(t-t_i)] , \quad (4.36)$$

where t_i is the time at which the reheating epoch begins, and $\Phi_i = \Phi(t_i)$. This solution indicates that the motion of $\Phi(t)$ is oscillatory, with an angular frequency $\tilde{\mu}$, and is damped by $e^{-\tilde{\Gamma}t/2}$, coming from the $\tilde{\Gamma}$ term in the equation of motion, while the amplitude is also attenuated as a function of $1/t$ from the Hubble damping term.

Now that we have this solution it is possible to find a simple relation describing the assumption that the $\sin(2\theta)$ term can be neglected in the equation of motion of Φ . This requires the following

relation be satisfied,

$$\frac{\tilde{\lambda}\phi_r^4}{\Lambda} \ll \text{amplitude of } \tilde{\mu}^2\Phi(t) . \quad (4.37)$$

This should be true throughout the reheating epoch, but is sufficient to be true at the end of reheating due to the damping of the inflaton's amplitude, predominantly by Hubble damping,

$$\frac{\tilde{\lambda}\phi_r^4}{\Lambda} \ll \tilde{\mu}^2\Phi_i \left(\frac{H_f}{H_i} \right) , \quad (4.38)$$

where t_f is the time at the end of the reheating epoch, and H_i and H_f are respectively the values of the Hubble parameter at t_i and t_f .

Behaviour of the Scalar Baryon

Now that we have determined the dynamics of the inflaton during reheating, we can utilise the solution to find an analytical solution for the phase of the scalar baryon. As found above, the equation of motion for θ is,

$$\left(\ddot{\theta} + \frac{2}{t}\dot{\theta} \right) + \tilde{\lambda}\phi_r^2 \sin(2\theta) + \frac{1}{\Lambda} \left[\tilde{\Gamma}\dot{\Phi}(t) + \tilde{\mu}^2\Phi(t) \right] = 0 . \quad (4.39)$$

Since the amplitude of $\tilde{\Gamma}\dot{\Phi}(t)$ is suppressed compared to the amplitude of $\tilde{\mu}^2\Phi(t)$ by a factor of approximately $\tilde{\Gamma}/\tilde{\mu}$, we can drop the $\tilde{\Gamma}\dot{\Phi}(t)$ term. Doing this, and defining some new parameters the equation of motion for θ becomes,

$$\left(\ddot{\theta} + \frac{2}{t}\dot{\theta} \right) + p \sin(2\theta) + q(t) \cos[\tilde{\mu}(t - t_i)] = 0 , \quad (4.40)$$

where we have defined,

$$p = \tilde{\lambda}\phi_r^2, \quad q(t) = \frac{\tilde{\mu}^2\Phi_i}{\tilde{\Lambda}} \left(\frac{t_i}{t} \right) e^{-\tilde{\Gamma}(t-t_i)/2} . \quad (4.41)$$

This equation is not simple to solve, so we shall first consider a few possible scenarios to determine some of its properties. First, consider the case when $p \ll q(t)$, we now have,

$$\left(\ddot{\theta} + \frac{2}{t}\dot{\theta} \right) = \frac{1}{t^2} \frac{d}{dt} (t^2\dot{\theta}) = -q(t) \cos[\tilde{\mu}(t - t_i)] , \quad (4.42)$$

which can be integrated to yield,

$$\dot{\theta}(t) = \left(\frac{\tilde{\mu}^2\Phi_i}{\tilde{\Lambda}} \right) \frac{t_i}{t^2} \left[1 - e^{-\tilde{\Gamma}(t-t_i)/2} \left\{ \cos[\tilde{\mu}(t - t_i)] + \tilde{\mu}t \sin[\tilde{\mu}(t - t_i)] \right\} \right] + \dots \quad (4.43)$$

where the ellipses includes terms subleading in $\tilde{\Gamma}/\tilde{\mu}$, which we have dropped for consistency with our approximations above. Upon observation we find that there is no dependence on θ , specifically the $\sin(2\theta)$ term is absent. Thus, we can see that, in this limit the motion of $\dot{\theta}$ is driven solely by the oscillation of the inflaton and simply oscillates around zero, not maintaining any finite value. This is to be expected since this limit is equivalent to removing the B violating term, associated with the scalar baryon potential, from the equation.

Now consider the limit $p \gg q(t)$, for which the equation of motion becomes,

$$\ddot{\theta} + \frac{2}{t}\dot{\theta} + p \sin(2\theta) = 0 . \quad (4.44)$$

In this case, if we start from a state with finite energy, the friction term will damp the motion of the phase θ until it settles into one of its potential minima, and again there will be no non-zero $\dot{\theta}$ which persists and can lead to a non-zero baryon number density. Of course, this is to be expected since in this limit the \mathcal{C} and \mathcal{CP} breaking term has been removed.

Therefore, we can conclude that for successful Baryogenesis, we require $p \simeq q(t)$ so that both the B breaking and the \mathcal{C} and \mathcal{CP} terms can contribute to the time evolution of θ . In particular, we need the condition $p \simeq q(t)$ to be satisfied towards the end of the reheating epoch, since if it is satisfied too early, $q(t)$ will evolve to be smaller than p by the end of the epoch and any directed motion of θ will come to a stop. Thus, we require $p \simeq q(t_f)$, which we shall name the Sweet Spot Condition (SSC),

$$\tilde{\lambda}\phi_r^2 \simeq \frac{\tilde{\mu}^2\Phi_i}{\Lambda} \left(\frac{H_f}{H_i} \right) . \quad (4.45)$$

The SSC must be compatible with the prior assumptions we have made, such as that derived in Eq. (4.38). Upon combining these two conditions we find the following requirement, $\phi_r^2/\Lambda^2 \ll 1$. This can easily be satisfied, and has the flow on effect that we can drop the tilde notation from this point on, that is, $\tilde{\lambda} \approx \lambda$, $\tilde{\mu} \approx \mu$, and $\tilde{\Gamma} \approx \Gamma$.

4.3.1 Approximate Analytical Solution

Considering the SSC found in the previous section, the equation of motion is still difficult to solve, so we will first attempt to find an approximate analytical solution. We shall then compare this with numerical calculations to test for consistency.

The requirement for non-zero driven motion near the end of reheating is defined by the SSC, which can be written as follows,

$$\lambda\phi_r^2 \simeq \frac{H_f}{H_i} \frac{\mu^2\Phi_i}{\Lambda} \approx 7 \times 10^{14} \text{ GeV} \frac{T_{\text{rh}}^2}{\Lambda} , \quad (4.46)$$

where we have used Eq. (4.3), Eq. (4.12), Eq. (4.13), and Eq. (4.45). If we substitute the SSC into the equation of motion for θ we get,

$$\frac{9}{4}H^4\theta'' - \frac{3}{2\Lambda}H^2\Gamma\Phi' + \frac{\mu^2}{\Lambda}\frac{H}{H_i}\Phi_0\cos\left[\frac{2\mu}{3H}\right] - \lambda\phi_r^2\sin(2\theta) \approx 0, \quad (4.47)$$

where the primes denote derivatives with respect to the Hubble rate. We obtain the following simplified equation,

$$\frac{9}{4}H^4\theta'' \approx p\left(\sin(2\theta) - \cos\left[\frac{2\mu}{3H}\right]\right), \quad (4.48)$$

where we have fixed $p = \lambda\phi_r^2 \simeq \frac{H_f}{H_i}\frac{\Phi_0\mu^2}{\Lambda}$ near the end of reheating, the damping term has been assumed to be suppressed, and the reheating epoch has been taken to be sufficiently long such that $H_f \ll H_i$. In order to aid finding an approximate analytical solution we have neglected the Hubble damping term. In doing this we find the following simplified expression for the dynamics of θ near the end of reheating,

$$\ddot{\theta} \approx p(\sin(2\theta) - \cos(\mu t)). \quad (4.49)$$

If we now assume the two oscillatory parts have different frequencies, with the inflaton oscillation rate μ being slower, we find that the average value of $\dot{\theta}$ is of order,

$$|\dot{\theta}| \approx \frac{p}{\mu} = \frac{\lambda\phi_r^2}{\mu}. \quad (4.50)$$

Therefore, the baryon number density generated at the end of reheating is given by,

$$n_B \approx \frac{\lambda\phi_r^4}{\mu}. \quad (4.51)$$

Combining the entropy density at the end of reheating and the approximate analytical expression for n_B derived in Eq. (4.51), we obtain the following expression for the asymmetry parameter,

$$\eta_B^{reh} \approx 0.02\frac{\lambda\phi_r^4}{\mu T_{rh}^3} \approx 0.1\frac{\phi_r^2}{\Lambda T_{rh}}, \quad (4.52)$$

where we have used the SSC in Eq. (4.46). This results assumes certain constraints on the parameters, namely the SSC, $\frac{\phi_r^2}{\Lambda^2} \ll 1$, and that the energy density of the scalar baryon must be less than that of the radiation epoch at the end of reheating, which corresponds to,

$$\lambda\phi_r^4 < \frac{\pi^2}{30}g_*T_{rh}^4. \quad (4.53)$$

It is found that a wide range of parameter choices can be used that satisfy the observed baryon asymmetry. We also find that when undertaking numerical calculations of Eq. (4.39), they are in reasonable agreement with this solution, with results being consistent to within an order of magnitude for most parameter choices where driven motion is observed. It is generically found that high reheating temperatures are required, that is $T_{rh} > 10^{13}$ GeV.

4.3.2 Phase Locked States

A more rigorous solution for θ can be found by drawing an analogy between our mechanism and a forced pendulum. The equation of motion for θ can be parametrised as follows,

$$\ddot{\theta} + f(t)\dot{\theta} + p \sin(2\theta) = -q(t) \cos[\mu(t - t_i)] , \quad (4.54)$$

where

$$p = \lambda\phi_r^2, \quad f(t) = \frac{2}{t}, \quad q(t) = \frac{\mu^2 \Phi_i}{\Lambda} \left(\frac{t_i}{t}\right) e^{-\Gamma(t-t_i)/2} . \quad (4.55)$$

Now we can observe that the motion of θ is identical to that of a forced pendulum. The term proportional to $\sin(2\theta)$ can be viewed as the gravitational force on the pendulum, when it is at an angle 2θ from the vertical, q the external pushing force, and f the friction term. There is an added complexity in our case, in that the strength of the external force $q(t)$ and the friction $f(t)$ on the pendulum depend on t .

As discussed above, Baryogenesis is realised in this scenario when the solution of the equations of motion is found to give $\dot{\theta}(t_f) \neq 0$, at the end of reheating t_f . This means we must adjust the timing and intensity of the external pushing to match the motion of the pendulum, which is the idea embodied by the SSC, corresponding to $p \approx q(t_f)$. If this is satisfied, the rotational motion of the pendulum around the fixed point arises with an almost constant angular velocity $\dot{\theta}$. This is the solution we wish to determine.

The time evolution of $q(t)$ toward the end of the reheating epoch can be expected to be slow, so to analyse the dynamics of θ within that time frame, it is sufficient to replace it with the constant $q(t_f)$. The same can be said of the coefficient of the $\dot{\theta}$ term, which we replace with $(2/t_f)$. Thus, we obtain the following simplified equation,

$$\ddot{\theta} + \frac{2}{t_f} \dot{\theta} + p \sin(2\theta) = -q(t_f) \cos[\mu(t - t_i)] , \quad (4.56)$$

where the coefficients of $\dot{\theta}$ and $\cos[\mu(t-t_i)]$ are now constants, $p = \lambda\phi_r^2$ and $q(t_f) = e^{-\frac{1}{3}} \frac{H_{end}}{H_{ini}} \frac{\Phi_0 \mu^2}{\Lambda}$, the damping term has been taken to be suppressed, and $H_{end} \ll H_i$ has been assumed.

The relevant solutions to the above equation in our Baryogenesis scenario are those that increase or decrease monotonously in time with only small amplitude modulations. Such solutions exist and are known as phase-locked states, which are found in the study of the chaotic behaviour of the forced pendulum. The conditions for phase-locked states to exist were considered in the study of chaotic behaviour of electric current passing through a Josephson junction [504]. Since it is convenient to follow the notation adopted in these studies [505], we change the variables as follows,

$$\Theta \equiv 2\theta, \quad \tau \equiv \sqrt{2p} \left[(t - t_i) - \frac{\pi}{\mu} \right], \quad \omega \equiv \frac{\mu}{\sqrt{2p}}, \quad Q \equiv \sqrt{\frac{p}{2}} t_f, \quad \gamma \equiv \frac{q(t_f)}{p}. \quad (4.57)$$

Thus, the equation of motion becomes,

$$\ddot{\Theta} + \frac{1}{Q} \dot{\Theta} + \sin \Theta = \gamma \cos(\omega\tau). \quad (4.58)$$

Our equation coincides exactly with that of the forced pendulum or Josephson junctions. The generic phase-locked state solution to the above equation has the following form, when $\gamma \approx 1$,

$$\Theta(\tau) = \Theta_0 + n\omega\tau - \sum_{m=1}^{\infty} \alpha_m \sin(m\omega\tau - \phi_m), \quad (4.59)$$

where n and m are integers. In the numerical calculations we performed only the phase-locked states with $m = 1$ appear. In such solutions the period of the amplitude modulation is equal to that of inflaton's oscillation. Hence the solution to our equation of motion is of the form,

$$\Theta = \Theta_n + n(\omega\tau - \phi) - \alpha \sin(\omega\tau - \phi). \quad (4.60)$$

For these solutions, we can calculate the baryon number density n_B as the time average of $\dot{\Theta}$. From this we arrive at the following,

$$n_B = \phi_r^2 \langle \dot{\theta} \rangle = \sqrt{\frac{p}{2}} \phi_r^2 \langle \dot{\Theta} \rangle = \sqrt{\frac{p}{2}} \phi_r^2 n\omega = (\mu\phi_r^2) \frac{n}{2}. \quad (4.61)$$

Interestingly, this result depends on the integer n , where $n/2$ is the number of rotations of the phase θ per oscillation of the inflaton. The value of n depends on the validity of the phase-locked state and its stability. This is not a number which is predicted by the theory and hence we must determine it using numerical simulations. Although, the approximate solution derived in the previous section can provide an approximate value for n , which is found to be surprisingly consistent with numerical calculations.

4.4 The Generated Asymmetry Parameter

Combining the entropy density at the end of reheating and the analytical expression for n_B derived in Eq. (4.61), we obtain the following relation for the asymmetry parameter,

$$\eta_B^{\text{reh}} = \frac{n_B}{s} \approx 0.01n \times \left(\frac{\mu\phi_r^2}{T_{\text{rh}}^3} \right). \quad (4.62)$$

This relation needs to be consistent with the observational constraint presented in Eq. (4.14). At the end of reheating, there is no further generation of baryon number, but for reheating temperatures greater than ~ 100 GeV, sphaleron redistribution must be considered [214, 440, 506]. This redistribution leads to a dilution factor of $\frac{28}{79}$, and hence the required asymmetry parameter at the end of reheating is,

$$\eta_B^{\text{reh}} = \frac{79}{28}\eta_B \simeq 2.4 \times 10^{-10}. \quad (4.63)$$

The rough analytical expression that we derived, is found to be a good approximation for large values of n . By comparing these two solutions we can get an approximation of the parameter n . Combining Eq. (4.52) and Eq. (4.62) we obtain,

$$n \approx 2 \frac{\lambda\phi_r^2}{\mu^2}, \quad (4.64)$$

which must be greater than one if driven motion is to be observed. For values of n of $\mathcal{O}(10)$ and greater, the approximate solution is in good agreement with numerical simulations, which is likely due to the consistency of these n values with the assumptions used. In this mechanism, large values of λ are naturally required to produce driven motion, which we shall address in future work. We also require high reheating temperatures for the numerical simulations to be valid, although this can also be seen when substituting the SSC into Eq. (4.64),

$$n \approx 2 \times 10^{-12} \text{ GeV}^{-1} \frac{T_{\text{rh}}^2}{\Lambda}, \quad (4.65)$$

where a large reheating temperature is required for driven motion. It would generally be expected that the cutoff scale Λ would be of order of the Planck scale, such that the derivative coupling interaction term is valid near the beginning of reheating.

4.4.1 Conversion to Standard Model Particles

So far we have identified the baryon number density n_B with that of the scalar baryon, implying $n_B(\text{scalar}) = \phi_r^2 \dot{\theta}$. However, the actual baryon number density in our universe is made up of

fermionic matter. Therefore, the baryon number density of the scalar baryon must be converted to that of fermionic baryons. There are multiple ways to accomplish this conversion.

If we insist on using SM fields directly for the scalar-fermion conversion, we require $SU(3)_C \times SU(2)_L \times U(1)_Y \times B$ invariant interactions.

The simplest and most realistic model may be to reinterpret the baryon number associated with the scalar as lepton number, and introduce the lepton-number preserving dimension four interaction,

$$\Delta\mathcal{L}_{\text{int}} = y_L \phi^* \bar{\nu}_R^c \nu_R + \text{h.c.} \quad (4.66)$$

which describes the decay of the complex scalar lepton into a $\nu_R \nu_R$ pair. This same interaction can be used to generate a large Majorana mass for ν_R , when ϕ obtains the vacuum expectation value, $\langle \phi \rangle = \phi_r / \sqrt{2}$, with this leptonic scalar being a component of the neutrino mass generating model known as the seesaw mechanism [507–509]. The lepton number generated via the decay of the complex scalar can then be converted to baryon number through redistribution by $B - L$ conserving sphaleron processes [207, 217], as in the usual Leptogenesis scenarios [510].

Alternatively, one could introduce the following dimension six interaction between the scalar and fermionic baryon number currents,

$$\Delta\mathcal{L}_{\text{int}} = \frac{i}{\Lambda^2} (\phi^* \overleftrightarrow{\partial}_\mu \phi) \frac{1}{3} (\bar{u} \gamma^\mu u + \bar{d} \gamma^\mu d) , \quad (4.67)$$

where u and d are four component Dirac fields. Identifying the charges carried by the scalar and fermionic currents requires the existence of a term in the interaction Lagrangian which would lead to both the ϕ and the quark fields transforming at the same time, which we will not show explicitly here. Rewriting the scalar baryon in the polar coordinate parametrisation, $\phi = \frac{1}{\sqrt{2}} \phi_r e^{i\theta}$, and ignoring the spatial dependence of θ we obtain,

$$\Delta\mathcal{L}_{\text{int}} = -\frac{\phi_r^2}{\Lambda^2} \dot{\theta} \frac{1}{3} (u^\dagger u + d^\dagger d) , \quad (4.68)$$

such an interaction term has been utilised in previously considered Baryogenesis scenarios [503]. This term is analogous to a chemical potential coupling to the baryonic current, shifting in favour of matter or antimatter, where $\mu_B = -\frac{\phi_r^2}{\Lambda^2}$.

4.5 Conclusions and Future Prospects

In this chapter, we have considered a new Baryogenesis mechanism that acts during reheating, which takes its inspiration from the ratchet models of molecular motors in biological systems. The mechanism we propose here is found to produce driven motion in an analogous framework

to that found for a forced pendulum, in which the driving force is supplied by the oscillation of the inflaton, the position of the motor is embodied in the phase θ of a complex scalar baryon field, and the required breaking of the reflection symmetry is realised via the coupling of the inflaton to the scalar baryon. The push out-of-equilibrium comes from the reheating epoch itself. In our analysis, we find a rigorous solution which is dependent on an indeterminable parameter n ,

$$\eta_B^{\text{reh}} \approx 0.01n \times \left(\frac{\mu\phi_r^2}{T_{\text{rh}}^3} \right), \quad (4.69)$$

The issue with this solution is that the parameter n must be determined using numerical calculations. This can be bypassed by considering the approximate solution we derived,

$$\eta_B^{\text{reh}} \approx 0.02 \frac{\lambda\phi_r^4}{\mu T_{\text{rh}}^3} \approx 0.1 \frac{\phi_r^2}{\Lambda T_{\text{rh}}}, \quad (4.70)$$

which is in good agreement with numerical simulations for $n > 10$. Using these relations we found that it is possible to replicate the observed baryon asymmetry. We were also able to find an estimate of the parameter n through comparing the approximate and rigorous solutions, which allows a more targeted approach at testing the allowed parameter space with the numerical methods.

Although, in order for this to be achieved a very unnatural choice of the coupling λ must be chosen. This issue can be alleviated by a change in the scalar baryon potential we have considered, which shall be discussed in future work.

High reheating temperatures are a generic requirement of our model, which can be seen in the SSC, but is also a result of difficulties with the numerical calculations when considering reheating temperatures less than approximately 5×10^{13} GeV. Reheating temperatures greater than 10^{14} GeV could be a possible issue due to the over production of gravitinos, if one considers a supersymmetric theory [511–515], but we do not consider this an issue in our non-supersymmetric scenario.

We have made the simplifying assumption of a uniform isotropic universe and as such have ignored the spatial dependence of the scalar fields Φ and ϕ , and consequently that of the phase θ , in our analysis. In reality, as the universe expands, different parts of the universe will lose causal contact with each other, possibly leading to the evolutions of Φ and θ obtaining spatial dependencies. More work needs to be done to see how much of an impact, if any, this can have on the baryon number density generated.

More analysis needs to be done on the possible chaotic nature of this mechanism, given its parallels with the forced pendulum. Preliminary work into the Poincare maps associated with the nature of this scenario have been conducted, and will be discussed in future work. Also,

a further refinement of the numerical procedure needs to be considered to see whether lower reheating temperatures, corresponding to longer integration times, can be calculable.

Hopefully, through further investigation of this mechanism we can get a better understanding of the allowed regions of the parameter space. We also plan to explore the application of the ratchet mechanism in other cosmological settings, for example during the radiation epoch following the end of reheating.

Chapter 5

Gravitational Waves and the Cosmic Neutrino Background

The Cosmic Neutrino Background ($C\nu B$) contains information from very early times which may help illuminate both the properties of the neutrino sector and the evolution of the universe. Unfortunately, the weakly interacting nature of neutrinos combined with the low temperature of the background today, makes the prospect for detection near impossible in the foreseeable future. Despite this, in the early universe, the dynamics of the $C\nu B$ could have had significant effects on the evolution of the universe. The prospect of gleaning indirect evidence of the $C\nu B$ is to be explored in this chapter, through considering the possible implications for gravitational wave propagation. Given the dawn of the new era of gravitational wave astronomy, this is an exciting possibility.

We argue that a $C\nu B$ that carries a non-zero lepton charge develops gravitational instabilities, which are fundamentally related to the mixed gravity-lepton number anomaly. In the presence of this background, we find that a gravitational Chern-Simons (CS) term is induced, which leads to interesting physical effects. Firstly, gravitational waves propagating in such a neutrino background exhibit birefringent behaviour leading to an enhancement or suppression of the gravitational wave amplitudes, depending on the polarisation, with the magnitude of this effect related to the size of the lepton asymmetry. Secondly, this modification can lead to negative energy graviton modes in the high frequency regime, which induce very fast vacuum decay that produces, for example, positive energy photons and negative energy gravitons. Both of these effects can provide bounds on the lepton asymmetry of the universe, and hence probe the dynamics of the early universe [5].

5.1 The Cosmic Neutrino Background

Along with the CMB, the existence of the $C\nu B$ is an inescapable prediction of the standard hot big bang cosmology [516]. At early times in the universe, the neutrinos are in thermal equilibrium with the SM plasma through the weak interactions alone. As the cross sections for these interactions are small compared to electromagnetic processes, the neutrinos will fall out of equilibrium well before the first generation of charged SM species. As given in Table 1.3, this occurs just prior to BBN, at around 2-3 MeV - with there being a temperature range due to the different decoupling times for each neutrino species. This is due to the additional weak interactions between electron neutrinos and the electrons and positrons present in the primordial plasma, which are not present for the muon and tau neutrinos because at these temperatures the population of charged muons and taus is thermally suppressed. The $C\nu B$ is assumed to be a highly homogeneous and isotropic distribution of relic neutrinos with the characteristic temperature,

$$T_\nu = \left(\frac{4}{11}\right)^{\frac{1}{3}} T_\gamma \approx 1.945 \text{ K} , \quad (5.1)$$

where $T_\gamma = 2.725 \text{ K}$ is the temperature of the CMB today. Unlike the CMB though, the $C\nu B$ is extremely hard to detect and its properties are largely unknown. The reason for the difference between the temperatures of the $C\nu B$ and CMB, despite both sectors evolving as radiation for most of the cosmological history, is the extra entropy injection produced by the electron-positron annihilation dominantly into photons, which occurs after neutrino decoupling.

The physics of the generation and evolution of the $C\nu B$ has been a closely studied area due to the window it could provide to early universe physics [517–519]. One interesting characteristic of the $C\nu B$ is that it may exhibit a neutrino-antineutrino asymmetry. Unlike the baryon asymmetry which is strongly constrained, the $C\nu B$ asymmetry can be relatively large. The associated lepton asymmetry is defined as follows,

$$\eta_{\nu_\alpha} = \frac{n_{\nu_\alpha} - \bar{n}_{\nu_\alpha}}{n_\gamma} \simeq \frac{\pi^2}{12\zeta(3)} \left(\xi_\alpha + \frac{\xi_\alpha^3}{\pi^2} \right) , \quad (5.2)$$

for each neutrino flavour $\alpha = e, \mu, \tau$. Here $\xi_\alpha = \mu_\alpha/T$ is the degeneracy parameter, μ_α being the chemical potential for α -neutrinos. In fact, such an asymmetry is generically expected to be of the order of the observed baryon-antibaryon asymmetry, $\eta_B = (n_B - \bar{n}_B)/n_\gamma \sim 10^{-10}$, due to the equilibration by sphalerons of the lepton and baryon asymmetries in the very early universe, which conserve $B - L$. However, there are also models [520, 521] which predict an asymmetry in the neutrino sector that is many orders of magnitude larger than η_B . If so, this would have interesting cosmological implications for the QCD phase transition [522] and large-scale magnetic fields [523]. There have been attempts to circumvent the sphaleron redistribution constraints in such scenarios [520], which include the possible suppression of the equilibrium

sphaleron processes when the lepton asymmetry is very large, $\xi \gg 1$ [524]. The presence of a large asymmetry can alter the cosmological expansion history, as it changes the effective number of relativistic species N_{eff} , due to the increased energy density in the neutrino sector associated with a non-zero neutrino chemical potential. That is,

$$\Delta N_{\text{eff}} \simeq \frac{30}{7} \left(\frac{\xi_\nu}{\pi} \right)^2 + \frac{15}{7} \left(\frac{\xi_\nu}{\pi} \right)^4, \quad (5.3)$$

which can be constrained by measurements of the CMB.

The most stringent bound on the relic lepton asymmetry comes from the successful theory of BBN. The BBN observables primarily constrain the electron neutrino asymmetry, due to the implications of a large electron neutrino asymmetry on the helium abundance [516, 525–538]. However, this bound applies to all flavours, since neutrino oscillations below ~ 10 MeV are sizeable enough to lead to an approximate flavour equilibrium before BBN, $\mu_e \approx \mu_\mu \approx \mu_\tau (\equiv \mu_\nu)$ [539–541]. Although it has been found in a recent analysis that larger η_{ν_μ, ν_τ} asymmetries may be allowed [542]. In any case, the updated analysis in [536] leads to the following bound on the common degeneracy parameter,

$$|\xi_\nu| \lesssim 0.049. \quad (5.4)$$

The lack of strong constraints on the size of the relic lepton asymmetry has led to the postulation of many ideas associated with the prospect of having a large asymmetry, and mechanisms that can generate it, while not being in conflict with BBN constraints [542]. If it would become possible to observe the size of the asymmetry this would then be a potential smoking gun for these unorthodox particle physics models [450, 521, 543–555].

The experimental observation of neutrino oscillations demonstrates that the neutrinos must carry mass, but the mechanism for the origin of these masses is not explained within the SM, and is still unknown. An interesting possibility is that the SM neutrinos have Majorana mass terms,

$$\mathcal{L}_m = \frac{1}{2} m_\nu \bar{\nu}_L^c \nu_L, \quad (5.5)$$

where no right-handed neutrinos are present. In this scenario, the SM exhibits a mixed gravity-lepton number quantum anomaly.

If the neutrinos have Majorana masses, as we assume in this chapter, then the cosmological leptonic asymmetry carried by the CνB is defined as the difference between the left-handed neutrinos and right-handed antineutrinos [544, 556]. Due to this Majorana mass term, which violates lepton number, the relic neutrino asymmetry would be expected to be washed out once the mixing process becomes important. This would occur when the mixing timescale becomes smaller than that associated with the spatial expansion $\frac{1}{H}$. This means that at very early times a large Majorana neutrino asymmetry could be present and consistent with current observations,

assuming that the rate of lepton number violation is such that it is sufficiently washed out before the freeze-out of equilibrium sphaleron processes.

5.2 Gravitational Waves and the Graviton Propagator

The recent observation of gravitational waves from black hole merger events by the LIGO collaboration [314], signals the beginning of the new era of gravitational wave astronomy [315, 316]. An important feature of upcoming gravitational wave astronomy experiments will be the ability to differentiate polarisations, hence allowing the exploration of possible astrophysical and cosmological birefringent effects. These can be the results or signs of very interesting physics [557–560]; including CS modified gravity theories [561, 562]. Future gravitational wave detectors, such as eLISA [322, 323], will be able to probe the polarisations of incoming signals from astrophysical sources.

5.2.1 Linearised Gravitational Waves

In order to obtain predictions associated with gravitational wave production and propagation we utilise the linearised gravity approach. Gravitational waves can be described as small perturbations around a general spacetime background. The assumption of a small amplitude allows the higher order terms in h , the metric perturbations, to be neglected, removing non-linear interaction terms - producing the linearised gravity description. The metric perturbations are defined as follows,

$$g_{\mu\nu} \simeq \bar{g}_{\mu\nu} + \frac{1}{M_p} h_{\mu\nu} , \quad (5.6)$$

where $h_{\mu\nu}$ is the metric perturbation and satisfies $|h_{\mu\nu}| \ll M_p$, and $M_p = 1/\sqrt{8\pi G}$ is the reduced Planck mass. In the linearised approximation, the Einstein-Hilbert action becomes,

$$\begin{aligned} S_\Lambda &= \int dx^4 \sqrt{-g} \frac{M_p^2}{2} R \\ &= \int dx^4 \sqrt{-g} \left(\frac{1}{2} h_{\mu\nu} \square h^{\mu\nu} - h_{\mu\nu} \partial^\mu \partial_\alpha h^{\nu\alpha} + h \partial_\mu \partial_\nu h^{\mu\nu} - \frac{1}{2} h \square h \right) \end{aligned} \quad (5.7)$$

$$= \frac{1}{2} \int dx^4 \sqrt{-g} \bar{h}_{\mu\nu} \square \bar{h}^{\mu\nu} , \quad (5.8)$$

where in Eq. (5.8) we have taken $\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2} h \eta_{\mu\nu}$ and assumed the harmonic gauge, $\partial_\mu \bar{h}^\mu_\nu = \frac{1}{2} \partial_\nu \bar{h}$. The graviton coupling to matter is of the form $\mathcal{L}_{\text{int}} \propto h^{\mu\nu} T_{\mu\nu}$.

From this action we can derive the propagator for the graviton. In this chapter, we wish to consider the alteration to the graviton propagator in the presence of an asymmetric CνB. We shall do this by calculating the graviton polarisation tensor, which is connected to the graviton

action in the following way,

$$S_\Lambda = \frac{1}{2} \int dx^4 \sqrt{-g} \bar{h}^{\mu\nu} \Pi_{\mu\nu\rho\sigma} \bar{h}^{\rho\sigma} , \quad (5.9)$$

where $\Pi_{\mu\nu\rho\sigma}$ is the graviton polarisation tensor.

The graviton h can be split into circularly polarised planewaves - h_R and h_L - in an isotropic and homogeneous universe, similarly to the quantised $U(1)_X$ gauge boson in Chapter 3. This shall be useful for our analysis of birefringent effects in this chapter.

Gravitational waves can have many sources, with both present-day and primordial origins of interest. The recent measurement of black hole mergers is an example of present-day sources that correspond to very high energy scenarios. Cosmological sources may include phase transitions in the early universe, which can produce a stochastic gravitational wave background. Also, inflation generates tensor modes that can be imprinted on the CMB in the measured B-mode polarisation.

5.3 Graviton Polarisation Tensor in a Lepton Asymmetric CνB

In what follows we wish to consider the possible observational implications of a lepton asymmetric CνB on the properties of gravitational waves, and the possible inducement of gravitational instabilities. A non-zero lepton asymmetry for active neutrinos implies an imbalance between neutrinos of left-handed chirality and antineutrinos of right-handed chirality, and as we shall demonstrate, leads to the inducement of the gravitational CS term in the effective gravitational action. The possible effects of a large lepton asymmetry on primordial gravitational radiation has been considered in other contexts [563], and gravity has also been utilised to produce a relic neutrino asymmetry [564].

To determine the implications of a universal lepton asymmetry on gravity, we want to consider the effects on the graviton propagator. To do this we shall consider the presence of a chiral chemical potential μ_ν , which parametrises this asymmetry, and the gravity-lepton number chiral quantum anomaly. This anomaly is present in the SM when considering Majorana neutrinos; due to the absence of right-handed neutrinos. In this scenario we can expect an induced parity violating contribution to the gravitational action, which can be found through calculating the contribution of the chiral chemical potential to the graviton polarisation tensor.

We calculate the inducement of the CS like term to the effective graviton Lagrangian through the 1-loop graviton polarization diagram depicted in Figure 5.1, which is influenced by the chemical potential μ_ν . The lepton asymmetry is enforced in the Lagrangian through the chiral chemical potential by the following term, $\mathcal{L}_{\mu_\nu} = \bar{\nu} \not{b} \gamma^5 \nu = \mu_\nu \bar{\nu} \gamma_0 \gamma^5 \nu$, where we have considered

the frame in which the CνB is at rest ($\not{b} = \mu_\nu \gamma_0$). The neutrino propagator is altered as follows,

$$S(p) = \frac{i}{\not{p} - m - \not{b}\gamma^5} = \frac{i}{\not{p} - m} \sum_{n=0}^{\infty} \left(-i\not{b}\gamma^5 \frac{i}{\not{p} - m} \right)^n \equiv S_0(p) + \sum_{n=1}^{\infty} S_n(p) , \quad (5.10)$$

where $S_0(p)$ is the usual fermion propagator in vacuum.




FIGURE 5.1: The 1-loop correction to the graviton propagator that is relevant in the parity violating background.

In the calculation that follows, we shall use the standard Feynman rules [565, 566], with the above modified neutrino propagator to first-order in μ_ν , $S(p) \approx S_0(p) - i\mu_\nu \frac{i}{\not{p}-m} \gamma_0 \gamma^5 \frac{i}{\not{p}-m}$. The higher order terms in b_μ , or μ_ν , are neglected because we are only interested in the linear terms in b_μ , which give the CS like term. Thus we find that the induced parity odd part of the graviton polarization tensor is,

$$\Pi_{\mu\nu\rho\sigma} = - \int \frac{d^4 p}{(2\pi)^4} (2p+k)_\nu (2p+k)_\sigma \left[\text{Tr}(\gamma_\mu S_0(p+k) \gamma_\rho S_1(p)) \right. \quad (5.11)$$

$$\left. + \text{Tr}(\gamma_\rho S_0(p) \gamma_\mu S_1(p+k)) \right] . \quad (5.12)$$

To evaluate the divergent loop integral in Eq. (5.12) we employ the dimensional regularization method ($d = 4 - \epsilon$, $\epsilon \rightarrow 0$) and use the relations provided in Appendix C. This gives,

$$\begin{aligned} \Pi_{\mu\nu\rho\sigma} = & \frac{\mu_\nu}{8\pi^2} k^\alpha \varepsilon_{\mu\rho\alpha 0} \int_0^1 dx \left[\frac{4\pi^2 \lambda^2}{M^2} \right]^\epsilon \left[8x^2(1-x)^2(1-2x)^2 \frac{k^2}{M^2} \Gamma(1+\epsilon) k_\nu k_\sigma \right. \\ & + (24x^2 - 44x + 18) \Gamma(\epsilon-1) M^2 \eta_{\nu\sigma} - 16x^2(1-x)^2 \Gamma(\epsilon) k^2 \eta_{\nu\sigma} \\ & \left. - (80x^4 - 192x^3 + 156x^2 - 50x + 5) \Gamma(\epsilon) k_\nu k_\sigma \right] , \end{aligned} \quad (5.13)$$

where $M^2 = m^2 - x(1-x)k^2$ and the limit $\epsilon \rightarrow 0$ is assumed. In simplifying this result we find a divergent quantity that is of the following form,

$$\Pi_{\mu\nu\rho\sigma}^{(div)} = -\frac{1}{\epsilon} \frac{\mu_\nu}{2\pi^2} k^\alpha \varepsilon_{\mu\rho\alpha 0} m^2 \eta_{\nu\sigma} , \quad (5.14)$$

where γ is Euler's constant. A straightforward inspection reveals that this divergent term does not satisfy the gravitational Ward identity, $k^\nu \Pi_{\mu\nu\rho\sigma}^{(div)} \neq 0$, and hence violates the gauge invariance of the effective gravitational action. This has also been observed previously in related calculations [567, 568]. The origin of this violation is rooted in the method of dimensional regularization, which violates local Lorentz invariance explicitly through the extrapolation to a non-integer number of spacetime dimensions $d = 4 - \epsilon$. Therefore, following the standard lore, we introduce non-invariant counter-terms to renormalise away this unphysical divergent term. The polarisation tensor then takes the following simple form,

$$\Pi_{\mu\nu\rho\sigma} = \mu_\nu \varepsilon_{\mu\rho\alpha 0} k^\alpha [k_\nu k_\sigma - k^2 \eta_{\nu\sigma}] C(k^2), \quad (5.15)$$

where

$$C(k^2) = \frac{1}{192\pi^2} - \frac{m^2}{16\pi^2(k^2)^{3/2}} \left[\sqrt{k^2} - \sqrt{4m^2 - k^2} \tan^{-1} \left(\frac{\sqrt{k^2}}{\sqrt{4m^2 - k^2}} \right) \right]. \quad (5.16)$$

This further reduces to,

$$C(k^2) = \begin{cases} -\frac{1}{1920\pi^2} \frac{k^2}{m^2}, & \text{if } k^2/m^2 \ll 1 \\ \frac{1}{192\pi^2}, & \text{if } k^2/m^2 \gg 1 \end{cases}. \quad (5.17)$$

From this polarization tensor we can determine the induced parity violating term in the effective action. We shall now investigate the two possible limiting cases separately.

In the limit $k^2/m_\nu^2 \ll 1$, this term reads,

$$S_{eff} \propto \frac{\mu_\nu}{m_\nu^2} \int d^4x \varepsilon_{\mu\rho\alpha 0} h^{\mu\nu} \partial^\alpha \square (\square h^{\rho\sigma} \eta_{\nu\sigma} - \partial_\nu \partial_\sigma h^{\rho\sigma}). \quad (5.18)$$

In the case of the harmonic gauge, taking $\bar{h}^{\mu\nu} = h^{\mu\nu} - \frac{1}{2} \eta^{\mu\nu} h$, for which $\partial_\mu \bar{h}^{\mu\nu} = 0$, the action reduces to,

$$S \propto \frac{\mu_\nu}{m_\nu^2} \int d^4x \varepsilon_{\mu\rho\alpha 0} \bar{h}^{\mu\nu} \partial^\alpha \square^2 \bar{h}^{\rho\sigma} \eta_{\nu\sigma}. \quad (5.19)$$

Note that this induced term contains more than two derivatives and thus is significant only in the ultraviolet regime. Indeed, taking the harmonic gauge we derive the modified equation of motion for the linearised graviton field,

$$\square \bar{h}_{ij} = \frac{1}{1920\pi^2} \varepsilon_{ilk} \partial^l \frac{\mu_\nu}{m_\nu^2 M_p^2} \square^2 \bar{h}^k{}_j. \quad (5.20)$$

The dispersion relations for left (−) and right (+) graviton polarisation modes then readily follow from Eq. (5.20),

$$(\omega^2 - |\mathbf{k}|^2) \mp \frac{\mu_\nu}{1920\pi^2 m_\nu^2 M_p^2} |\mathbf{k}| (\omega^2 - |\mathbf{k}|^2)^2 = 0 . \quad (5.21)$$

One can then see that for large enough momenta $|\mathbf{k}|$ there are modes with imaginary frequencies,

$$(\omega^2 - |\mathbf{k}|^2) = -\frac{1920\pi^2 m_\nu^2 M_p^2}{\mu_\nu |\mathbf{k}|} , \quad (5.22)$$

which are unstable. Such potentially unstable modes, however, have extremely small wavelengths,

$$|\mathbf{k}| \gtrsim \frac{1920\pi^2 M_p^2}{\mu_\nu} . \quad (5.23)$$

In this trans-Planckian regime, the Einstein's theory itself is believed to be untrustworthy, so we do not consider this limit any further.

It is more interesting to consider the opposite limit, $k^2/m_\nu^2 \gg 1$. In this limit we obtain the following contribution to the graviton action,

$$\begin{aligned} S_{eff} &= -\frac{\mu_\nu}{192\pi^2} \int d^4x \varepsilon_{\mu\rho\alpha 0} h^{\mu\nu} \partial^\alpha (\square h^{\rho\sigma} \eta_{\nu\sigma} - \partial_\nu \partial_\sigma h^{\rho\sigma}) \\ &= \frac{\mu_\nu}{48\pi^2} \int d^4x K^0 , \end{aligned} \quad (5.24)$$

which contains the same number of derivatives as the standard kinetic term in the weak field approximation. In fact, K^0 is the linearised 0th component of the four dimensional CS topological current,

$$K^\beta = \varepsilon^{\beta\alpha\mu\nu} (\Gamma_{\alpha\rho}^\sigma \partial_\mu \Gamma_{\nu\sigma}^\rho - \frac{2}{3} \Gamma_{\alpha\rho}^\sigma \Gamma_{\mu\lambda}^\rho \Gamma_{\nu\sigma}^\lambda) . \quad (5.25)$$

Therefore, the presence of an asymmetry in the CνB replicates CS modified gravity,

$$S_{CS} = \int d^4x (\partial_\mu \theta) K^\mu = \int d^4x \theta (*RR) , \quad (5.26)$$

where we can make the identification $\partial_0 \theta = \frac{\mu_\nu}{48\pi^2}$ and,

$$*RR := R\tilde{R} = *R^\mu{}_\nu{}^{\rho\sigma} R^\nu{}_{\mu\rho\sigma} , \quad (5.27)$$

where the dual tensor is given by $*R^\mu{}_\nu{}^{\rho\sigma} := \frac{1}{2} \varepsilon^{\rho\sigma\alpha\beta} R^\nu{}_{\mu\alpha\beta}$.

This is the expected result in a parity violating background, and is found in both gravitational and electromagnetic cases [566–577].

5.4 Chern-Simons Modified Gravity and Observational Implications

CS modified gravity is an area of active research due to it naturally appearing in the context of anomaly cancellation in String Theory, via the Green-Schwarz mechanism [74], and in Loop Quantum Gravity, as well as providing many interesting phenomenological implications [561, 562]. It is also a useful model independent way of parametrising parity violating effects in the cosmological setting. This modification to the gravitational action is analogous to the CS terms associated with the Green-Schwarz mechanism utilised in Chapter 3, but with the Riemann tensor replacing the field strength tensor. There are many potential observational consequences of this extension that have been explored, from which limits have been imposed on the CS coupling parameter.

The terms added to the gravitational action for CS modified gravity are given below,

$$S_{CS} \propto \int d^4x \sqrt{-g} \theta^* RR, \quad (5.28)$$

where $\kappa = 1/8\pi G$, and θ denotes the CS coupling, which we have assumed to be a non-dynamical CS extension. In the case of a dynamical CS extension, a kinetic and potential term for θ are required,

$$S_\theta \propto \int d^4x \sqrt{-g} [g^{\mu\nu} (\nabla_\mu \theta) (\nabla_\nu \theta) + 2V(\theta)]. \quad (5.29)$$

Although, experimental tests of CS modified gravity have been mostly confined to the non-dynamical scenario [578–581]. This is due to the increased complexity associated with the dynamical case, which has not been explored as extensively in the literature. Fortunately, the case we consider in our work is of the non-dynamical type, and hence we can consider the effects that have been explored in this simpler scenario.

An interesting phenomenon associated with CS gravity is birefringent propagation of gravitational waves. The CS correction leads to an exponential enhancement and suppression of the left and right circularly polarised waves, which depend on the wave number, CS coupling and the integrated history of the propagation [562].

5.4.1 Birefringent Propagation through an Asymmetric CνB

Planned gravitational wave detectors, such as eLISA, DECIGO and BBO, can potentially measure the polarization of observed gravitational waves, and hence potentially this birefringence effect. We now wish to consider the magnitude of the birefringent effect induced by the CS

term in Eq. (5.24), and determine the possible sources that could be constrained by this effect. To this end, we parametrise the gravitational waves as, $h_{ij} = \frac{A_{ij}}{a(\eta)} \exp[-i(\phi(\eta) - \mathbf{k} \cdot \mathbf{x})]$, which can be decomposed into the two circularly polarised states - e_{ij}^R and e_{ij}^L . The two possible circularly polarised states are, $e_{ij}^R = \frac{1}{\sqrt{2}}(e_{ij}^+ + ie_{ij}^\times)$ and $e_{ij}^L = \frac{1}{\sqrt{2}}(e_{ij}^+ - ie_{ij}^\times)$, which satisfy $n_i \varepsilon^{ijk} e_{kl}^{R,L} = i\lambda_{R,L}(e_l^j)^{R,L}$, where $\lambda_{R,L} = \pm 1$. The phase factor $\lambda_{R,L}$ leads to exponential suppression or enhancement of the left and right circular polarisations of the the propagating gravitational waves, the magnitude of which we shall now calculate.

From the equations of motion for the action $S = S_{EH} + S_{eff}$ we obtain, for a general θ [582],

$$\begin{aligned} (i\phi_{,\eta\eta}^{R,L} + (\phi_{,\eta}^{R,L})^2 + \mathcal{H}_{,\eta} + \mathcal{H}^2 - |\mathbf{k}|^2) \left(1 - \frac{\lambda_{R,L} \kappa \theta_{,\eta}}{a^2}\right) \\ = \frac{i\lambda_{R,L} |\mathbf{k}|}{a^2} (\theta_{,\eta\eta} - 2\mathcal{H}\theta_{,\eta})(\phi_{,\eta}^{R,L} - i\mathcal{H}). \end{aligned} \quad (5.30)$$

We first solve the above equation assuming propagation in the matter dominated epoch, $a(\eta) = a_0 \eta^2 = \frac{a_0}{1+z}$, as we are considering sources within the range of possible near future observatories - $z < 30$. The accumulated phase over propagation, to first order in θ , is given by,

$$\Delta\phi_{mat}^{R,L} = i\lambda_{R,L} |\mathbf{k}| H_0 \int_{\eta}^1 \left[\frac{1}{4} \theta_{,\eta\eta} - \frac{1}{\eta} \theta_{,\eta} \right] \frac{d\eta}{\eta^4}. \quad (5.31)$$

In the case considered in this work, we make the following identification $\theta_{,\eta} = \left(\frac{a(\eta_0)}{a(\eta)}\right)^2 \frac{\mu_0}{48\pi^2 M_p^2}$, where $\mu_0 = a(\eta)\mu_\nu$ is the chemical potential at present. Thus, for the asymmetric CνB,

$$\Delta\phi_{mat}^{R,L} = -i \frac{1}{288\pi^2} \frac{\mu_\nu H_0}{M_p^2} \left(\frac{|\mathbf{k}|}{1 \text{ GeV}} \right) (1+z)^4. \quad (5.32)$$

Hence the ratio of the wave amplitudes of the two polarisation states is given by,

$$\frac{h_R}{h_L} \propto e^{-2|\Delta\phi_{mat}^{R,L}|}. \quad (5.33)$$

Taking into account the current bounds on the CνB asymmetry parameter, $\xi < 0.049$, we find $|i\Delta\phi_{mat}^{R,L}| \lesssim 10^{-87} \left(\frac{|\mathbf{k}|}{1 \text{ GeV}}\right)$, for $z \sim 30$. This accumulated phase difference is too small to be observable by any conceivable gravitational wave detector.

This leads us to shift to the more interesting scenario, which is the propagation of gravitational waves from sources in the very early universe. The effect would be thought to be of a higher magnitude because both the chemical potential would have been larger and the waves would have a longer propagation time over which a phase difference can be accumulated. Conceivably, any early universe sources could provide constraints, if the two polarisations of the signal are differentiable. Some examples of this would be the inflationary gravitational waves, and those

produced by collisions of bubbles formed during phase transitions. Therefore, we now consider gravitational waves produced at very early times, during the radiation dominated epoch. The accumulated phase now reads,

$$\Delta\phi_{rad}^{R,L} = i\lambda_{R,L} \frac{|\mathbf{k}|}{\Omega_{r,0} H_0^2} \int_{\eta}^1 \left[\frac{1}{2} \theta_{,\eta\eta} - \frac{1}{\eta} \theta_{,\eta} \right] \frac{d\eta}{\eta^2}, \quad (5.34)$$

where $\Omega_{r,0} \sim 9.2 \cdot 10^{-5}$ is the radiation density parameter today. After taking the integral we find,

$$\Delta\phi_{rad}^{R,L} \simeq -i\lambda_{R,L} \xi_{\nu} \left(\frac{|\mathbf{k}|}{1 \text{ GeV}} \right) \left(\frac{T_s}{1 \text{ TeV}} \right)^4, \quad (5.35)$$

where we have redefined the redshift in terms of the temperature at which the gravitational waves are produced, T_s , or when the asymmetry is generated, whichever is lowest.

From Eq. (5.35), it can be seen that if an asymmetry is present in the $C\nu B$, which equilibrium sphalerons transitions may assure, then it is possible to get significant birefringent behaviour in the propagation of gravitational waves from primordial sources, depending on the momenta $|\mathbf{k}|$ of the gravitational waves and size of the asymmetry.

If one is to assume that the characteristic momenta of the gravitational waves is of the order of the Hubble rate at the time of production ($|\mathbf{k}| \sim H$), and the asymmetry is already present, then we get the following interesting constraint on the gravitational waves produced from the source, at a temperature T_s ,

$$\Delta\phi_{rad}^{R,L} \simeq -i\lambda_{R,L} \xi_{\nu} \left(\frac{T_s}{10^6 \text{ GeV}} \right)^5, \quad (5.36)$$

hence very early sources and large asymmetry are required, if a relative enhancement or suppression of order one is to be observed.

5.5 Induced Ghost-like Modes and Vacuum Decay

Another interesting consequence of the induced CS term in Eq. (5.24) is that short-scale gravitational fluctuations exhibit negative energy modes, which if present lead to a rapid decay of a vacuum state, for example, into negative energy graviton and photons [581]. Since in this setting the graviton energy would not be bounded from below, the phase space for this process is formally infinite [583, 584], and as such will develop very rapidly. We investigate the production of two photons and a negative energy graviton via this process, to provide constraints on the

neutrino asymmetry at early times. The relevant effective interaction is of the form,

$$\begin{aligned} S_{\text{int}} &\sim \frac{1}{m_*} \int d^4x h_{\mu\nu}^{\text{can}} T^{\mu\nu} \\ &= \frac{1}{m_*} \int d^4x \frac{1}{2} h^{\text{can}} F_{\mu\nu} F^{\mu\nu} - h_{\mu\nu}^{\text{can}} F^{\mu\alpha} F_{\alpha}^{\nu}, \end{aligned} \quad (5.37)$$

where the canonically normalised graviton field is $h_{\mu\nu}^{\text{can}} = m_{\text{can}} h_{\mu\nu}^{\text{can}}$, with the definition,

$$m_{\text{can}} = M_p \sqrt{1 + \lambda_{R,L} \frac{|\mathbf{k}|}{am_{CS}}}, \quad (5.38)$$

where m_{CS} is the analogous CS mass scale,

$$m_{CS}(t) = \frac{M_p^2}{\mu_\nu} = \frac{M_p^2}{\xi T} = \frac{a(t) M_p^2}{\mu_0}. \quad (5.39)$$

5.5.1 Photon Energy Spectrum from Induced Vacuum Decay

To obtain a finite result for the decay rate we need to constrain the phase space. In the absence of a fundamental physical reason for such a truncation, we follow [583, 584], and simply cut-off the three momenta at $|\mathbf{k}|_{\text{max}} = \Lambda$. In the analysis that follows, we consider decays into this mode as it will have the largest contribution to the energy density of the generated photons. In addition, we take the reasonable approximation,

$$m_{\text{can}} \simeq \sqrt{\frac{|\mathbf{k}| \mu_\nu}{a}}, \quad (5.40)$$

and consider the dynamics of our scenario prior to BBN and after reheating, when the universe is radiation dominated and evolves as follows,

$$a(t) = a_0 \sqrt{t} = \sqrt{2\Omega_{r,0}^{1/2} H_0 t}, \quad (5.41)$$

where $\Omega_{r,0} \sim 9.2 \cdot 10^{-5}$ is the radiation density parameter today.

The time at which this ghost term is no longer present will be defined as t_* and is found in terms of the scale factor as,

$$1 \simeq \frac{\Lambda}{a(t_*) m_{CS}(t_*)} \Rightarrow a(t_*) \simeq \sqrt{\frac{\mu_0 \Lambda}{M_p^2}} \quad \text{or} \quad a(t_*) \simeq \frac{\xi_\nu T_* \Lambda}{M_p^2}, \quad (5.42)$$

where T_* is the temperature at which the ghost terms stop contributing.

This fixes the time at which the ghost modes no longer exist, and decay of the vacuum ceases. We can reinterpret this as a temperature, so that it is possible to associate this with the maximal

reheating temperature, and also ensure it does not have adverse implications on BBN. If we assume that the asymmetry is produced during or after the reheating epoch, and prior to BBN, the scale factor has a $\frac{1}{T}$ dependence, if we ignore the decoupling of radiation degrees of freedom. The scale factor takes the following form,

$$a(t) \simeq \left(\frac{90\Omega_{r,0}}{g^*\pi^2} \right)^{\frac{1}{4}} \frac{\sqrt{H_0 M_p}}{T}, \quad (5.43)$$

where $g^* \simeq 106.75$. Equating Eq. (5.42) and (5.43) to find the temperature at which this effect ends, we find,

$$\begin{aligned} T_* &= \left(\sqrt{\frac{90\Omega_{r,0}}{g^*\pi^2}} \frac{H_0 M_p^3}{\xi_\nu^2} \right)^{\frac{1}{4}} \sqrt{\frac{M_p}{\Lambda}} \\ &\simeq \frac{440}{\sqrt{\xi_\nu}} \text{ GeV} \sqrt{\frac{M_p}{\Lambda}}. \end{aligned} \quad (5.44)$$

Given that the maximum reheating temperature is $T_{\text{rh}} \sim 10^{15}$ GeV, Eq. (5.44) implies we can constrain the production temperature of neutrino asymmetries satisfying $\xi_\nu \gtrsim 2 \cdot 10^{-25} \frac{M_p}{\Lambda}$, with smaller ξ 's not generating ghost like modes after reheating. We also assume here that ξ is approximately constant, and hence is the same parameter currently constrained by BBN measurements, in the calculation of the lepton asymmetry stored in the $C\nu B$.

Next we compute the spectrum of photons generated by the induced vacuum decay, and then subsequently the energy density, which can be constrained by experiment. It is given by,

$$\frac{1}{a^3} \frac{d}{dt} (a^3 n(k, t)) = \Gamma \delta \left(\frac{|\mathbf{k}|}{\Lambda} - 1 \right), \quad (5.45)$$

where $n(k, t)$ is the number of photons per unit logarithmic wave number $|\mathbf{k}|$ and Γ is the total decay width, which we take to approximately be,

$$\Gamma \sim \frac{\Lambda^6}{m_{can}^2} = \frac{a(t)\Lambda^6}{|\mathbf{k}|\mu_\nu} = \frac{a(t)^2\Lambda^5}{\mu_0}. \quad (5.46)$$

Since the above decay rate is much faster than the expansion rate of the universe, we may safely assume that the decay is approximately instantaneous. Therefore, we fix the scale factor in Eq. (5.46) at time t_a , when the asymmetric background is first produced. We then integrate Eq. (5.45) between the time t_a and when the ghost terms are no longer present, t_* ,

$$|\mathbf{k}| n_*(|\mathbf{k}|) \sim \frac{a(t_*)^2 \Lambda \Gamma_a}{5\Omega_{r,0}^{1/2} H_0}. \quad (5.47)$$

Taking into account the dilution factor due to the expansion of the universe since the end of photon production to today, $\left(\frac{a(t_*)}{a_0}\right)^3 = a(t_*)^3$, we obtain,

$$|\mathbf{k}|n_0(|\mathbf{k}|) \sim \frac{a(t_*)^5 \Lambda \Gamma_a}{5\Omega_{r,0}^{1/2} H_0}. \quad (5.48)$$

Therefore, the energy density for a given momenta $|\mathbf{k}|$ is,

$$\frac{dE}{d^3x d \ln |\mathbf{k}|} \sim |\mathbf{k}|n_0(|\mathbf{k}|) \sim \frac{\xi^4 T_*^5}{10T_a^2} \sqrt{\frac{M_p^3}{H_0}} \left(\frac{\Lambda}{M_p}\right)^{11}. \quad (5.49)$$

We can now obtain a bound on the energy density in the produced photons, through the observation that the universe is not radiation dominated today,

$$\frac{dE}{d^3x d \ln |\mathbf{k}|} \lesssim M_p^2 H_0^2. \quad (5.50)$$

This means we get the following constraint on ξ_ν , assuming the asymmetry is generated above the characteristic temperature T_* , when requiring consistency with observation,

$$\xi_\nu \lesssim 2 \cdot 10^{-41} \left(\frac{T_a}{10^{15} \text{ GeV}}\right)^{4/3} \left(\frac{M_p}{\Lambda}\right)^{17/3}, \quad (5.51)$$

for which it is assumed $T_a \gtrsim \frac{440}{\sqrt{\xi_\nu}} \text{ GeV} \sqrt{\frac{M_p}{\Lambda}}$. Equivalently,

$$T_* \gtrsim 10^{23} \text{ GeV} \left(\frac{T_a}{10^{15} \text{ GeV}}\right)^{-2/3} \left(\frac{\Lambda}{M_p}\right)^{17/6}. \quad (5.52)$$

Thus we arrive at the conclusion that, unless $\Lambda \ll M_p$, the resulting photon energy density from the induced vacuum decay can hardly be accommodated with observation. Substituting the constraint in Eq. (5.51) into that for the asymmetry stored in the CνB as a function of ξ_ν , in Eq. (5.2), we find the following bound,

$$\eta_\nu \lesssim 10^{-41} \left(\frac{T_a}{10^{15} \text{ GeV}}\right)^{4/3} \left(\frac{M_p}{\Lambda}\right)^{17/3}. \quad (5.53)$$

If we instead assume that $T_a \lesssim \frac{440}{\sqrt{\xi_\nu}} \text{ GeV} \sqrt{\frac{M_p}{\Lambda}}$, and hence vacuum decay does not occur, then we get the following constraint on η_ν ,

$$\eta_\nu \lesssim 0.033 \left(\frac{2000 \text{ GeV}}{T_a}\right)^2 \frac{M_p}{\Lambda}, \quad (5.54)$$

where $\eta_\nu \lesssim 0.033$ is the current upper limit from BBN constraints.

5.6 Conclusions and Future Prospects

In this chapter we have argued that a relic neutrino background with non-zero lepton number can lead to gravitationally observable effects [5]. We have explicitly calculated the parity odd part of the graviton polarization tensor in the lepton asymmetric $C\nu B$, which replicates the gravitational CS term in the effective action. The observable implications of this were then explored, wherein the derived gravitational instabilities are related to the gravity-lepton number mixed anomaly.

The induced CS term leads to birefringent behaviour, causing an enhancement or suppression of the gravitational wave amplitudes depending on the polarisation. While this effect is negligible for local sources, we demonstrate that it could be sizeable for gravitational waves produced in the very early universe,

$$\Delta\phi_{rad}^{R,L} \simeq -i\lambda_{R,L}\xi_\nu \left(\frac{|\mathbf{k}|}{1 \text{ GeV}}\right) \left(\frac{T_s}{1 \text{ TeV}}\right)^4, \quad (5.55)$$

which when considering a source produced with momenta $|\mathbf{k}| \sim H$, with H being the Hubble rate at the time of production,

$$\Delta\phi_{rad}^{R,L} \simeq -i\lambda_{R,L}\xi_\nu \left(\frac{T_s}{10^6 \text{ GeV}}\right)^5, \quad (5.56)$$

which immediately indicates the need for very early sources, if interesting constraints are to be obtained.

In addition to the above, we have also argued that short-scale gravitational fluctuations in the presence of asymmetric $C\nu B$ exhibit negative energy modes, which can lead to the rapid decay of the vacuum state into negative energy graviton and positive energy photons. Since the graviton energy is not bounded from below, the phase space for this process is formally infinite, that is the instability is expected to develop very rapidly. Conservatively, we introduced a comoving cut-off Λ and computed the spectrum of produced photons as a function of the neutrino chemical potential. From the constraints on the radiation energy density today, we have obtained an interesting bound on the neutrino degeneracy parameter,

$$\xi_\nu \lesssim 2 \cdot 10^{-41} \left(\frac{T_a}{10^{15} \text{ GeV}}\right)^{4/3} \left(\frac{M_p}{\Lambda}\right)^{17/3}, \quad (5.57)$$

which unless $\Lambda \ll M_p$, would effectively rule out the existence of an asymmetric $C\nu B$ that is produced early enough for ghost modes to be present. If we assume there are no ghost modes

associated with the $C\nu B$ at any point in the early universe we get the following constraint,

$$\eta_\nu \lesssim 0.033 \left(\frac{2000 \text{ GeV}}{T_a} \right)^2 \frac{M_p}{\Lambda} . \quad (5.58)$$

We believe that the findings reported in this chapter will prove to be useful for gaining a greater understanding of the properties of the $C\nu B$, and possibly allow constraints to be placed on particle physics models containing a lepton asymmetry. Being able to constrain or obtain indirect measurements of the size of the lepton asymmetry would help illuminate the properties of the neutrinos and possibly the mechanism for Baryogenesis.

This paper [5] explores the exciting new possibility of using gravitational wave phenomena to uncover information about the properties of the fundamental particles of nature. It will be interesting to consider these $C\nu B$ effects further, and also if similar phenomena could be induced by dark matter or other potential relics.

Chapter 6

Concluding Remarks and Outlook

The Standard Models of Particle Physics and Cosmology have been highly successful at describing and reproducing the observed dynamics and properties of the Universe, but they are incomplete. There are still many mysteries of nature yet to be solved, for which new physics beyond the standard paradigms is required. In this thesis we have tried to propose solutions for some of these problems by considering particle physics ideas, specifically in relation to quantum anomalies, to early universe cosmology. This was motivated by the knowledge of the strong intertwining of cosmological and particle dynamics at very early times, when the microscopic dynamics of the fundamental particles directly dictated the evolution of the universe. Through considering a combination of observables from terrestrial collider searches and cosmological observables it may be possible to piece together the answers to many of the open questions of our universe.

In Chapter 2, we proposed a new class of natural inflation models which provided a solution to the hierarchy problem through a hidden scale invariance realised through the introduction of a dilaton field [1]. Given the scale invariant symmetry of the theory, the inflationary potential naturally contains a flat direction in the classical limit, which is lifted by quantum corrections. Thus inflation can naturally, without fine-tuning, proceed when the inflaton field evolves along this direction. We find that in the conformal limit, the inflaton potential is linear, which gives predictions in agreement with observations. Therefore, this model provides a successful inflationary scenario within which a solution to the hierarchy problem of the Standard Model can be found.

Chapter 3 presented an unorthodox mechanism for the origin of the matter-antimatter asymmetry as well as dark matter; one that acts during the inflationary epoch [2, 3]. This mechanism forogenesis involved the introduction of an anomalous gauge interaction and sterile fermion to the Standard Model. The anomalies associated with the new gauge field provided the X charge violation, and the corresponding counter terms violated \mathcal{C} and \mathcal{CP} in the cosmological setting,

while the inflationary epoch provided the push out-of-equilibrium. It was found that this scenario for cogenesis can successfully reproduce the observed values of the baryon asymmetry and dark matter abundance for the two possible cases considered - gauged B and gauged $B - L$ charge - for certain parameter spaces. The general mechanism for cogenesis developed here could be applied to more complex models involving other or extra anomalous gauge symmetries and additional sterile or non-sterile fermionic states. It is possible that these additions could lead to a lessening of the parameter constraints imposed by the observed matter-antimatter asymmetry, through extra contributions to the luminous matter generation. Given that this model involves the introduction of new gauge bosons and a dark sector to the Standard Model, this mechanism could have potential avenues for experimental investigation at terrestrial collider searches and direct detection experiments. Therefore, it may be possible to utilise both terrestrial and cosmological measurements to constrain this mechanism.

In Chapter 4, we presented a model for Baryogenesis during reheating that utilises the Ratchet Mechanism [4]. We introduced a theory containing two fundamental scalars, an inflaton consistent with the Starobinsky inflationary mechanism, and a complex scalar baryon with a symmetric potential; with the two scalars interacting via a derivative coupling. The scalar baryon potential violates B , and the violation of C and \mathcal{CP} is introduced by the derivative coupling interaction. The push out-of-equilibrium in this mechanism is provided by the reheating epoch, which is caused by the coherent oscillation of the inflaton in its potential. In order for a non-zero baryon number density to be produced, driven motion must be induced in the phase of the complex scalar baryon. The inflaton-scalar baryon system was found to act analogously to a forced pendulum, with driven motion achieved near the end of reheating for parameters consistent with the Sweet Spot Condition. This result implied a high reheating temperature as a generic requirement of our model. Further analysis of this mechanism could provide interesting cosmological phenomenology beyond the reheating epoch through the decays and interactions of the baryonic scalar.

Chapter 5 discussed a novel way to utilise gravitational waves to illuminate the properties of the illusive Cosmic Neutrino Background [5]. We explicitly calculated the parity odd part of the graviton polarization tensor in the presence of a lepton asymmetric Cosmic Neutrino Background, which replicates the gravitational Chern-Simons term in the effective action, in the vanishing neutrino mass limit. The induced Chern-Simons term causes birefringent behaviour in gravitational wave propagation leading to an enhancement or suppression of the gravitational wave amplitude depending on the polarisation. While this effect is negligibly small for local sources, we demonstrated that it could be sizeable for gravitational waves produced in the very early universe with a momenta $|\mathbf{k}| \sim H$. We also argued that a relic neutrino background with non-zero lepton number exhibits gravitational instabilities that are related to the gravity-lepton number mixed anomaly. The induced negative energy modes, lead to a rapid decay of a vacuum state into photons and gravitons, from which we could derive observational bounds. From the

constraints on the radiation energy density today, we were able to obtain an interesting bound on the neutrino degeneracy parameter. We believe that the findings reported in this Chapter will prove to be useful for further understanding of the properties of the Cosmic Neutrino Background and allow constraints to be placed on particle physics models that generate a lepton asymmetry. Being able to constrain or obtain indirect measurements of the size of the lepton asymmetry would help illuminate the properties of the neutrinos and possibly the mechanism for Baryogenesis. This work explored the exciting new possibility of using gravitational wave phenomena to uncover information about the properties of the fundamental particles of nature.

In this thesis, we sought to demonstrate the importance of particle physics in the evolution of the early universe, and some of the interesting ways in which this could be explored; predominantly through the consideration of quantum anomalies. Increased investigation into such phenomena, as well as other particle physics applications to cosmology, will undoubtedly further increase our understanding of nature. It is clear that this approach, in concert with terrestrial particle physics phenomenology and experimental searches, is the clear way forward in our endeavour to understand the fundamental nature of the world around us.

Appendix A

Baryon and Lepton Number Anomalies in the Standard Model

A.1 Baryon Number Anomalies

The introduction of a gauged baryon number leads to the inclusion of quantum anomalies in the theory, refer to Figure 1.2. The anomalies, for the baryonic current, are given by the following,

For $SU(3)^2U(1)_B$,

$$\mathcal{A}_1(SU(3)^2U(1)_B) = Tr[\lambda^a \lambda^b B] = 3 \times \frac{3}{2} \left(\sum_{left} B_i - \sum_{right} B_i \right) = 0. \quad (\text{A.1})$$

For $SU(2)^2U(1)_B$,

$$\mathcal{A}_2(SU(2)^2U(1)_B) = Tr[\tau^a \tau^b B] = \frac{3 \times 3}{2} B_Q = \frac{3}{2}. \quad (\text{A.2})$$

For $U(1)_Y^2U(1)_B$,

$$\mathcal{A}_3(U(1)_Y^2U(1)_B) = Tr[YYB] = 3 \times 3(2Y_Q^2 B_Q - Y_u^2 B_u - Y_d^2 B_d) = -\frac{3}{2}. \quad (\text{A.3})$$

For $U(1)_B^2U(1)_Y$,

$$\mathcal{A}_4(U(1)_B^2U(1)_Y) = Tr[BBY] = 3 \times 3(2B_Q^2 Y_Q - B_u^2 Y_u - B_d^2 Y_d) = 0. \quad (\text{A.4})$$

For $U(1)_B^3$,

$$\mathcal{A}_5(U(1)_B^3) = Tr[BBB] = 3 \times 3(2B_Q^3 - B_u^3 - B_d^3) = 0 . \quad (\text{A.5})$$

For $U(1)_B$,

$$\mathcal{A}_6(U(1)_B) = Tr[B] = 3 \times 3(2B_Q - B_u - B_d) = 0 , \quad (\text{A.6})$$

where the factor of 3×3 is a result of there being three generations of quarks and three colours for each quark. The δ^{ab} terms are not included in the anomalies.

A.2 Lepton Number Anomalies

When including right handed neutrinos into the SM the quantum anomalies for a gauged lepton number, or leptonic current, are the following,

For $SU(3)^2U(1)_L$,

$$\mathcal{A}_1(SU(3)^2U(1)_L) = Tr[\lambda^a \lambda^b L] = \frac{3}{2} \left(\sum_{left} L_i - \sum_{right} L_i \right) = 0 . \quad (\text{A.7})$$

For $SU(2)^2U(1)_L$,

$$\mathcal{A}_2(SU(2)^2U(1)_L) = Tr[\tau^a \tau^b L] = \frac{3}{2} L_L = \frac{3}{2} . \quad (\text{A.8})$$

For $U(1)_Y^2U(1)_L$,

$$\mathcal{A}_3(U(1)_Y^2U(1)_L) = Tr[YYL] = 3(2Y_L^2 L_L - Y_e^2 L_e - Y_\nu^2 L_\nu) = -\frac{3}{2} . \quad (\text{A.9})$$

For $U(1)_L^2U(1)_Y$,

$$\mathcal{A}_4(U(1)_L^2U(1)_Y) = Tr[LLY] = 3(2L_L^2 Y_L - L_e^2 Y_e - L_\nu^2 Y_\nu) = 0 . \quad (\text{A.10})$$

For $U(1)_L^3$,

$$\mathcal{A}_5(U(1)_L^3) = Tr[LLL] = 3(2L_L^3 - L_e^3 - L_\nu^3) = 0 . \quad (\text{A.11})$$

For $U(1)_L$,

$$\mathcal{A}_6(U(1)_L) = Tr[L] = 3(2L_L - L_e - L_\nu) = 0 . \quad (\text{A.12})$$

If the right handed neutrinos are not included in the SM, \mathcal{A}_5 and \mathcal{A}_6 will be non-zero. That is, $\mathcal{A}_5 = 3$ and $\mathcal{A}_6 = 3$, where \mathcal{A}_6 is to the graviton-lepton anomaly.

A.3 Mixed Gauged Baryon and Lepton Number Anomalies

If these two gauge groups are introduced then the interactions between the leptonic and baryonic currents must also be anomaly free,

For $U(1)_B^2 U(1)_L$,

$$\mathcal{A}(U(1)_B^2 U(1)_L) = \text{Tr}[BBL] = 0 . \quad (\text{A.13})$$

For $U(1)_B U(1)_L^2$,

$$\mathcal{A}(U(1)_B U(1)_L^2) = \text{Tr}[LLB] = 0 . \quad (\text{A.14})$$

For $U(1)_B U(1)_L U(1)_Y$,

$$\mathcal{A}(U(1)_B U(1)_L U(1)_Y) = \text{Tr}[BLY] = 0 . \quad (\text{A.15})$$

These will only be non-zero if fermions such as leptoquarks are added to the SM. There are no fermions in the SM which can couple to both a leptophobic gauge boson and a leptophilic gauge boson.

Some recent models have introduced leptoquarks along with gauged baryon and lepton number symmetries into the SM [76, 77]. To ensure that these mixed interactions don't lead to new gauge anomalies, the number of types of leptoquarks and the quantum numbers they carry are such that these quantum corrections remain zero. They can also be used to cancel the gauge anomalies that are also present with these gauge bosons in combination with the SM gauge fields.

Appendix B

Further Details of Chapter 3 Calculations

B.1 F_+ Coefficients, Eq. (3.20)

Matching superhorizon modes with the plane waves, we obtain the following relation,

$$C_1 = \frac{\Gamma(\frac{3-\Omega_k}{4})}{2^{\frac{-1}{4}(1-\Omega_k)}\sqrt{\pi}} \left(\frac{1}{\sqrt{2k}} - C_2 \frac{2^{\frac{-1}{4}(1+\Omega_k)}\sqrt{\pi}}{\Gamma(\frac{3+\Omega_k}{4})} \right). \quad (\text{B.1})$$

The Wronskian normalisation implies:

$$\sqrt{\frac{2}{\Omega_k}} C_1 C_2 \sin\left(\frac{\pi}{4}(1+\Omega_k)\right) + C_2^2 \sqrt{\frac{\pi}{\Omega_k}} \frac{1}{\Gamma(\frac{1+\Omega_k}{2})} = \frac{1}{2k}. \quad (\text{B.2})$$

Solving the above conditions we find that the coefficients for the F_+ modes are,

$$C_1 = \frac{2^{-\frac{1}{4}(1+\Omega_k)}\Gamma(\frac{3-\Omega_k}{4})}{\sqrt{\pi k}} - \frac{2^{-\frac{1}{2}(\Omega_k+3)}\Gamma(\frac{1+\Omega_k}{4})\Gamma(\frac{3-\Omega_k}{4})}{\Gamma(\frac{3+\Omega_k}{4})} \sqrt{\frac{\Omega_k}{\pi k}}, \quad (\text{B.3})$$

and

$$C_2 = \frac{\Gamma(\frac{1+\Omega_k}{4})}{2\sqrt{2\pi}} \sqrt{\frac{\Omega_k}{k}} = \frac{\Gamma(\frac{1+\Omega_k}{4})}{2\sqrt{2\pi}} \left(\frac{k}{\kappa}\right)^{\frac{1}{4}}. \quad (\text{B.4})$$

B.2 F_- Coefficients, Eq. (3.21)

Similarly as above, we obtain the following relations from the matching,

$$C_4 = \frac{\Gamma(\frac{3-i\Omega_k}{4})}{2^{\frac{-1}{4}(1-i\Omega_k)}\sqrt{\pi}} \left(\frac{1}{\sqrt{2k}} - C_3 \frac{2^{\frac{-1}{4}(1+i\Omega_k)}\sqrt{\pi}}{\Gamma(\frac{3+i\Omega_k}{4})} \right), \quad (\text{B.5})$$

and the Wronskian normalisation,

$$C_3^2 + |C_4|^2 + 2C_3 e^{\frac{-\pi\Omega_k}{4}} \sqrt{2\pi} \text{Im} \left(\frac{\sqrt{i}C_4^*}{\Gamma(\frac{1+i\Omega_k}{2})} \right) = \frac{e^{\frac{-\pi\Omega_k}{4}}}{k} \sqrt{\frac{\Omega_k}{2}}. \quad (\text{B.6})$$

These two equations determine the coefficients for the F_- modes,

$$C_3 = \frac{1}{2\sqrt{2k}P(k)} \left(\sqrt{\Omega_k} e^{-\frac{\pi\Omega_k}{4}} - \frac{1}{\pi} \left| \Gamma\left(\frac{3-i\Omega_k}{4}\right) \right|^2 \right), \quad (\text{B.7})$$

and

$$C_4 = \frac{\Gamma(\frac{3-i\Omega_k}{4})}{2^{\frac{-1}{4}(1-i\Omega_k)}\sqrt{2\pi k}} \left(1 - \frac{\sqrt{\pi}}{2^{\frac{1}{4}(5+i\Omega_k)}P(k)\Gamma(\frac{3+i\Omega_k}{4})} \left(\sqrt{\Omega_k} e^{-\frac{\pi\Omega_k}{4}} - \frac{1}{\pi} \left| \Gamma\left(\frac{3-i\Omega_k}{4}\right) \right|^2 \right) \right), \quad (\text{B.8})$$

where

$$P(k) = \frac{2^{3/4}}{\sqrt{\pi}} \left(2\pi e^{-\frac{\pi\Omega_k}{4}} \text{Im} \left[\frac{\sqrt{i}}{2^{\frac{i\Omega_k}{4}}\Gamma(\frac{1+i\Omega_k}{4})} \right] - \text{Re} \left[\frac{\Gamma(\frac{3-i\Omega_k}{4})}{2^{\frac{i\Omega_k}{4}}} \right] \right). \quad (\text{B.9})$$

Appendix C

Further Details of Chapter 5 Calculations

C.1 Dimensional Regularisation Integrals and Useful Relations

The following dimensional regularisation Integrals were utilised in Chapter 5,

$$i \int \frac{d^N p}{(2\pi)^N} \frac{1}{(p^2 - m^2)^2} = -\frac{1}{16\pi^2} \left[\frac{4\pi^2 \lambda^2}{M^2} \right]^\epsilon \Gamma(\epsilon) , \quad (\text{C.1})$$

$$i \int \frac{d^N p}{(2\pi)^N} \frac{1}{(p^2 - m^2)^3} = \frac{1}{32\pi^2} \left[\frac{4\pi^2 \lambda^2}{M^2} \right]^\epsilon \frac{\Gamma(1 + \epsilon)}{M^2} , \quad (\text{C.2})$$

$$i \int \frac{d^N p}{(2\pi)^N} \frac{p_\mu p_\nu}{(p^2 - m^2)^2} = \frac{1}{32\pi^2} \left[\frac{4\pi^2 \lambda^2}{M^2} \right]^\epsilon M^2 \Gamma(\epsilon - 1) g_{\mu\nu} , \quad (\text{C.3})$$

$$i \int \frac{d^N p}{(2\pi)^N} \frac{p_\mu p_\nu}{(p^2 - m^2)^3} = -\frac{1}{64\pi^2} \left[\frac{4\pi^2 \lambda^2}{M^2} \right]^\epsilon \Gamma(\epsilon) g_{\mu\nu} , \quad (\text{C.4})$$

$$i \int \frac{d^N p}{(2\pi)^N} \frac{p_\mu p_\nu p_\rho p_\sigma}{(p^2 - m^2)^3} = \frac{1}{128\pi^2} \left[\frac{4\pi^2 \lambda^2}{M^2} \right]^\epsilon M^2 \Gamma(\epsilon - 1) (g_{\mu\nu} g_{\rho\sigma} + g_{\mu\sigma} g_{\nu\rho} + g_{\mu\rho} g_{\nu\sigma}) . \quad (\text{C.5})$$

Some other useful relations are,

$$\text{Tr}(\gamma_\mu \gamma_\alpha \gamma_\rho \gamma_\beta \gamma^5) = -4i \epsilon_{\mu\alpha\rho\beta} , \quad (\text{C.6})$$

$$\text{Tr}(\gamma_\mu(\not{p} + \not{k} + m)\gamma_\rho\sigma_{\alpha\beta}\gamma^5(\not{p} + m)) = 4\{\varepsilon_{\mu\rho\alpha\beta}[m^2 - p^2 - (kp)] - k^\lambda[\varepsilon_{\alpha\beta\rho\lambda}p_\mu - \varepsilon_{\alpha\beta\mu\lambda}p_\rho]\} , \quad (\text{C.7})$$

$$\text{Tr}(\gamma_\rho(\not{p} - \not{k} + m)\gamma_\mu\sigma_{\alpha\beta}\gamma^5(\not{p} + m)) = 4\{\varepsilon_{\mu\rho\alpha\beta}[p^2 - m^2 - (kp)] - k^\lambda[\varepsilon_{\alpha\beta\rho\lambda}p_\mu - \varepsilon_{\alpha\beta\mu\lambda}p_\rho]\} . \quad (\text{C.8})$$

Upon taking $\epsilon \rightarrow 0$ the following are obtained,

$$\Gamma(1 + \epsilon)|_{\epsilon \rightarrow 0} \simeq 1, \quad \Gamma(\epsilon)|_{\epsilon \rightarrow 0} \simeq \frac{1}{\epsilon} - \gamma, \quad \Gamma(\epsilon - 1)|_{\epsilon \rightarrow 0} \simeq -\frac{1}{\epsilon} + \gamma - 1 , \quad (\text{C.9})$$

$$\eta_{\mu\nu}\eta^{\mu\nu} \simeq 4 - 2\epsilon , \quad (\text{C.10})$$

$$\left[\frac{4\pi\lambda^2}{M^2} \right]^\epsilon |_{\epsilon \rightarrow 0} \simeq 1 + \epsilon \ln \left(\frac{4\pi\lambda^2}{M^2} \right) . \quad (\text{C.11})$$

Bibliography

- [1] Neil D. Barrie, Archil Kobakhidze, and Shelley Liang. Natural Inflation with Hidden Scale Invariance. *Phys. Lett.*, B756:390–393, 2016. doi: 10.1016/j.physletb.2016.03.056.
- [2] Neil D. Barrie and Archil Kobakhidze. Inflationary Baryogenesis in a Model with Gauged Baryon Number. *JHEP*, 09:163, 2014. doi: 10.1007/JHEP09(2014)163.
- [3] Neil D. Barrie and Archil Kobakhidze. Generating luminous and dark matter during inflation. *Modern Physics Letters A*, 32(14):1750087, 2017. doi: 10.1142/S0217732317500870.
- [4] Kazuharu Bamba, Neil D. Barrie, Akio Sugamoto, Tatsu Takeuchi, and Kimiko Yamashita. Ratchet Baryogenesis with an Analogy to the Forced Pendulum. arXiv:1610.03268, 2016.
- [5] Neil D. Barrie and Archil Kobakhidze. Gravitational Instabilities of the Cosmic Neutrino Background with Non-zero Lepton Number. *Phys. Lett.*, B772:459–463, 2017. doi: 10.1016/j.physletb.2017.07.012.
- [6] Neil D. Barrie, Archil Kobakhidze, Shelley Liang, Matthew Talia, and Lei Wu. Heavy Leptonium as the Origin of the 750 GeV Diphoton Excess. arXiv:1604.02803, 2016.
- [7] Neil D. Barrie, Archil Kobakhidze, Matthew Talia, and Lei Wu. 750 GeV Composite Axion as the LHC Diphoton Resonance. *Phys. Lett.*, B755:343–347, 2016. doi: 10.1016/j.physletb.2016.02.010.
- [8] Neil D. Barrie, Akio Sugamoto, and Kimiko Yamashita. Construction of a model of monopolium and its search via multiphoton channels at LHC. *PTEP*, 2016(11):113B02, 2016. doi: 10.1093/ptep/ptw155.
- [9] Chen-Ning Yang and Robert L. Mills. Conservation of Isotopic Spin and Isotopic Gauge Invariance. *Phys. Rev.*, 96:191–195, 1954. doi: 10.1103/PhysRev.96.191.
- [10] Ryoyu Utiyama. Invariant theoretical interpretation of interaction. *Phys. Rev.*, 101:1597–1607, 1956. doi: 10.1103/PhysRev.101.1597.
- [11] T. D. Lee and Chen-Ning Yang. Question of Parity Conservation in Weak Interactions. *Phys. Rev.*, 104:254–258, 1956. doi: 10.1103/PhysRev.104.254.

-
- [12] Julian S. Schwinger. A Theory of the Fundamental Interactions. *Annals Phys.*, 2:407–434, 1957. doi: 10.1016/0003-4916(57)90015-5.
- [13] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, and R. P. Hudson. Experimental Test of Parity Conservation in Beta Decay. *Phys. Rev.*, 105:1413–1414, 1957. doi: 10.1103/PhysRev.105.1413.
- [14] S. L. Glashow. Partial Symmetries of Weak Interactions. *Nucl. Phys.*, 22:579–588, 1961. doi: 10.1016/0029-5582(61)90469-2.
- [15] J. Goldstone. Field Theories with Superconductor Solutions. *Nuovo Cim.*, 19:154–164, 1961. doi: 10.1007/BF02812722.
- [16] Yoichiro Nambu and G. Jona-Lasinio. Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. 1. *Phys. Rev.*, 122:345–358, 1961. doi: 10.1103/PhysRev.122.345.
- [17] Yoichiro Nambu and G. Jona-Lasinio. Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. 2. *Phys. Rev.*, 124:246–254, 1961. doi: 10.1103/PhysRev.124.246.
- [18] Jeffrey Goldstone, Abdus Salam, and Steven Weinberg. Broken Symmetries. *Phys. Rev.*, 127:965–970, 1962. doi: 10.1103/PhysRev.127.965.
- [19] Ziro Maki, Masami Nakagawa, and Shoichi Sakata. Remarks on the unified model of elementary particles. *Prog. Theor. Phys.*, 28:870–880, 1962. doi: 10.1143/PTP.28.870.
- [20] Murray Gell-Mann. Symmetries of baryons and mesons. *Phys. Rev.*, 125:1067–1084, 1962. doi: 10.1103/PhysRev.125.1067.
- [21] Philip W. Anderson. Plasmons, Gauge Invariance, and Mass. *Phys. Rev.*, 130:439–442, 1963. doi: 10.1103/PhysRev.130.439.
- [22] Nicola Cabibbo. Unitary Symmetry and Leptonic Decays. *Phys. Rev. Lett.*, 10:531–533, 1963. doi: 10.1103/PhysRevLett.10.531.
- [23] G. Zweig. An SU(3) model for strong interaction symmetry and its breaking. Version 2. In *Developments in the Quark Theory of Hadrons. Vol. 1. 1964 - 1978*, pages 22–101. *Developments in the Quark Theory of Hadrons, Volume 1*. Edited by D. Lichtenberg and S. Rosen. pp. 22-101, 1964.
- [24] Murray Gell-Mann. A Schematic Model of Baryons and Mesons. *Phys. Lett.*, 8:214–215, 1964. doi: 10.1016/S0031-9163(64)92001-3.
- [25] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble. Global Conservation Laws and Massless Particles. *Phys. Rev. Lett.*, 13:585–587, 1964. doi: 10.1103/PhysRevLett.13.585.

- [26] Abdus Salam and John Clive Ward. Electromagnetic and weak interactions. *Phys. Lett.*, 13:168–171, 1964. doi: 10.1016/0031-9163(64)90711-5.
- [27] T. W. B. Kibble. Symmetry breaking in nonAbelian gauge theories. *Phys. Rev.*, 155:1554–1561, 1967. doi: 10.1103/PhysRev.155.1554.
- [28] Sidney R. Coleman and J. Mandula. All Possible Symmetries of the S Matrix. *Phys. Rev.*, 159:1251–1256, 1967. doi: 10.1103/PhysRev.159.1251.
- [29] Steven Weinberg. A Model of Leptons. *Phys. Rev. Lett.*, 19:1264–1266, 1967. doi: 10.1103/PhysRevLett.19.1264.
- [30] Abdus Salam. Weak and Electromagnetic Interactions. *Conf. Proc.*, C680519:367–377, 1968.
- [31] Kenneth G. Wilson. The Renormalization Group and Strong Interactions. *Phys. Rev.*, D3:1818, 1971. doi: 10.1103/PhysRevD.3.1818.
- [32] S. L. Glashow, J. Iliopoulos, and L. Maiani. Weak Interactions with Lepton-Hadron Symmetry. *Phys. Rev.*, D2:1285–1292, 1970. doi: 10.1103/PhysRevD.2.1285.
- [33] C. Bouchiat, J. Iliopoulos, and P. Meyer. An Anomaly Free Version of Weinberg’s Model. *Phys. Lett.*, 38B:519–523, 1972. doi: 10.1016/0370-2693(72)90532-1.
- [34] Steven Weinberg. Effects of a neutral intermediate boson in semileptonic processes. *Phys. Rev.*, D5:1412–1417, 1972. doi: 10.1103/PhysRevD.5.1412.
- [35] C. G. Bollini and J. J. Giambiagi. Dimensional Renormalization: The Number of Dimensions as a Regularizing Parameter. *Nuovo Cim.*, B12:20–26, 1972. doi: 10.1007/BF02895558.
- [36] Gerard ’t Hooft and M. J. G. Veltman. Regularization and Renormalization of Gauge Fields. *Nucl. Phys.*, B44:189–213, 1972. doi: 10.1016/0550-3213(72)90279-9.
- [37] K. G. Wilson and John B. Kogut. The Renormalization group and the epsilon expansion. *Phys. Rept.*, 12:75–200, 1974. doi: 10.1016/0370-1573(74)90023-4.
- [38] Makoto Kobayashi and Toshihide Maskawa. CP Violation in the Renormalizable Theory of Weak Interaction. *Prog. Theor. Phys.*, 49:652–657, 1973. doi: 10.1143/PTP.49.652.
- [39] Jogesh C. Pati and Abdus Salam. Lepton Number as the Fourth Color. *Phys. Rev.*, D10:275–289, 1974. doi: 10.1103/PhysRevD.10.275,10.1103/PhysRevD.11.703.2. [Erratum: *Phys. Rev.*D11,703(1975)].
- [40] John Iliopoulos. The Making of the Standard Theory. *Adv. Ser. Direct. High Energy Phys.*, 26:29–59, 2016. doi: 10.1142/9789814733519_0002.

-
- [41] K. A. Olive et al. Review of Particle Physics. *Chin. Phys.*, C38:090001, 2014. doi: 10.1088/1674-1137/38/9/090001.
- [42] Peter W. Higgs. Broken Symmetries and the Masses of Gauge Bosons. *Phys. Rev. Lett.*, 13:508–509, 1964. doi: 10.1103/PhysRevLett.13.508.
- [43] Peter W. Higgs. Broken symmetries, massless particles and gauge fields. *Phys. Lett.*, 12: 132–133, 1964. doi: 10.1016/0031-9163(64)91136-9.
- [44] F. Englert and R. Brout. Broken Symmetry and the Mass of Gauge Vector Mesons. *Phys. Rev. Lett.*, 13:321–323, 1964. doi: 10.1103/PhysRevLett.13.321.
- [45] Peter W. Higgs. Spontaneous Symmetry Breakdown without Massless Bosons. *Phys. Rev.*, 145:1156–1163, 1966. doi: 10.1103/PhysRev.145.1156.
- [46] Lyndon Evans and Philip Bryant. LHC Machine. *JINST*, 3:S08001, 2008. doi: 10.1088/1748-0221/3/08/S08001.
- [47] G. Aad et al. The ATLAS Experiment at the CERN Large Hadron Collider. *JINST*, 3: S08003, 2008. doi: 10.1088/1748-0221/3/08/S08003.
- [48] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012. doi: 10.1016/j.physletb.2012.08.020.
- [49] S. Chatrchyan et al. The CMS Experiment at the CERN LHC. *JINST*, 3:S08004, 2008. doi: 10.1088/1748-0221/3/08/S08004.
- [50] Serguei Chatrchyan et al. Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett.*, B716:30–61, 2012. doi: 10.1016/j.physletb.2012.08.021.
- [51] Steven Weinberg. *The Quantum theory of fields. Vol. 1: Foundations*. Cambridge University Press, 2005. ISBN 9780521670531, 9780511252044.
- [52] Steven Weinberg. *The quantum theory of fields. Vol. 2: Modern applications*. Cambridge University Press, 2013. ISBN 9781139632478, 9780521670548, 9780521550024.
- [53] L. H. Ryder. *Quantum Field Theory*. Cambridge University Press, 1996. ISBN 9780521478144, 9781139632393, 9780521237642.
- [54] Matthew D. Schwartz. *Quantum Field Theory and the Standard Model*. Cambridge University Press, 2014. ISBN 1107034736, 9781107034730.
- [55] W. N. Cottingham and D. A. Greenwood. *An introduction to the standard model of particle physics*. Cambridge University Press, 2007. ISBN 9780511271366, 9780521852494.

- [56] T. Morii, C. S. Lim, and S. N. Mukherjee. *The physics of the standard model and beyond*. River Edge, USA: World Scientific, 2004.
- [57] E. Leader and E. Predazzi. An Introduction to gauge theories and modern particle physics. Vol. 2: CP violation, QCD and hard processes. *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.*, 4:1–464, 1996.
- [58] E. Leader and E. Predazzi. *An Introduction to gauge theories and modern particle physics. Vol. 1: Electroweak interactions, the new particles and the parton model*. Cambridge University Press, 2011. ISBN 9780511885730, 9780521468404.
- [59] Sidney R. Coleman and Erick J. Weinberg. Radiative Corrections as the Origin of Spontaneous Symmetry Breaking. *Phys. Rev.*, D7:1888–1910, 1973. doi: 10.1103/PhysRevD.7.1888.
- [60] Steven Weinberg. Implications of Dynamical Symmetry Breaking. *Phys. Rev.*, D13:974–996, 1976. doi: 10.1103/PhysRevD.13.974.
- [61] Leonard Susskind. Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory. *Phys. Rev.*, D20:2619–2625, 1979. doi: 10.1103/PhysRevD.20.2619.
- [62] Steven Weinberg. Implications of Dynamical Symmetry Breaking: An Addendum. *Phys. Rev.*, D19:1277–1280, 1979. doi: 10.1103/PhysRevD.19.1277.
- [63] D. A. Kirzhnits and Andrei D. Linde. Macroscopic Consequences of the Weinberg Model. *Phys. Lett.*, 42B:471–474, 1972. doi: 10.1016/0370-2693(72)90109-8.
- [64] L. Dolan and R. Jackiw. Symmetry Behavior at Finite Temperature. *Phys. Rev.*, D9:3320–3341, 1974. doi: 10.1103/PhysRevD.9.3320.
- [65] Steven Weinberg. Gauge and Global Symmetries at High Temperature. *Phys. Rev.*, D9:3357–3378, 1974. doi: 10.1103/PhysRevD.9.3357.
- [66] R. D. Peccei and Helen R. Quinn. CP Conservation in the Presence of Instantons. *Phys. Rev. Lett.*, 38:1440–1443, 1977. doi: 10.1103/PhysRevLett.38.1440.
- [67] M. Lindner. Implications of Triviality for the Standard Model. *Z. Phys.*, C31:295, 1986. doi: 10.1007/BF01479540.
- [68] E. Noether. Invariant variation problems. *Transport Theory and Statistical Physics*, 1:186–207, 1971. doi: 10.1080/00411457108231446.
- [69] S. Rajpoot. Gauge symmetries of electroweak interactions. *Int.J.Theor.Phys.*, 27:689, 1988. doi: 10.1007/BF00669312.

- [70] Stephen L. Adler. Axial vector vertex in spinor electrodynamics. *Phys.Rev.*, 177:2426–2438, 1969. doi: 10.1103/PhysRev.177.2426.
- [71] J.S. Bell and R. Jackiw. A PCAC puzzle: $\pi^0 \rightarrow \gamma \gamma$ in the sigma model. *Nuovo Cim.*, A60:47–61, 1969. doi: 10.1007/BF02823296.
- [72] Gerard 't Hooft. Symmetry Breaking Through Bell-Jackiw Anomalies. *Phys.Rev.Lett.*, 37:8–11, 1976. doi: 10.1103/PhysRevLett.37.8.
- [73] John Preskill. Gauge anomalies in an effective field theory. *Annals Phys.*, 210:323–379, 1991. doi: 10.1016/0003-4916(91)90046-B.
- [74] Michael B. Green and John H. Schwarz. Anomaly Cancellation in Supersymmetric D=10 Gauge Theory and Superstring Theory. *Phys.Lett.*, B149:117–122, 1984. doi: 10.1016/0370-2693(84)91565-X.
- [75] Pavel Fileviez Perez, Tao Han, Tong Li, and Michael J. Ramsey-Musolf. Leptoquarks and Neutrino Masses at the LHC. *Nucl.Phys.*, B819:139–176, 2009. doi: 10.1016/j.nuclphysb.2009.04.009.
- [76] P. V. Dong and H. N. Long. A simple model of gauged lepton and baryon charges. *Physics International*, 6, 1:23–32, 2010. doi: 10.3844/pisp.2015.23.32.
- [77] Michael Duerr, Pavel Fileviez Perez, and Mark B. Wise. Gauge Theory for Baryon and Lepton Numbers with Leptoquarks. *Phys.Rev.Lett.*, 110:231801, 2013. doi: 10.1103/PhysRevLett.110.231801.
- [78] E.C.G. Stueckelberg. Interaction energy in electrodynamics and in the field theory of nuclear forces. *Helv.Phys.Acta*, 11:225–244, 1938. doi: 10.5169/seals-110852.
- [79] Steven Weinberg. *Gravitation and Cosmology*. John Wiley and Sons, New York, 1972. ISBN 0471925675, 9780471925675.
- [80] Scott Dodelson. *Modern Cosmology*. Academic Press, Amsterdam, 2003. ISBN 9780122191411.
- [81] Mark Trodden and Sean M. Carroll. TASI lectures: Introduction to cosmology. In *Progress in string theory. Proceedings, Summer School, TASI 2003, Boulder, USA, June 2-27, 2003*, pages 703–793, 2004.
- [82] V. Mukhanov. *Physical Foundations of Cosmology*. Cambridge University Press, Oxford, 2005. ISBN 0521563984, 9780521563987.
- [83] Arno A. Penzias and Robert Woodrow Wilson. A Measurement of excess antenna temperature at 4080-Mc/s. *Astrophys. J.*, 142:419–421, 1965. doi: 10.1086/148307.

- [84] John C. Mather et al. A Preliminary measurement of the Cosmic Microwave Background spectrum by the Cosmic Background Explorer (COBE) satellite. *Astrophys. J.*, 354:L37–L40, 1990. doi: 10.1086/185717.
- [85] George F. Smoot et al. Structure in the COBE differential microwave radiometer first year maps. *Astrophys. J.*, 396:L1–L5, 1992. doi: 10.1086/186504.
- [86] D. N. Spergel et al. First year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Determination of cosmological parameters. *Astrophys. J. Suppl.*, 148:175–194, 2003. doi: 10.1086/377226.
- [87] D. N. Spergel et al. Wilkinson Microwave Anisotropy Probe (WMAP) three year results: implications for cosmology. *Astrophys. J. Suppl.*, 170:377, 2007. doi: 10.1086/513700.
- [88] E. Komatsu et al. Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation. *Astrophys. J. Suppl.*, 180:330–376, 2009. doi: 10.1088/0067-0049/180/2/330.
- [89] E. Komatsu et al. Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation. *Astrophys. J. Suppl.*, 192:18, 2011. doi: 10.1088/0067-0049/192/2/18.
- [90] C. L. Bennett et al. Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results. *Astrophys. J. Suppl.*, 208:20, 2013. doi: 10.1088/0067-0049/208/2/20.
- [91] G. Hinshaw et al. Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results. *Astrophys. J. Suppl.*, 208:19, 2013. doi: 10.1088/0067-0049/208/2/19.
- [92] P. A. R. Ade et al. Planck 2013 results. I. Overview of products and scientific results. *Astron. Astrophys.*, 571:A1, 2014. doi: 10.1051/0004-6361/201321529.
- [93] R. Adam et al. Planck 2015 results. I. Overview of products and scientific results. *Astron. Astrophys.*, 594:A1, 2016. doi: 10.1051/0004-6361/201527101.
- [94] P. A. R. Ade et al. Planck 2013 results. XVI. Cosmological parameters. *Astron. Astrophys.*, 571:A16, 2014. doi: 10.1051/0004-6361/201321591.
- [95] J. B. Hartle. *An introduction to Einstein's general relativity*. San Francisco, USA: Addison-Wesley, 2003.
- [96] Edwin Hubble. A relation between distance and radial velocity among extra-galactic nebulae. *Proc. Nat. Acad. Sci.*, 15:168–173, 1929. doi: 10.1073/pnas.15.3.168.

- [97] G. Gamow. Expanding universe and the origin of elements. *Phys. Rev.*, 70:572–573, 1946. doi: 10.1103/PhysRev7.0.572.
- [98] R. A. Alpher, H. Bethe, and G. Gamow. The origin of chemical elements. *Phys. Rev.*, 73:803–804, 1948. doi: 10.1103/PhysRev.73.803.
- [99] Robert V. Wagoner, William A. Fowler, and Fred Hoyle. On the Synthesis of elements at very high temperatures. *Astrophys. J.*, 148:3–49, 1967. doi: 10.1086/149126.
- [100] David N. Schramm and Robert V. Wagoner. Element production in the early universe. *Ann. Rev. Nucl. Part. Sci.*, 27:37–74, 1977. doi: 10.1146/annurev.ns.27.120177.000345.
- [101] Jong-Mann Yang, Michael S. Turner, G. Steigman, D. N. Schramm, and Keith A. Olive. Primordial Nucleosynthesis: A Critical Comparison of Theory and Observation. *Astrophys. J.*, 281:493–511, 1984. doi: 10.1086/162123.
- [102] Ann Merchant Boesgaard and Gary Steigman. Big Bang Nucleosynthesis: Theories and Observations. *Ann. Rev. Astron. Astrophys.*, 23:319–378, 1985. doi: 10.1146/annurev.aa.23.090185.001535.
- [103] Craig J. Copi, David N. Schramm, and Michael S. Turner. Big bang nucleosynthesis and the baryon density of the universe. *Science*, 267:192–199, 1995. doi: 10.1126/science.7809624.
- [104] Gary Steigman. Primordial Nucleosynthesis in the Precision Cosmology Era. *Ann. Rev. Nucl. Part. Sci.*, 57:463–491, 2007. doi: 10.1146/annurev.nucl.56.080805.140437.
- [105] Fabio Iocco, Gianpiero Mangano, Gennaro Miele, Ofelia Pisanti, and Pasquale D. Serpico. Primordial Nucleosynthesis: from precision cosmology to fundamental physics. *Phys. Rept.*, 472:1–76, 2009. doi: 10.1016/j.physrep.2009.02.002.
- [106] Richard H. Cyburt, Brian D. Fields, and Keith A. Olive. An Update on the big bang nucleosynthesis prediction for Li-7: The problem worsens. *JCAP*, 0811:012, 2008. doi: 10.1088/1475-7516/2008/11/012.
- [107] Richard H. Cyburt, Brian D. Fields, Keith A. Olive, and Tsung-Han Yeh. Big Bang Nucleosynthesis: 2015. *Rev. Mod. Phys.*, 88:015004, 2016. doi: 10.1103/RevModPhys.88.015004.
- [108] Daniel Baumann. On the Quantum Origin of Structure in the Inflationary Universe. arXiv: 0710.3187, 2007.
- [109] Alan H. Guth. The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems. *Phys. Rev.*, D23:347–356, 1981. doi: 10.1103/PhysRevD.23.347.

- [110] Alexei A. Starobinsky. A New Type of Isotropic Cosmological Models Without Singularity. *Phys. Lett.*, B91:99–102, 1980. doi: 10.1016/0370-2693(80)90670-X.
- [111] D. Kazanas. Dynamics of the Universe and Spontaneous Symmetry Breaking. *Astrophys. J.*, 241:L59–L63, 1980. doi: 10.1086/183361.
- [112] K. Sato. First Order Phase Transition of a Vacuum and Expansion of the Universe. *Mon. Not. Roy. Astron. Soc.*, 195:467–479, 1981.
- [113] Andrei D. Linde. A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems. *Phys. Lett.*, B108:389–393, 1982. doi: 10.1016/0370-2693(82)91219-9.
- [114] Andrei D. Linde. Coleman-Weinberg Theory and a New Inflationary Universe Scenario. *Phys. Lett.*, B114:431–435, 1982. doi: 10.1016/0370-2693(82)90086-7.
- [115] Andreas Albrecht and Paul J. Steinhardt. Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking. *Phys. Rev. Lett.*, 48:1220–1223, 1982. doi: 10.1103/PhysRevLett.48.1220.
- [116] Alan H. Guth and S. Y. Pi. Fluctuations in the New Inflationary Universe. *Phys. Rev. Lett.*, 49:1110–1113, 1982. doi: 10.1103/PhysRevLett.49.1110.
- [117] Michael S. Turner. Coherent Scalar Field Oscillations in an Expanding Universe. *Phys. Rev.*, D28:1243, 1983. doi: 10.1103/PhysRevD.28.1243.
- [118] James M. Bardeen, Paul J. Steinhardt, and Michael S. Turner. Spontaneous Creation of Almost Scale - Free Density Perturbations in an Inflationary Universe. *Phys. Rev.*, D28:679, 1983. doi: 10.1103/PhysRevD.28.679.
- [119] Andrei D. Linde. The Inflationary Universe. *Rept. Prog. Phys.*, 47:925–986, 1984. doi: 10.1088/0034-4885/47/8/002.
- [120] Robert H. Brandenberger. Quantum Field Theory Methods and Inflationary Universe Models. *Rev. Mod. Phys.*, 57:1, 1985. doi: 10.1103/RevModPhys.57.1.
- [121] Andrei D. Linde. Eternal Chaotic Inflation. *Mod. Phys. Lett.*, A1:81, 1986. doi: 10.1142/S0217732386000129.
- [122] Daniel Baumann. Inflation. In *Physics of the large and the small, TASI 09, proceedings of the Theoretical Advanced Study Institute in Elementary Particle Physics, Boulder, Colorado, USA, 1-26 June 2009*, pages 523–686, 2011. doi: 10.1142/9789814327183_0010.
- [123] Gerard 't Hooft. Magnetic Monopoles in Unified Gauge Theories. *Nucl. Phys.*, B79:276–284, 1974. doi: 10.1016/0550-3213(74)90486-6.

- [124] Ya. B. Zeldovich and M. Yu. Khlopov. On the Concentration of Relic Magnetic Monopoles in the Universe. *Phys. Lett.*, B79:239–241, 1978. doi: 10.1016/0370-2693(78)90232-0.
- [125] John Preskill. Cosmological Production of Superheavy Magnetic Monopoles. *Phys. Rev. Lett.*, 43:1365, 1979. doi: 10.1103/PhysRevLett.43.1365.
- [126] Viatcheslav F. Mukhanov and G. V. Chibisov. Quantum Fluctuations and a Nonsingular Universe. *JETP Lett.*, 33:532–535, 1981. [Pisma Zh. Eksp. Teor. Fiz.33,549(1981)].
- [127] Keith A. Olive. Inflation. *Phys. Rept.*, 190:307–403, 1990. doi: 10.1016/0370-1573(90)90144-Q.
- [128] Edward W. Kolb and Michael S. Turner. The Early Universe. *Front. Phys.*, 69:1–547, 1990.
- [129] Antonio Riotto. Inflation and the theory of cosmological perturbations. In *Astroparticle physics and cosmology. Proceedings: Summer School, Trieste, Italy, Jun 17-Jul 5 2002*, pages 317–413, 2002.
- [130] Robert H. Brandenberger. Lectures on the theory of cosmological perturbations. *Lect. Notes Phys.*, 646:127–167, 2004.
- [131] Andrei D. Linde. Particle physics and inflationary cosmology. *Contemp. Concepts Phys.*, 5:1–362, 1990.
- [132] Karim A. Malik and David Wands. Cosmological perturbations. *Phys. Rept.*, 475:1–51, 2009. doi: 10.1016/j.physrep.2009.03.001.
- [133] Robert H. Brandenberger. Alternatives to the inflationary paradigm of structure formation. *Int. J. Mod. Phys. Conf. Ser.*, 01:67–79, 2011. doi: 10.1142/S2010194511000109.
- [134] Robert H. Brandenberger. Cosmology of the Very Early Universe. *AIP Conf. Proc.*, 1268: 3–70, 2010. doi: 10.1063/1.3483879.
- [135] Robert H. Brandenberger. Unconventional Cosmology. *Lect. Notes Phys.*, 863:333, 2013. doi: 10.1007/978-3-642-33036-0_12.
- [136] F. Lucchin and S. Matarrese. Power Law Inflation. *Phys. Rev.*, D32:1316, 1985. doi: 10.1103/PhysRevD.32.1316.
- [137] Katherine Freese, Joshua A. Frieman, and Angela V. Olinto. Natural inflation with pseudo - Nambu-Goldstone bosons. *Phys. Rev. Lett.*, 65:3233–3236, 1990. doi: 10.1103/PhysRevLett.65.3233.
- [138] John D. Barrow and Kei-ichi Maeda. Extended inflationary universes. *Nucl. Phys.*, B341: 294–308, 1990. doi: 10.1016/0550-3213(90)90272-F.

- [139] A. L. Berkin and Kei-Ichi Maeda. Inflation in generalized Einstein theories. *Phys. Rev.*, D44:1691–1704, 1991. doi: 10.1103/PhysRevD.44.1691.
- [140] Andrei D. Linde. Axions in inflationary cosmology. *Phys. Lett.*, B259:38–47, 1991. doi: 10.1016/0370-2693(91)90130-I.
- [141] L. F. Abbott, Edward Farhi, and Mark B. Wise. Particle Production in the New Inflationary Cosmology. *Phys. Lett.*, B117:29, 1982. doi: 10.1016/0370-2693(82)90867-X.
- [142] S. W. Hawking. The Development of Irregularities in a Single Bubble Inflationary Universe. *Phys. Lett.*, B115:295, 1982. doi: 10.1016/0370-2693(82)90373-2.
- [143] Alexei A. Starobinsky. Dynamics of Phase Transition in the New Inflationary Universe Scenario and Generation of Perturbations. *Phys. Lett.*, B117:175–178, 1982. doi: 10.1016/0370-2693(82)90541-X.
- [144] Fred C. Adams, J. Richard Bond, Katherine Freese, Joshua A. Frieman, and Angela V. Olinto. Natural inflation: Particle physics models, power law spectra for large scale structure, and constraints from COBE. *Phys. Rev.*, D47:426–455, 1993. doi: 10.1103/PhysRevD.47.426.
- [145] Andrei D. Linde. Chaotic Inflation. *Phys. Lett.*, B129:177–181, 1983. doi: 10.1016/0370-2693(83)90837-7.
- [146] Andrei D. Linde. Hybrid inflation. *Phys. Rev.*, D49:748–754, 1994. doi: 10.1103/PhysRevD.49.748.
- [147] Edmund J. Copeland, Andrew R. Liddle, David H. Lyth, Ewan D. Stewart, and David Wands. False vacuum inflation with Einstein gravity. *Phys. Rev.*, D49:6410–6433, 1994. doi: 10.1103/PhysRevD.49.6410.
- [148] Arjun Berera. Warm inflation. *Phys. Rev. Lett.*, 75:3218–3221, 1995. doi: 10.1103/PhysRevLett.75.3218.
- [149] P. Binetruy and G. R. Dvali. D term inflation. *Phys. Lett.*, B388:241–246, 1996. doi: 10.1016/S0370-2693(96)01083-0.
- [150] G. R. Dvali and S. H. Henry Tye. Brane inflation. *Phys. Lett.*, B450:72–82, 1999. doi: 10.1016/S0370-2693(99)00132-X.
- [151] David H. Lyth and Antonio Riotto. Particle physics models of inflation and the cosmological density perturbation. *Phys. Rept.*, 314:1–146, 1999. doi: 10.1016/S0370-1573(98)00128-8.
- [152] Andrew R. Liddle, Anupam Mazumdar, and Franz E. Schunck. Assisted inflation. *Phys. Rev.*, D58:061301, 1998. doi: 10.1103/PhysRevD.58.061301.

- [153] C. Armendariz-Picon, T. Damour, and Viatcheslav F. Mukhanov. k - inflation. *Phys. Lett.*, B458:209–218, 1999. doi: 10.1016/S0370-2693(99)00603-6.
- [154] Anupam Mazumdar. Extra dimensions and inflation. *Phys.Lett.*, B469:55–60, 1999. doi: 10.1016/S0370-2693(99)01256-3.
- [155] Lotfi Boubekeur and David.H. Lyth. Hilltop inflation. *JCAP*, 0507:010, 2005. doi: 10.1088/1475-7516/2005/07/010.
- [156] S. Dimopoulos, S. Kachru, J. McGreevy, and Jay G. Wacker. N-flation. *JCAP*, 0808:003, 2008. doi: 10.1088/1475-7516/2008/08/003.
- [157] Jerome Martin, Christophe Ringeval, and Vincent Vennin. Encyclopædia Inflationaris. *Phys. Dark Univ.*, 5-6:75–235, 2014. doi: 10.1016/j.dark.2014.01.003.
- [158] L. F. Abbott and Mark B. Wise. Constraints on Generalized Inflationary Cosmologies. *Nucl. Phys.*, B244:541–548, 1984. doi: 10.1016/0550-3213(84)90329-8.
- [159] Fred C. Adams, Katherine Freese, and Alan H. Guth. Constraints on the scalar field potential in inflationary models. *Phys. Rev.*, D43:965–976, 1991. doi: 10.1103/PhysRevD.43.965.
- [160] David H. Lyth. What would we learn by detecting a gravitational wave signal in the cosmic microwave background anisotropy? *Phys. Rev. Lett.*, 78:1861–1863, 1997. doi: 10.1103/PhysRevLett.78.1861.
- [161] David N. Spergel and Matias Zaldarriaga. CMB polarization as a direct test of inflation. *Phys. Rev. Lett.*, 79:2180–2183, 1997. doi: 10.1103/PhysRevLett.79.2180.
- [162] Andrew R Liddle and Samuel M Leach. How long before the end of inflation were observable perturbations produced? *Phys. Rev.*, D68:103503, 2003. doi: 10.1103/PhysRevD.68.103503.
- [163] Laila Alabidi and David H. Lyth. Inflation models and observation. *JCAP*, 0605:016, 2006. doi: 10.1088/1475-7516/2006/05/016.
- [164] Jessica L. Cook and Lorenzo Sorbo. Particle production during inflation and gravitational waves detectable by ground-based interferometers. *Phys. Rev.*, D85:023534, 2012. doi: 10.1103/PhysRevD.86.069901,10.1103/PhysRevD.85.023534. [Erratum: *Phys. Rev.D*86,069901(2012)].
- [165] David H. Lyth. The CMB modulation from inflation. *JCAP*, 1308:007, 2013. doi: 10.1088/1475-7516/2013/08/007.

- [166] Jérôme Martin, Christophe Ringeval, Roberto Trotta, and Vincent Vennin. The Best Inflationary Models After Planck. *JCAP*, 1403:039, 2014. doi: 10.1088/1475-7516/2014/03/039.
- [167] Diederik Roest. Universality classes of inflation. *JCAP*, 1401:007, 2014. doi: 10.1088/1475-7516/2014/01/007.
- [168] Mario Galante, Renata Kallosh, Andrei Linde, and Diederik Roest. Unity of Cosmological Inflation Attractors. *Phys. Rev. Lett.*, 114(14):141302, 2015. doi: 10.1103/PhysRevLett.114.141302.
- [169] P. Binetruy, E. Kiritsis, J. Mabillard, M. Pieroni, and C. Rosset. Universality classes for models of inflation. *JCAP*, 1504(04):033, 2015. doi: 10.1088/1475-7516/2015/04/033.
- [170] Valerie Domcke, Mauro Pieroni, and Pierre Binétruy. Primordial gravitational waves for universality classes of pseudoscalar inflation. *JCAP*, 1606:031, 2016. doi: 10.1088/1475-7516/2016/06/031.
- [171] Hong-Yee Chiu. Symmetry between particle and anti-particle populations in the universe. *Phys. Rev. Lett.*, 17:712, 1966. doi: 10.1103/PhysRevLett.17.712.
- [172] G. Steigman. Observational tests of antimatter cosmologies. *Ann. Rev. Astron. Astrophys.*, 14:339–372, 1976. doi: 10.1146/annurev.aa.14.090176.002011.
- [173] A. Yu. Ignatiev, N. V. Krasnikov, V. A. Kuzmin, and A. N. Tavkhelidze. Universal CP Noninvariant Superweak Interaction and Baryon Asymmetry of the Universe. *Phys. Lett.*, B76:436–438, 1978. doi: 10.1016/0370-2693(78)90900-0.
- [174] D. Toussaint, S. B. Treiman, Frank Wilczek, and A. Zee. Matter - Antimatter Accounting, Thermodynamics, and Black Hole Radiation. *Phys. Rev.*, D19:1036–1045, 1979. doi: 10.1103/PhysRevD.19.1036.
- [175] Savas Dimopoulos and Leonard Susskind. On the Baryon Number of the Universe. *Phys. Rev.*, D18:4500–4509, 1978. doi: 10.1103/PhysRevD.18.4500.
- [176] Motohiko Yoshimura. Origin of Cosmological Baryon Asymmetry. *Phys. Lett.*, B88:294–298, 1979. doi: 10.1016/0370-2693(79)90471-4.
- [177] Steven Weinberg. Cosmological Production of Baryons. *Phys. Rev. Lett.*, 42:850–853, 1979. doi: 10.1103/PhysRevLett.42.850.
- [178] Edward W. Kolb and Stephen Wolfram. Baryon Number Generation in the Early Universe. *Nucl. Phys.*, B172:224, 1980. doi: 10.1016/0550-3213(80)90167-4,10.1016/0550-3213(82)90012-8. [Erratum: *Nucl. Phys.*B195,542(1982)].

- [179] A. D. Dolgov and Ya. B. Zeldovich. Cosmology and Elementary Particles. *Rev. Mod. Phys.*, 53:1–41, 1981. doi: 10.1103/RevModPhys.53.1.
- [180] A. D. Dolgov. NonGUT baryogenesis. *Phys. Rept.*, 222:309–386, 1992. doi: 10.1016/0370-1573(92)90107-B.
- [181] W. Buchmuller and T. Yanagida. Baryogenesis and the scale of B-L breaking. *Phys. Lett.*, B302:240–244, 1993. doi: 10.1016/0370-2693(93)90391-T.
- [182] W. Buchmuller and T. Yanagida. Quark lepton mass hierarchies and the baryon asymmetry. *Phys. Lett.*, B445:399–402, 1999. doi: 10.1016/S0370-2693(98)01480-4.
- [183] W. Buchmuller. Some aspects of baryogenesis and lepton number violation. In *Recent developments in particle physics and cosmology: Proceedings. NATO ASI 2000. Cascais, Portugal, July 26 - Jul 7, 2000*, pages 281–314, 2000.
- [184] W. Buchmuller, P. Di Bari, and M. Plumacher. Cosmic microwave background, matter - antimatter asymmetry and neutrino masses. *Nucl. Phys.*, B643:367–390, 2002. doi: 10.1016/S0550-3213(02)00737-X,10.1016/j.nuclphysb.2007.11.030. [Erratum: Nucl. Phys.B793,362(2008)].
- [185] P. A. M. Dirac. A Theory of Electrons and Protons. *Proc. Roy. Soc. Lond.*, A126:360, 1930. doi: 10.1098/rspa.1930.0013.
- [186] J.H. Christenson, J.W. Cronin, V.L. Fitch, and R. Turlay. Evidence for the 2 pi Decay of the K^0 Meson. *Phys.Rev.Lett.*, 13:138–140, 1964. doi: 10.1103/PhysRevLett.13.138.
- [187] Giancarlo D’Ambrosio and Gino Isidori. CP violation in kaon decays. *Int.J.Mod.Phys.*, A13:1–94, 1998. doi: 10.1142/S0217751X98000020.
- [188] R. Aaij et al. Evidence for CP violation in time-integrated $D^0 \rightarrow h^- h^+$ decay rates. *Phys.Rev.Lett.*, 108:111602, 2012. doi: 10.1103/PhysRevLett.108.129903,10.1103/PhysRevLett.108.111602.
- [189] R Aaij et al. First observation of CP violation in the decays of B_s^0 mesons. *Phys.Rev.Lett.*, 110:221601, 2013. doi: 10.1103/PhysRevLett.110.221601.
- [190] Guennadi Borissov, Robert Fleischer, and Marie-Hélène Schune. Rare decays and cp violation in the bs system. *Annual Review of Nuclear and Particle Science*, 63(1):null, 2013. doi: 10.1146/annurev-nucl-102912-144527.
- [191] S.P. Ahlen, S. Barwick, J.J. Beatty, C.R. Bower, G. Gerbier, et al. New Limit on the Low-Energy Anti-Proton / Proton Ratio in the Galactic Cosmic Radiation. *Phys.Rev.Lett.*, 61:145–148, 1988. doi: 10.1103/PhysRevLett.61.145.

- [192] J. Alcaraz et al. Search for anti-helium in cosmic rays. *Phys.Lett.*, B461:387–396, 1999. doi: 10.1016/S0370-2693(99)00874-6.
- [193] P. A. R. Ade et al. Planck 2015 results. XIII. Cosmological parameters. *Astron. Astrophys.*, 594:A13, 2016. doi: 10.1051/0004-6361/201525830.
- [194] J. Beringer et al. Review of Particle Physics (RPP). *Phys.Rev.*, D86:010001, 2012. doi: 10.1103/PhysRevD.86.010001.
- [195] Vimal Simha and Gary Steigman. Constraining The Early-Universe Baryon Density And Expansion Rate. *JCAP*, 0806:016, 2008. doi: 10.1088/1475-7516/2008/06/016.
- [196] Gary Steigman. Primordial Nucleosynthesis: The Predicted and Observed Abundances and Their Consequences. *PoS*, NICXI:001, 2010.
- [197] Brian D. Fields, Paolo Molaro, and Subir Sarkar. Big-Bang Nucleosynthesis. *Chin. Phys.*, C38:339–344, 2014.
- [198] F.W. Stecker, D.L. Morgan, and J. Bredekamp. Possible evidence for the existence of antimatter on a cosmological scale in the universe. *Phys.Rev.Lett.*, 27:1469–1472, 1971. doi: 10.1103/PhysRevLett.27.1469.
- [199] Scott Dodelson and Lawrence M. Widrow. Baryogenesis in a Baryon Symmetric Universe. *Phys. Rev.*, D42:326–342, 1990. doi: 10.1103/PhysRevD.42.326.
- [200] Scott Dodelson and Lawrence M. Widrow. Baryon Symmetric Baryogenesis. *Phys. Rev. Lett.*, 64:340–343, 1990. doi: 10.1103/PhysRevLett.64.340.
- [201] D.L. Morgan and V.W. Hughes. Atomic processes involved in matter-antimatter annihilation. *Phys.Rev.*, D2:1389–1399, 1970. doi: 10.1103/PhysRevD.2.1389.
- [202] Laurent Canetti, Marco Drewes, and Mikhail Shaposhnikov. Matter and Antimatter in the Universe. *New J.Phys.*, 14:095012, 2012. doi: 10.1088/1367-2630/14/9/095012.
- [203] Andrew G. Cohen, A. De Rujula, and S.L. Glashow. A Matter - antimatter universe? *Astrophys.J.*, 495:539–549, 1998. doi: 10.1086/305328.
- [204] Andrew G. Cohen and A. De Rujula. Scars on the CBR? arXiv:astro-ph/9709132, 1997.
- [205] A. D. Sakharov. Violation of CP Invariance, c Asymmetry, and Baryon Asymmetry of the Universe. *Pisma Zh. Eksp. Teor. Fiz.*, 5:32–35, 1967. doi: 10.1070/PU1991v034n05ABEH002497. [Usp. Fiz. Nauk161,61(1991)].
- [206] Andrei D. Linde. Phase Transitions in Gauge Theories and Cosmology. *Rept. Prog. Phys.*, 42:389, 1979. doi: 10.1088/0034-4885/42/3/001.

- [207] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov. On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe. *Phys. Lett.*, B155:36, 1985. doi: 10.1016/0370-2693(85)91028-7.
- [208] M. E. Shaposhnikov. Structure of the High Temperature Gauge Ground State and Electroweak Production of the Baryon Asymmetry. *Nucl. Phys.*, B299:797–817, 1988. doi: 10.1016/0550-3213(88)90373-2.
- [209] M. E. Shaposhnikov. Baryon Asymmetry of the Universe in Standard Electroweak Theory. *Nucl. Phys.*, B287:757–775, 1987. doi: 10.1016/0550-3213(87)90127-1.
- [210] Neil Turok and John Zadrozny. Dynamical generation of baryons at the electroweak transition. *Phys. Rev. Lett.*, 65:2331–2334, 1990. doi: 10.1103/PhysRevLett.65.2331.
- [211] Michael Dine, Patrick Huet, and Robert L. Singleton, Jr. Baryogenesis at the electroweak scale. *Nucl. Phys.*, B375:625–648, 1992. doi: 10.1016/0550-3213(92)90113-P.
- [212] Greg W. Anderson and Lawrence J. Hall. The Electroweak phase transition and baryogenesis. *Phys. Rev.*, D45:2685–2698, 1992. doi: 10.1103/PhysRevD.45.2685.
- [213] Greg W. Anderson. Remarks on the electroweak phase transition. In *1st Yale-Texas Workshop on Baryon Number Violation at the Electroweak Scale New Haven, Connecticut, March 19-21, 1992*, pages 0134–143, 1992.
- [214] Andrew G. Cohen, D. B. Kaplan, and A. E. Nelson. Progress in electroweak baryogenesis. *Ann. Rev. Nucl. Part. Sci.*, 43:27–70, 1993. doi: 10.1146/annurev.ns.43.120193.000331.
- [215] Glennys R. Farrar and M. E. Shaposhnikov. Baryon asymmetry of the universe in the standard electroweak theory. *Phys. Rev.*, D50:774, 1994. doi: 10.1103/PhysRevD.50.774.
- [216] James M. Cline. Recent progress in electroweak baryogenesis. In *Strong and electroweak matter '98. Proceedings, Conference, SEWM'98, Copenhagen, Denmark, December 2-5, 1998*, pages 70–80, 1998.
- [217] Mark Trodden. Electroweak baryogenesis. *Rev. Mod. Phys.*, 71:1463–1500, 1999. doi: 10.1103/RevModPhys.71.1463.
- [218] Rabindra N. Mohapatra and Goran Senjanovic. Broken Symmetries at High Temperature. *Phys. Rev.*, D20:3390–3398, 1979. doi: 10.1103/PhysRevD.20.3390.
- [219] K. Sato. Cosmological Baryon Number Domain Structure and the First Order Phase Transition of a Vacuum. *Phys. Lett.*, B99:66–70, 1981. doi: 10.1016/0370-2693(81)90805-4.
- [220] Nikita Blinov. *Phase transitions: applications to physics beyond the Standard Model*. PhD thesis, British Columbia U., 2015.

- [221] M.B. Gavela, P. Hernandez, J. Orloff, and O. Pene. Standard model CP violation and baryon asymmetry. *Mod.Phys.Lett.*, A9:795–810, 1994. doi: 10.1142/S0217732394000629.
- [222] Michael Dine, Robert Leigh, Patrick Huet, Andrei Linde, and Dmitri Linde. Towards the theory of the electroweak phase transition. *Phys.Rev.*, D46:550–571, 1992. doi: 10.1103/PhysRevD.46.550.
- [223] K. Kajantie, M. Laine, K. Rummukainen, and Mikhail E. Shaposhnikov. Is there a hot electroweak phase transition at $m(H)$ larger or equal to $m(W)$? *Phys.Rev.Lett.*, 77:2887–2890, 1996. doi: 10.1103/PhysRevLett.77.2887.
- [224] Michael Dine and Alexander Kusenko. The Origin of the matter - antimatter asymmetry. *Rev. Mod. Phys.*, 76:1, 2003. doi: 10.1103/RevModPhys.76.1.
- [225] Antonio Riotto. Theories of baryogenesis. In *Proceedings, Summer School in High-energy physics and cosmology: Trieste, Italy, June 29-July 17, 1998*, pages 326–436, 1998.
- [226] Antonio Riotto and Mark Trodden. Recent progress in baryogenesis. *Ann.Rev.Nucl.Part.Sci.*, 49:35–75, 1999. doi: 10.1146/annurev.nucl.49.1.35.
- [227] Wilfried Buchmuller and Stefan Fredenhagen. Elements of baryogenesis. In *Current topics in astrophysical physics: The cosmic microwave background. Proceedings, NATO Advanced Study Institute, 8th Course, Erice, Italy, December 5-16, 1999*, pages 17–35, 2000.
- [228] James M. Cline. Baryogenesis. In *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime Les Houches, France, July 31-August 25, 2006*, 2006.
- [229] S.N. Gninenko, D.S. Gorbunov, and M.E. Shaposhnikov. Search for GeV-scale sterile neutrinos responsible for active neutrino oscillations and baryon asymmetry of the Universe. *Adv.High Energy Phys.*, 2012:718259, 2012. doi: 10.1155/2012/718259.
- [230] Wei-Min Yang. A Model of Four Generation Fermions and Cold Dark Matter and Matter-Antimatter Asymmetry. *Phys.Rev.*, D87:095003, 2013. doi: 10.1103/PhysRevD.87.095003.
- [231] Ian Affleck and Michael Dine. A New Mechanism for Baryogenesis. *Nucl.Phys.*, B249:361, 1985. doi: 10.1016/0550-3213(85)90021-5.
- [232] Rouzbeh Allahverdi and Anupam Mazumdar. A mini review on Affleck-Dine baryogenesis. *New J.Phys.*, 14:125013, 2012. doi: 10.1088/1367-2630/14/12/125013.
- [233] Karim Benakli and Sacha Davidson. Baryogenesis in models with a low quantum gravity scale. *Phys.Rev.*, D60:025004, 1999. doi: 10.1103/PhysRevD.60.025004.

- [234] Motohiko Yoshimura. Unified Gauge Theories and the Baryon Number of the Universe. *Phys. Rev. Lett.*, 41:281–284, 1978. doi: 10.1103/PhysRevLett.41.281. [Erratum: *Phys. Rev. Lett.*42,746(1979)].
- [235] Edward W. Kolb and Michael S. Turner. Grand Unified Theories and the Origin of the Baryon Asymmetry. *Ann. Rev. Nucl. Part. Sci.*, 33:645–696, 1983. doi: 10.1146/annurev.ns.33.120183.003241.
- [236] Rouzbeh Allahverdi, Bhaskar Dutta, and Kuver Sinha. Cladogenesis: Baryon-Dark Matter Coincidence from Branchings in Moduli Decay. *Phys.Rev.*, D83:083502, 2011. doi: 10.1103/PhysRevD.83.083502.
- [237] Matthew R. Buckley and Lisa Randall. Xogenesis. *JHEP*, 09:009, 2011. doi: 10.1007/JHEP09(2011)009.
- [238] Jessie Shelton and Kathryn M. Zurek. Darkogenesis: A baryon asymmetry from the dark matter sector. *Phys. Rev.*, D82:123512, 2010. doi: 10.1103/PhysRevD.82.123512.
- [239] Mattias Blennow, Basudeb Dasgupta, Enrique Fernandez-Martinez, and Nuria Rius. Aidnogenesis via Leptogenesis and Dark Sphalerons. *JHEP*, 03:014, 2011. doi: 10.1007/JHEP03(2011)014.
- [240] Laurent Canetti, Marco Drewes, Tibor Frossard, and Mikhail Shaposhnikov. Dark Matter, Baryogenesis and Neutrino Oscillations from Right Handed Neutrinos. *Phys.Rev.*, D87:093006, 2013. doi: 10.1103/PhysRevD.87.093006.
- [241] S. M. Boucenna and S. Morisi. Theories relating baryon asymmetry and dark matter: A mini review. *Front.in Phys.*, 1:33, 2014. doi: 10.3389/fphy.2013.00033.
- [242] Clifford Cheung and Yue Zhang. Electroweak Cogenesis. *JHEP*, 1309:002, 2013. doi: 10.1007/JHEP09(2013)002.
- [243] M. A. Luty. Baryogenesis via leptogenesis. *Phys. Rev.*, D45:455–465, 1992. doi: 10.1103/PhysRevD.45.455.
- [244] Apostolos Pilaftsis. Heavy Majorana neutrinos and baryogenesis. *Int. J. Mod. Phys.*, A14:1811–1858, 1999. doi: 10.1142/S0217751X99000932.
- [245] T. Asaka, Koichi Hamaguchi, M. Kawasaki, and T. Yanagida. Leptogenesis in inflationary universe. *Phys. Rev.*, D61:083512, 2000. doi: 10.1103/PhysRevD.61.083512.
- [246] W. Buchmuller and M. Plumacher. Neutrino masses and the baryon asymmetry. *Int. J. Mod. Phys.*, A15:5047–5086, 2000. doi: 10.1016/S0217-751X(00)00293-5,10.1142/S0217751X00002935.

- [247] W. Buchmuller, P. Di Bari, and M. Plumacher. Leptogenesis for pedestrians. *Annals Phys.*, 315:305–351, 2005. doi: 10.1016/j.aop.2004.02.003.
- [248] W. Buchmuller, R. D. Peccei, and T. Yanagida. Leptogenesis as the origin of matter. *Ann. Rev. Nucl. Part. Sci.*, 55:311–355, 2005. doi: 10.1146/annurev.nucl.55.090704.151558.
- [249] Sacha Davidson, Enrico Nardi, and Yosef Nir. Leptogenesis. *Phys.Rept.*, 466:105–177, 2008. doi: 10.1016/j.physrep.2008.06.002.
- [250] Raghavan Rangarajan and D. V. Nanopoulos. Inflationary baryogenesis. *Phys. Rev. D*, 64:063511, 2001. doi: 10.1103/PhysRevD.64.063511.
- [251] Stephon Haigh-Solom Alexander, Michael E. Peskin, and Mohammad M. Sheikh-Jabbari. Leptogenesis from gravity waves in models of inflation. *Phys.Rev.Lett.*, 96:081301, 2006. doi: 10.1103/PhysRevLett.96.081301.
- [252] Stephon Alexander, Antonino Marciano, and David Spergel. Chern-Simons Inflation and Baryogenesis. *JCAP*, 1304:046, 2013. doi: 10.1088/1475-7516/2013/04/046.
- [253] F. Zwicky. Die Rotverschiebung von extragalaktischen Nebeln. *Helv. Phys. Acta*, 6:110–127, 1933. doi: 10.1007/s10714-008-0707-4. [Gen. Rel. Grav.41,207(2009)].
- [254] Vera C. Rubin and W. Kent Ford, Jr. Rotation of the Andromeda Nebula from a Spectroscopic Survey of Emission Regions. *Astrophys. J.*, 159:379–403, 1970. doi: 10.1086/150317.
- [255] V. C. Rubin, N. Thonnard, and W. K. Ford, Jr. Rotational properties of 21 SC galaxies with a large range of luminosities and radii, from NGC 4605 / $R = 4\text{kpc}$ / to UGC 2885 / $R = 122\text{kpc}$ /. *Astrophys. J.*, 238:471, 1980. doi: 10.1086/158003.
- [256] V. C. Rubin, D. Burstein, W. K. Ford, Jr., and N. Thonnard. Rotation velocities of 16 SA galaxies and a comparison of Sa, Sb, and SC rotation properties. *Astrophys. J.*, 289:81, 1985. doi: 10.1086/162866.
- [257] Massimo Persic and Paolo Salucci. Rotation curves of 967 spiral galaxies. *Astrophys. J. Suppl.*, 99:501, 1995. doi: 10.1086/192195.
- [258] Massimo Persic, Paolo Salucci, and Fulvio Stel. The Universal rotation curve of spiral galaxies: 1. The Dark matter connection. *Mon. Not. Roy. Astron. Soc.*, 281:27, 1996. doi: 10.1093/mnras/281.1.27,10.1093/mnras/278.1.27.
- [259] Julio F. Navarro, Carlos S. Frenk, and Simon D. M. White. The Structure of cold dark matter halos. *Astrophys. J.*, 462:563–575, 1996. doi: 10.1086/177173.
- [260] W. J. G. de Blok, F. Walter, E. Brinks, C. Trachternach, S-H. Oh, and R. C. Kennicutt, Jr. High-Resolution Rotation Curves and Galaxy Mass Models from THINGS. *Astron. J.*, 136:2648–2719, 2008. doi: 10.1088/0004-6256/136/6/2648.

- [261] Dennis J. Hegyi and Keith A. Olive. Can Galactic Halos Be Made of Baryons? *Phys. Lett.*, B126:28, 1983. doi: 10.1016/0370-2693(83)90009-6.
- [262] Dennis J. Hegyi and Keith A. Olive. A Case Against Baryons in Galactic Halos. *Astrophys. J.*, 303:56–65, 1986. doi: 10.1086/164051.
- [263] G. Steigman, Keith A. Olive, and D. N. Schramm. Cosmological Constraints on Superweak Particles. *Phys. Rev. Lett.*, 43:239–242, 1979. doi: 10.1103/PhysRevLett.43.239.
- [264] Keith A. Olive, David N. Schramm, and Gary Steigman. Limits on New Superweakly Interacting Particles from Primordial Nucleosynthesis. *Nucl. Phys.*, B180:497–515, 1981. doi: 10.1016/0550-3213(81)90065-1.
- [265] Gary Steigman and Michael S. Turner. Cosmological Constraints on the Properties of Weakly Interacting Massive Particles. *Nucl. Phys.*, B253:375–386, 1985. doi: 10.1016/0550-3213(85)90537-1.
- [266] E. Aprile et al. Physics reach of the XENON1T dark matter experiment. *JCAP*, 1604(04):027, 2016. doi: 10.1088/1475-7516/2016/04/027.
- [267] D. S. Akerib et al. Results from a search for dark matter in the complete LUX exposure. *Phys. Rev. Lett.*, 118(2):021303, 2017. doi: 10.1103/PhysRevLett.118.021303.
- [268] Joel R. Primack, David Seckel, and Bernard Sadoulet. Detection of Cosmic Dark Matter. *Ann. Rev. Nucl. Part. Sci.*, 38:751–807, 1988. doi: 10.1146/annurev.ns.38.120188.003535.
- [269] Keith A. Olive. TASI lectures on dark matter. In *Particle physics and cosmology: The quest for physics beyond the standard model(s). Proceedings, Theoretical Advanced Study Institute, TASI 2002, Boulder, USA, June 3-28, 2002*, pages 797–851, 2003.
- [270] Lars Bergstrom. Dark Matter Candidates. *New J. Phys.*, 11:105006, 2009. doi: 10.1088/1367-2630/11/10/105006.
- [271] Katherine Garrett and Gintaras Duda. Dark Matter: A Primer. *Adv. Astron.*, 2011:968283, 2011. doi: 10.1155/2011/968283.
- [272] Mariangela Lisanti. Lectures on Dark Matter Physics. In *Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings (TASI 2015): Boulder, CO, USA, June 1-26, 2015*, pages 399–446, 2017. doi: 10.1142/9789813149441_0007.
- [273] David N. Schramm and Gary Steigman. Relic Neutrinos and the Density of the Universe. *Astrophys. J.*, 243:1, 1981. doi: 10.1086/158559.
- [274] M. Milgrom. A Modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *Astrophys. J.*, 270:365–370, 1983. doi: 10.1086/161130.

- [275] Paolo Gondolo and Graciela Gelmini. Cosmic abundances of stable particles: Improved analysis. *Nucl. Phys.*, B360:145–179, 1991. doi: 10.1016/0550-3213(91)90438-4.
- [276] Stephen M. Barr. Baryogenesis, sphalerons and the cogeneration of dark matter. *Phys. Rev.*, D44:3062–3066, 1991. doi: 10.1103/PhysRevD.44.3062.
- [277] Vadim A. Kuzmin. A Simultaneous solution to baryogenesis and dark matter problems. *Phys. Part. Nucl.*, 29:257–265, 1998. doi: 10.1134/1.953070. [Phys. Atom. Nucl.61,1107(1998)].
- [278] David H. Oaknin and Ariel Zhitnitsky. Baryon asymmetry, dark matter and quantum chromodynamics. *Phys. Rev.*, D71:023519, 2005. doi: 10.1103/PhysRevD.71.023519.
- [279] Ryuichiro Kitano and Ian Low. Dark matter from baryon asymmetry. *Phys. Rev.*, D71:023510, 2005. doi: 10.1103/PhysRevD.71.023510.
- [280] Glennys R. Farrar and Gabrijela Zaharijas. Dark matter and the baryon asymmetry. *Phys. Rev. Lett.*, 96:041302, 2006. doi: 10.1103/PhysRevLett.96.041302.
- [281] David E. Kaplan, Markus A. Luty, and Kathryn M. Zurek. Asymmetric Dark Matter. *Phys. Rev.*, D79:115016, 2009. doi: 10.1103/PhysRevD.79.115016.
- [282] Pei-Hong Gu, Utpal Sarkar, and Xinmin Zhang. Visible and Dark Matter Genesis and Cosmic Positron/Electron Excesses. *Phys. Rev.*, D80:076003, 2009. doi: 10.1103/PhysRevD.80.076003.
- [283] Hooman Davoudiasl, David E. Morrissey, Kris Sigurdson, and Sean Tulin. Hylogenesis: A Unified Origin for Baryonic Visible Matter and Antibaryonic Dark Matter. *Phys. Rev. Lett.*, 105:211304, 2010. doi: 10.1103/PhysRevLett.105.211304.
- [284] N. Haba and S. Matsumoto. Baryogenesis from Dark Sector. *Prog. Theor. Phys.*, 125:1311–1316, 2011. doi: 10.1143/PTP.125.1311.
- [285] Pei-Hong Gu, Manfred Lindner, Utpal Sarkar, and Xinmin Zhang. WIMP Dark Matter and Baryogenesis. *Phys. Rev.*, D83:055008, 2011. doi: 10.1103/PhysRevD.83.055008.
- [286] Lawrence J. Hall, John March-Russell, and Stephen M. West. A Unified Theory of Matter Genesis: Asymmetric Freeze-In. arXiv:1010.0245, 2010.
- [287] Bhaskar Dutta and Jason Kumar. Asymmetric Dark Matter from Hidden Sector Baryogenesis. *Phys. Lett.*, B699:364–367, 2011. doi: 10.1016/j.physletb.2011.04.036.
- [288] Adam Falkowski, Joshua T. Ruderman, and Tomer Volansky. Asymmetric Dark Matter from Leptogenesis. *JHEP*, 05:106, 2011. doi: 10.1007/JHEP05(2011)106.

-
- [289] Jonathan J. Heckman and Soo-Jong Rey. Baryon and Dark Matter Genesis from Strongly Coupled Strings. *JHEP*, 06:120, 2011. doi: 10.1007/JHEP06(2011)120.
- [290] Michael L. Graesser, Ian M. Shoemaker, and Luca Vecchi. Asymmetric WIMP dark matter. *JHEP*, 10:110, 2011. doi: 10.1007/JHEP10(2011)110.
- [291] Nicole F. Bell, Kalliopi Petraki, Ian M. Shoemaker, and Raymond R. Volkas. Pangenesis in a Baryon-Symmetric Universe: Dark and Visible Matter via the Affleck-Dine Mechanism. *Phys. Rev.*, D84:123505, 2011. doi: 10.1103/PhysRevD.84.123505.
- [292] Clifford Cheung and Kathryn M. Zurek. Affleck-Dine Cogenesis. *Phys. Rev.*, D84:035007, 2011. doi: 10.1103/PhysRevD.84.035007.
- [293] John March-Russell and Matthew McCullough. Asymmetric Dark Matter via Spontaneous Co-Genesis. *JCAP*, 1203:019, 2012. doi: 10.1088/1475-7516/2012/03/019.
- [294] Yanou Cui, Lisa Randall, and Brian Shuve. Emergent Dark Matter, Baryon, and Lepton Numbers. *JHEP*, 08:073, 2011. doi: 10.1007/JHEP08(2011)073.
- [295] Anupam Mazumdar. The origin of dark matter, matter-anti-matter asymmetry, and inflation. arXiv:1106.5408, 2011.
- [296] Tongyan Lin, Hai-Bo Yu, and Kathryn M. Zurek. On Symmetric and Asymmetric Light Dark Matter. *Phys. Rev.*, D85:063503, 2012. doi: 10.1103/PhysRevD.85.063503.
- [297] Kalliopi Petraki, Mark Trodden, and Raymond R. Volkas. Visible and dark matter from a first-order phase transition in a baryon-symmetric universe. *JCAP*, 1202:044, 2012. doi: 10.1088/1475-7516/2012/02/044.
- [298] Wan-Zhe Feng, Anupam Mazumdar, and Pran Nath. Baryogenesis from dark matter. *Phys. Rev.*, D88(3):036014, 2013. doi: 10.1103/PhysRevD.88.036014.
- [299] Kathryn M. Zurek. Asymmetric Dark Matter: Theories, Signatures, and Constraints. *Phys. Rept.*, 537:91–121, 2014. doi: 10.1016/j.physrep.2013.12.001.
- [300] Pierre Ramond. Neutrinos: A Glimpse beyond the standard model. *Nucl. Phys. Proc. Suppl.*, 77:3–9, 1999. doi: 10.1016/S0920-5632(99)00382-5.
- [301] Y. Fukuda et al. Measurement of a small atmospheric muon-neutrino / electron-neutrino ratio. *Phys. Lett.*, B433:9–18, 1998. doi: 10.1016/S0370-2693(98)00476-6.
- [302] Y. Fukuda et al. Study of the atmospheric neutrino flux in the multi-GeV energy range. *Phys. Lett.*, B436:33–41, 1998. doi: 10.1016/S0370-2693(98)00876-4.
- [303] Y. Fukuda et al. Evidence for oscillation of atmospheric neutrinos. *Phys. Rev. Lett.*, 81:1562–1567, 1998. doi: 10.1103/PhysRevLett.81.1562.

- [304] Y. Fukuda et al. Measurements of the solar neutrino flux from Super-Kamiokande's first 300 days. *Phys. Rev. Lett.*, 81:1158–1162, 1998. doi: 10.1103/PhysRevLett.81.1158. [Erratum: *Phys. Rev. Lett.*81,4279(1998)].
- [305] Y. Fukuda et al. Constraints on neutrino oscillation parameters from the measurement of day night solar neutrino fluxes at Super-Kamiokande. *Phys. Rev. Lett.*, 82:1810–1814, 1999. doi: 10.1103/PhysRevLett.82.1810.
- [306] Y. Fukuda et al. Measurement of the solar neutrino energy spectrum using neutrino electron scattering. *Phys. Rev. Lett.*, 82:2430–2434, 1999. doi: 10.1103/PhysRevLett.82.2430.
- [307] Q. R. Ahmad et al. Measurement of the rate of $\nu_e + d \rightarrow p + p + e^-$ interactions produced by ${}^8\text{B}$ solar neutrinos at the Sudbury Neutrino Observatory. *Phys. Rev. Lett.*, 87:071301, 2001. doi: 10.1103/PhysRevLett.87.071301.
- [308] Q. R. Ahmad et al. Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory. *Phys. Rev. Lett.*, 89:011301, 2002. doi: 10.1103/PhysRevLett.89.011301.
- [309] Q. R. Ahmad et al. Measurement of day and night neutrino energy spectra at SNO and constraints on neutrino mixing parameters. *Phys. Rev. Lett.*, 89:011302, 2002. doi: 10.1103/PhysRevLett.89.011302.
- [310] B. T. Cleveland, Timothy Daily, Raymond Davis, Jr., James R. Distel, Kenneth Lande, C. K. Lee, Paul S. Wildenhain, and Jack Ullman. Measurement of the solar electron neutrino flux with the Homestake chlorine detector. *Astrophys. J.*, 496:505–526, 1998. doi: 10.1086/305343.
- [311] V. N. Gavrin. Solar neutrino results from SAGE. *Nucl. Phys. Proc. Suppl.*, 91:36–43, 2001. doi: 10.1016/S0920-5632(00)00920-8.
- [312] E. Bellotti. First results from GNO. *Nucl. Phys. Proc. Suppl.*, 91:44–49, 2001. doi: 10.1016/S0920-5632(00)00921-X.
- [313] Kevork N. Abazajian. Sterile neutrinos in cosmology. arXiv:1705.01837, 2017.
- [314] J. Aasi et al. Advanced LIGO. *Class. Quant. Grav.*, 32:074001, 2015. doi: 10.1088/0264-9381/32/7/074001.
- [315] B. P. Abbott et al. Binary Black Hole Mergers in the first Advanced LIGO Observing Run. *Phys. Rev.*, X6(4):041015, 2016. doi: 10.1103/PhysRevX.6.041015.
- [316] B. P. Abbott et al. The Rate of Binary Black Hole Mergers Inferred from Advanced LIGO Observations Surrounding GW150914. *Astrophys. J.*, 833(1):L1, 2016. doi: 10.3847/2041-8205/833/1/L1.

- [317] B. P. Abbott et al. Observation of Gravitational Waves from a Binary Black Hole Merger. *Phys. Rev. Lett.*, 116(6):061102, 2016. doi: 10.1103/PhysRevLett.116.061102.
- [318] B. P. Abbott et al. Properties of the Binary Black Hole Merger GW150914. *Phys. Rev. Lett.*, 116(24):241102, 2016. doi: 10.1103/PhysRevLett.116.241102.
- [319] B. P. Abbott et al. Tests of general relativity with GW150914. *Phys. Rev. Lett.*, 116(22):221101, 2016. doi: 10.1103/PhysRevLett.116.221101.
- [320] B. P. Abbott et al. GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence. *Phys. Rev. Lett.*, 116(24):241103, 2016. doi: 10.1103/PhysRevLett.116.241103.
- [321] A. Albert et al. Search for High-energy Neutrinos from Gravitational Wave Event GW151226 and Candidate LVT151012 with ANTARES and IceCube. *Phys. Rev.*, D96(2):022005, 2017. doi: 10.1103/PhysRevD.96.022005.
- [322] Pau Amaro Seoane et al. The Gravitational Universe. arXiv:1305.5720, 2013.
- [323] Chiara Caprini et al. Science with the space-based interferometer eLISA. II: Gravitational waves from cosmological phase transitions. *JCAP*, 1604(04):001, 2016. doi: 10.1088/1475-7516/2016/04/001.
- [324] Eldad Gildener. Gauge Symmetry Hierarchies. *Phys. Rev.*, D14:1667, 1976. doi: 10.1103/PhysRevD.14.1667.
- [325] Steven Weinberg. Gauge Hierarchies. *Phys. Lett.*, B82:387–391, 1979. doi: 10.1016/0370-2693(79)90248-X.
- [326] Gerard 't Hooft. Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking. *NATO Sci. Ser. B*, 59:135–157, 1980. doi: 10.1007/978-1-4684-7571-5_9.
- [327] Eldad Gildener. Gauge Symmetry Hierarchies Revisited. *Phys. Lett.*, B92:111–114, 1980. doi: 10.1016/0370-2693(80)90316-0.
- [328] C. Wetterich. Fine Tuning Problem and the Renormalization Group. *Phys. Lett.*, B140:215–222, 1984. doi: 10.1016/0370-2693(84)90923-7.
- [329] M. J. G. Veltman. The Infrared - Ultraviolet Connection. *Acta Phys. Polon.*, B12:437, 1981.
- [330] William A. Bardeen. On naturalness in the standard model. In *Ontake Summer Institute on Particle Physics Ontake Mountain, Japan, August 27-September 2, 1995*, 1995.
- [331] Hajime Aoki and Satoshi Iso. Revisiting the Naturalness Problem – Who is afraid of quadratic divergences? *Phys. Rev.*, D86:013001, 2012. doi: 10.1103/PhysRevD.86.013001.

- [332] Stephen P. Martin. A Supersymmetry primer. *Adv.Ser.Direct.High Energy Phys.*, 21: 1–153, 2010. doi: 10.1142/9789814307505_0001.
- [333] Krzysztof A. Meissner and Hermann Nicolai. Conformal Symmetry and the Standard Model. *Phys. Lett.*, B648:312–317, 2007. doi: 10.1016/j.physletb.2007.03.023.
- [334] Y. Fujii. Scalar-tensor theory of gravitation and spontaneous breakdown of scale invariance. *Phys. Rev.*, D9:874–876, 1974. doi: 10.1103/PhysRevD.9.874.
- [335] F. Englert, C. Truffin, and R. Gastmans. Conformal Invariance in Quantum Gravity. *Nucl. Phys.*, B117:407–432, 1976. doi: 10.1016/0550-3213(76)90406-5.
- [336] W. Buchmuller and N. Dragon. Scale Invariance and Spontaneous Symmetry Breaking. *Phys. Lett.*, B195:417–422, 1987. doi: 10.1016/0370-2693(87)90041-4.
- [337] C. Wetterich. Cosmology and the Fate of Dilatation Symmetry. *Nucl. Phys.*, B302: 668–696, 1988. doi: 10.1016/0550-3213(88)90193-9.
- [338] W. Buchmuller and N. Dragon. Dilatons in Flat and Curved Space-time. *Nucl. Phys.*, B321:207–231, 1989. doi: 10.1016/0550-3213(89)90249-6.
- [339] Robert Foot, Archil Kobakhidze, and Raymond R. Volkas. Electroweak Higgs as a pseudo-Goldstone boson of broken scale invariance. *Phys. Lett.*, B655:156–161, 2007. doi: 10.1016/j.physletb.2007.06.084.
- [340] Robert Foot, Archil Kobakhidze, Kristian L. McDonald, and Raymond R. Volkas. Neutrino mass in radiatively-broken scale-invariant models. *Phys. Rev.*, D76:075014, 2007. doi: 10.1103/PhysRevD.76.075014.
- [341] Robert Foot, Archil Kobakhidze, Kristian L. McDonald, and Raymond R. Volkas. A Solution to the hierarchy problem from an almost decoupled hidden sector within a classically scale invariant theory. *Phys. Rev.*, D77:035006, 2008. doi: 10.1103/PhysRevD.77.035006.
- [342] Krzysztof A. Meissner and Hermann Nicolai. Effective action, conformal anomaly and the issue of quadratic divergences. *Phys. Lett.*, B660:260–266, 2008. doi: 10.1016/j.physletb.2007.12.035.
- [343] Mikhail Shaposhnikov and Daniel Zenhausern. Scale invariance, unimodular gravity and dark energy. *Phys. Lett.*, B671:187–192, 2009. doi: 10.1016/j.physletb.2008.11.054.
- [344] Mikhail Shaposhnikov and Daniel Zenhausern. Quantum scale invariance, cosmological constant and hierarchy problem. *Phys. Lett.*, B671:162–166, 2009. doi: 10.1016/j.physletb.2008.11.041.
- [345] M. E. Shaposhnikov and F. V. Tkachov. Quantum scale-invariant models as effective field theories. arXiv:0905.4857, 2009.

- [346] Robert Foot, Archil Kobakhidze, and Raymond R. Volkas. Stable mass hierarchies and dark matter from hidden sectors in the scale-invariant standard model. *Phys. Rev.*, D82:035005, 2010. doi: 10.1103/PhysRevD.82.035005.
- [347] Robert Foot, Archil Kobakhidze, and Raymond R. Volkas. Cosmological constant in scale-invariant theories. *Phys. Rev.*, D84:075010, 2011. doi: 10.1103/PhysRevD.84.075010.
- [348] Robert Foot and Archil Kobakhidze. Electroweak Scale Invariant Models with Small Cosmological Constant. *Int. J. Mod. Phys.*, A30(21):1550126, 2015. doi: 10.1142/S0217751X15501262.
- [349] Diego Blas, Mikhail Shaposhnikov, and Daniel Zenhausern. Scale-invariant alternatives to general relativity. *Phys. Rev.*, D84:044001, 2011. doi: 10.1103/PhysRevD.84.044001.
- [350] Ichiro Oda. Classically Scale-invariant B-L Model and Conformal Gravity. *Phys. Lett.*, B724:160–164, 2013. doi: 10.1016/j.physletb.2013.06.014.
- [351] Roberta Armillis, Alexander Monin, and Mikhail Shaposhnikov. Spontaneously Broken Conformal Symmetry: Dealing with the Trace Anomaly. *JHEP*, 10:030, 2013. doi: 10.1007/JHEP10(2013)030.
- [352] Itzhak Bars, Paul Steinhardt, and Neil Turok. Local Conformal Symmetry in Physics and Cosmology. *Phys. Rev.*, D89(4):043515, 2014. doi: 10.1103/PhysRevD.89.043515.
- [353] Carlos Tamarit. Running couplings with a vanishing scale anomaly. *JHEP*, 12:098, 2013. doi: 10.1007/JHEP12(2013)098.
- [354] Csaba Csaki, Nemanja Kaloper, Javi Serra, and John Terning. Inflation from Broken Scale Invariance. *Phys.Rev.Lett.*, 113:161302, 2014. doi: 10.1103/PhysRevLett.113.161302.
- [355] Rutger H. Boels and Wadim Wormsbecher. Spontaneously broken conformal invariance in observables. arXiv:1507.08162, 2015.
- [356] Pedro G. Ferreira, Christopher T. Hill, and Graham G. Ross. Scale-Independent Inflation and Hierarchy Generation. *Phys. Lett.*, B763:174–178, 2016. doi: 10.1016/j.physletb.2016.10.036.
- [357] Georgios K. Karananas and Mikhail Shaposhnikov. Scale invariant alternatives to general relativity. II. Dilaton properties. *Phys. Rev.*, D93(8):084052, 2016. doi: 10.1103/PhysRevD.93.084052.
- [358] K. Kannike, M. Raidal, C. Spethmann, and H. Veermäe. The evolving Planck mass in classically scale-invariant theories. *JHEP*, 04:026, 2017. doi: 10.1007/JHEP04(2017)026.

- [359] Georgios K. Karananas and Javier Rubio. On the geometrical interpretation of scale-invariant models of inflation. *Phys. Lett.*, B761:223–228, 2016. doi: 10.1016/j.physletb.2016.08.037.
- [360] Pedro G. Ferreira, Christopher T. Hill, and Graham G. Ross. Weyl Current, Scale-Invariant Inflation and Planck Scale Generation. *Phys. Rev.*, D95(4):043507, 2017. doi: 10.1103/PhysRevD.95.043507.
- [361] D. M. Ghilencea, Z. Lalak, and P. Olszewski. Standard Model with spontaneously broken quantum scale invariance. arXiv:1612.09120, 2016.
- [362] Pedro G. Ferreira, Christopher T. Hill, and Graham G. Ross. No fifth force in a scale invariant universe. *Phys. Rev.*, D95(6):064038, 2017. doi: 10.1103/PhysRevD.95.064038.
- [363] Archil Kobakhidze and Shelley Liang. Standard Model with hidden scale invariance and light dilaton. arXiv:1701.04927, 2017.
- [364] Alberto Salvio. Inflationary Perturbations in No-Scale Theories. *Eur. Phys. J.*, C77(4):267, 2017. doi: 10.1140/epjc/s10052-017-4825-6.
- [365] P. A. R. Ade et al. Planck 2015 results. XX. Constraints on inflation. *Astron. Astrophys.*, 594:A20, 2016. doi: 10.1051/0004-6361/201525898.
- [366] P. J. E. Peebles. Tests of Cosmological Models Constrained by Inflation. *Astrophys. J.*, 284:439–444, 1984. doi: 10.1086/162425.
- [367] Robert H. Brandenberger and Jerome Martin. Trans-Planckian Issues for Inflationary Cosmology. *Class. Quant. Grav.*, 30:113001, 2013. doi: 10.1088/0264-9381/30/11/113001.
- [368] Gabriela Barenboim and Oscar Vives. Transplanckian masses in inflation. *Nucl. Part. Phys. Proc.*, 273-275:446–451, 2016. doi: 10.1016/j.nuclphysbps.2015.09.065.
- [369] Clifford Cheung, Paolo Creminelli, A. Liam Fitzpatrick, Jared Kaplan, and Leonardo Senatore. The Effective Field Theory of Inflation. *JHEP*, 03:014, 2008. doi: 10.1088/1126-6708/2008/03/014.
- [370] Steven Weinberg. Effective Field Theory for Inflation. *Phys. Rev.*, D77:123541, 2008. doi: 10.1103/PhysRevD.77.123541.
- [371] Lev Kofman, Andrei D. Linde, and Alexei A. Starobinsky. Reheating after inflation. *Phys. Rev. Lett.*, 73:3195–3198, 1994. doi: 10.1103/PhysRevLett.73.3195.
- [372] Alan H. Guth and David I. Kaiser. Inflationary cosmology: Exploring the Universe from the smallest to the largest scales. *Science*, 307:884–890, 2005. doi: 10.1126/science.1107483.

- [373] Sean M. Carroll. *Spacetime and geometry: An introduction to general relativity*. San Francisco, USA: Addison-Wesley, 2004. ISBN 0805387323, 9780805387322.
- [374] Alexei A. Starobinsky. Multicomponent de Sitter (Inflationary) Stages and the Generation of Perturbations. *JETP Lett.*, 42:152–155, 1985.
- [375] L.A. Kofman and D. Yu. Pogosian. Nonflat Perturbations in Inflationary Cosmology. *Phys.Lett.*, B214:508–514, 1988. doi: 10.1016/0370-2693(88)90109-8.
- [376] Juan Garcia-Bellido and David Wands. Metric perturbations in two field inflation. *Phys. Rev.*, D53:5437–5445, 1996. doi: 10.1103/PhysRevD.53.5437.
- [377] Laura E. Allen, Sujata Gupta, and David Wands. Non-gaussian perturbations from multi-field inflation. *JCAP*, 0601:006, 2006. doi: 10.1088/1475-7516/2006/01/006.
- [378] Filippo Vernizzi and David Wands. Non-gaussianities in two-field inflation. *JCAP*, 0605:019, 2006. doi: 10.1088/1475-7516/2006/05/019.
- [379] Thorsten Battefeld and Richard Easther. Non-Gaussianities in Multi-field Inflation. *JCAP*, 0703:020, 2007. doi: 10.1088/1475-7516/2007/03/020.
- [380] Shuichiro Yokoyama, Teruaki Suyama, and Takahiro Tanaka. Primordial Non-Gaussianity in Multi-Scalar Inflation. *Phys. Rev.*, D77:083511, 2008. doi: 10.1103/PhysRevD.77.083511.
- [381] Shuichiro Yokoyama, Teruaki Suyama, and Takahiro Tanaka. Primordial Non-Gaussianity in Multi-Scalar Slow-Roll Inflation. *JCAP*, 0707:013, 2007. doi: 10.1088/1475-7516/2007/07/013.
- [382] David Wands. Multiple field inflation. *Lect.Notes Phys.*, 738:275–304, 2008. doi: 10.1007/978-3-540-74353-8_8.
- [383] David Langlois. Cosmological perturbations from multi-field inflation. *J.Phys.Conf.Ser.*, 140:012004, 2008. doi: 10.1088/1742-6596/140/1/012004.
- [384] Ki-Young Choi, Jinn-Ouk Gong, and Donghui Jeong. Evolution of the curvature perturbation during and after multi-field inflation. *JCAP*, 0902:032, 2009. doi: 10.1088/1475-7516/2009/02/032.
- [385] Takeshi Chiba and Masahide Yamaguchi. Extended Slow-Roll Conditions and Primordial Fluctuations: Multiple Scalar Fields and Generalized Gravity. *JCAP*, 0901:019, 2009. doi: 10.1088/1475-7516/2009/01/019.
- [386] David I. Kaiser. Conformal Transformations with Multiple Scalar Fields. *Phys.Rev.*, D81:084044, 2010. doi: 10.1103/PhysRevD.81.084044.

- [387] Leonardo Senatore and Matias Zaldarriaga. The Effective Field Theory of Multifield Inflation. *JHEP*, 04:024, 2012. doi: 10.1007/JHEP04(2012)024.
- [388] Peter Adshead and Mark Wyman. Chromo-Natural Inflation: Natural inflation on a steep potential with classical non-Abelian gauge fields. *Phys.Rev.Lett.*, 108:261302, 2012. doi: 10.1103/PhysRevLett.108.261302.
- [389] M.M. Sheikh-Jabbari. Gauge-flation Vs Chromo-Natural Inflation. *Phys.Lett.*, B717:6–9, 2012. doi: 10.1016/j.physletb.2012.09.014.
- [390] Peter Adshead, Emil Martinec, and Mark Wyman. Perturbations in Chromo-Natural Inflation. *JHEP*, 1309:087, 2013. doi: 10.1007/JHEP09(2013)087.
- [391] Mark P. Hertzberg and Johanna Karouby. Generating the Observed Baryon Asymmetry from the Inflaton Field. *Phys.Rev.*, D89(6):063523, 2014. doi: 10.1103/PhysRevD.89.063523.
- [392] Robert Foot, Archil Kobakhidze, Kristian L. McDonald, and Raymond R. Volkas. Poincaré protection for a natural electroweak scale. *Phys. Rev.*, D89(11):115018, 2014. doi: 10.1103/PhysRevD.89.115018.
- [393] Archil Kobakhidze and Kristian L. McDonald. Comments on the Hierarchy Problem in Effective Theories. *JHEP*, 07:155, 2014. doi: 10.1007/JHEP07(2014)155.
- [394] Juan Garcia-Bellido, Javier Rubio, Mikhail Shaposhnikov, and Daniel Zenhausern. Higgs-Dilaton Cosmology: From the Early to the Late Universe. *Phys. Rev.*, D84:123504, 2011. doi: 10.1103/PhysRevD.84.123504.
- [395] Renata Kallosh and Andrei Linde. Universality Class in Conformal Inflation. *JCAP*, 1307:002, 2013. doi: 10.1088/1475-7516/2013/07/002.
- [396] Alberto Salvio and Alessandro Strumia. Agravity. *JHEP*, 06:080, 2014. doi: 10.1007/JHEP06(2014)080.
- [397] John Ellis, Marcos A. G. Garcia, Dimitri V. Nanopoulos, and Keith A. Olive. A No-Scale Inflationary Model to Fit Them All. *JCAP*, 1408:044, 2014. doi: 10.1088/1475-7516/2014/08/044.
- [398] Kristjan Kannike, Antonio Racioppi, and Martti Raidal. Embedding inflation into the Standard Model - more evidence for classical scale invariance. *JHEP*, 06:154, 2014. doi: 10.1007/JHEP06(2014)154.
- [399] Kristjan Kannike, Gert Hütsi, Liberato Pizza, Antonio Racioppi, Martti Raidal, Alberto Salvio, and Alessandro Strumia. Dynamically Induced Planck Scale and Inflation. *JHEP*, 05:065, 2015. doi: 10.1007/JHEP05(2015)065.

- [400] Mehmet Ozkan and Diederik Roest. Universality Classes of Scale Invariant Inflation. arXiv:1507.03603, 2015.
- [401] Kristjan Kannike, G. Hütsi, L. Pizza, A. Racioppi, M. Raidal, A. Salvio, and Alessandro Strumia. Dynamically Induced Planck Scale and Inflation. *PoS*, EPS-HEP2015:379, 2015.
- [402] Arsham Farzinnia and Seyen Kouwn. Classically scale invariant inflation, supermassive WIMPs, and adimensional gravity. *Phys. Rev.*, D93(6):063528, 2016. doi: 10.1103/PhysRevD.93.063528.
- [403] Massimiliano Rinaldi and Luciano Vanzo. Inflation and reheating in theories with spontaneous scale invariance symmetry breaking. *Phys. Rev.*, D94(2):024009, 2016. doi: 10.1103/PhysRevD.94.024009.
- [404] S. Sonego and V. Faraoni. Coupling to the curvature for a scalar field from the equivalence principle. *Class. Quant. Grav.*, 10:1185–1187, 1993. doi: 10.1088/0264-9381/10/6/015.
- [405] A. A. Grib and E. A. Poberii. On the difference between conformal and minimal couplings in general relativity. *Helv. Phys. Acta*, 68:380–395, 1995.
- [406] Valerio Faraoni. Nonminimal coupling of the scalar field and inflation. *Phys. Rev.*, D53:6813–6821, 1996. doi: 10.1103/PhysRevD.53.6813.
- [407] Mark P. Hertzberg. On Inflation with Non-minimal Coupling. *JHEP*, 11:023, 2010. doi: 10.1007/JHEP11(2010)023.
- [408] Mauro Pieroni. β -function formalism for inflationary models with a non minimal coupling with gravity. *JCAP*, 1602(02):012, 2016. doi: 10.1088/1475-7516/2016/02/012.
- [409] Michal Artymowski and Antonio Racioppi. Scalar-tensor linear inflation. *JCAP*, 1704(04):007, 2017. doi: 10.1088/1475-7516/2017/04/007.
- [410] Orest Hrycyna. What ξ ? Cosmological constraints on the non-minimal coupling constant. *Phys. Lett.*, B768:218–227, 2017. doi: 10.1016/j.physletb.2017.02.062.
- [411] Fedor L. Bezrukov and Mikhail Shaposhnikov. The Standard Model Higgs boson as the inflaton. *Phys. Lett.*, B659:703–706, 2008. doi: 10.1016/j.physletb.2007.11.072.
- [412] Alexander Yu. Kamenshchik and Christian F. Steinwachs. Question of quantum equivalence between Jordan frame and Einstein frame. *Phys. Rev.*, D91(8):084033, 2015. doi: 10.1103/PhysRevD.91.084033.
- [413] Nobuchika Okada, Mansoor Ur Rehman, and Qaisar Shafi. Tensor to Scalar Ratio in Non-Minimal ϕ^4 Inflation. *Phys. Rev.*, D82:043502, 2010. doi: 10.1103/PhysRevD.82.043502.

- [414] S. D. Odintsov. Two loop effective potential in quantum field theory in curved space-time. *Phys. Lett.*, B306:233–236, 1993. doi: 10.1016/0370-2693(93)90073-Q.
- [415] Kristjan Kannike, Antonio Racioppi, and Martti Raidal. Linear inflation from quartic potential. *JHEP*, 01:035, 2016. doi: 10.1007/JHEP01(2016)035.
- [416] Qing-Guo Huang, Ke Wang, and Sai Wang. Inflation model constraints from data released in 2015. *Phys. Rev.*, D93(10):103516, 2016. doi: 10.1103/PhysRevD.93.103516.
- [417] Gordan Krnjaic. Can the Baryon Asymmetry Arise From Initial Conditions? arXiv:1606.05344, 2016.
- [418] Koichi Funakubo. CP violation and baryogenesis at the electroweak phase transition. *Prog. Theor. Phys.*, 96:475–520, 1996. doi: 10.1143/PTP.96.475.
- [419] Wilfried Buchmuller. Baryogenesis: 40 Years Later. In *Proceedings on 13th International Symposium on Particles, strings, and cosmology (PASCOS 2007): London, UK, July 2-7, 2007*, 2007.
- [420] Steven Weinberg. *Cosmology*. Oxford, UK: Oxford Univ. Pr. (2008) 593 p, 2008. ISBN 9780198526827.
- [421] Mikhail Shaposhnikov. Baryogenesis. *J. Phys. Conf. Ser.*, 171:012005, 2009. doi: 10.1088/1742-6596/171/1/012005.
- [422] Pavel Fileviez Perez. New Paradigm for Baryon and Lepton Number Violation. *Phys. Rept.*, 597:1–30, 2015. doi: 10.1016/j.physrep.2015.09.001.
- [423] A. D. Dolgov and Andrei D. Linde. Baryon Asymmetry in Inflationary Universe. *Phys. Lett.*, B116:329, 1982. doi: 10.1016/0370-2693(82)90292-1.
- [424] Steven Weinberg. Baryon and Lepton Nonconserving Processes. *Phys.Rev.Lett.*, 43:1566–1570, 1979. doi: 10.1103/PhysRevLett.43.1566.
- [425] Pran Nath and Pavel Fileviez Perez. Proton stability in grand unified theories, in strings and in branes. *Phys.Rept.*, 441:191–317, 2007. doi: 10.1016/j.physrep.2007.02.010.
- [426] Ilja Dorsner and Pavel Fileviez Perez. How long could we live? *Phys.Lett.*, B625:88–95, 2005. doi: 10.1016/j.physletb.2005.08.039.
- [427] Kalliopi Petraki and Raymond R. Volkas. Review of asymmetric dark matter. *Int.J.Mod.Phys.*, A28:1330028, 2013. doi: 10.1142/S0217751X13300287.
- [428] T. P. Cheng and L. F. Li. *Gauge Theory of Elementary Particle Physics*. Oxford, Uk: Clarendon (1984) 536 P. (Oxford Science Publications), 1984. ISBN 9780198519614.

- [429] Gerald V. Dunne. Aspects of Chern-Simons theory. In *Topological Aspects of Low-dimensional Systems: Proceedings, Les Houches Summer School of Theoretical Physics, Session 69: Les Houches, France, July 7-31 1998*, 1998.
- [430] V.A. Rubakov and M.E. Shaposhnikov. Electroweak baryon number nonconservation in the early universe and in high-energy collisions. *Usp.Fiz.Nauk*, 166:493–537, 1996. doi: 10.1070/PU1996v039n05ABEH000145.
- [431] Erick J. Weinberg. Classical solutions in quantum field theories. *Ann.Rev.Nucl.Part.Sci.*, 42:177–210, 1992.
- [432] Peter Brockway Arnold and Larry D. McLerran. Sphalerons, Small Fluctuations and Baryon Number Violation in Electroweak Theory. *Phys.Rev.*, D36:581, 1987. doi: 10.1103/PhysRevD.36.581.
- [433] Peter Brockway Arnold. An Introduction to baryon violation in standard electroweak theory. In *Testing the Standard Model - TASI-90: Theoretical Advanced Study Inst. in Elementary Particle Physics Boulder, Colorado, June 3-29, 1990*, pages 0719–742, 1990.
- [434] Frans R. Klinkhamer and N.S. Manton. A Saddle Point Solution in the Weinberg-Salam Theory. *Phys.Rev.*, D30:2212, 1984. doi: 10.1103/PhysRevD.30.2212.
- [435] Dmitri Diakonov. Instantons at work. *Prog.Part.Nucl.Phys.*, 51:173–222, 2003. doi: 10.1016/S0146-6410(03)90014-7.
- [436] Olivier Espinosa. High-Energy Behavior of Baryon and Lepton Number Violating Scattering Amplitudes and Breakdown of Unitarity in the Standard Model. *Nucl.Phys.*, B343: 310–340, 1990. doi: 10.1016/0550-3213(90)90473-Q.
- [437] M. Hellmund and J. Kripfganz. The Decay of the sphalerons. *Nucl.Phys.*, B373:749–760, 1992. doi: 10.1016/0550-3213(92)90274-F.
- [438] Y. Burnier, M. Laine, and M. Shaposhnikov. Baryon and lepton number violation rates across the electroweak crossover. *JCAP*, 0602:007, 2006. doi: 10.1088/1475-7516/2006/02/007.
- [439] Luis Bento. Sphaleron relaxation temperatures. *JCAP*, 0311:002, 2003. doi: 10.1088/1475-7516/2003/11/002.
- [440] S. Yu. Khlebnikov and M.E. Shaposhnikov. The Statistical Theory of Anomalous Fermion Number Nonconservation. *Nucl.Phys.*, B308:885–912, 1988. doi: 10.1016/0550-3213(88)90133-2.
- [441] Azadeh Maleknejad, Mahdiyar Noorbala, and M. M. Sheikh-Jabbari. Inflato-Natural Leptogenesis: Leptogenesis in Chromo-Natural and Gauge Inflatons. arXiv:1208.2807, 2012.

- [442] Azadeh Maleknejad. Chiral Gravity Waves and Leptogenesis in Inflationary Models with non-Abelian Gauge Fields. *Phys. Rev.*, D90(2):023542, 2014. doi: 10.1103/PhysRevD.90.023542.
- [443] Masahiro Ibe, Shigeki Matsumoto, and Tsutomu T. Yanagida. The GeV-scale dark matter with B–L asymmetry. *Phys. Lett.*, B708:112–118, 2012. doi: 10.1016/j.physletb.2012.01.032.
- [444] S. M. Barr. The Unification and Cogeneration of Dark Matter and Baryonic Matter. *Phys. Rev.*, D85:013001, 2012. doi: 10.1103/PhysRevD.85.013001.
- [445] Shinya Kanemura, Toshinori Matsui, and Hiroaki Sugiyama. Neutrino mass and dark matter from gauged $U(1)_{B-L}$ breaking. *Phys. Rev.*, D90:013001, 2014. doi: 10.1103/PhysRevD.90.013001.
- [446] Zackaria Chacko, Yanou Cui, Sungwoo Hong, and Takemichi Okui. Hidden dark matter sector, dark radiation, and the CMB. *Phys. Rev.*, D92:055033, 2015. doi: 10.1103/PhysRevD.92.055033.
- [447] Alexandre Alves, Asher Berlin, Stefano Profumo, and Farinaldo S. Queiroz. Dirac-fermionic dark matter in $U(1)_X$ models. *JHEP*, 10:076, 2015. doi: 10.1007/JHEP10(2015)076.
- [448] Nobuchika Okada and Satomi Okada. Z' -portal right-handed neutrino dark matter in the minimal $U(1)_X$ extended Standard Model. *Phys. Rev.*, D95(3):035025, 2017. doi: 10.1103/PhysRevD.95.035025.
- [449] Wan-Zhe Feng and Pran Nath. Baryogenesis and Dark Matter in $U(1)$ Extensions. *Mod. Phys. Lett.*, A32:1740005, 2017. doi: 10.1142/S0217732317400053.
- [450] Bruce A. Campbell, Sacha Davidson, John R. Ellis, and Keith A. Olive. On the baryon, lepton flavor and right-handed electron asymmetries of the universe. *Phys. Lett.*, B297:118–124, 1992. doi: 10.1016/0370-2693(92)91079-O.
- [451] Massimo Giovannini and M.E. Shaposhnikov. Primordial hypermagnetic fields and triangle anomaly. *Phys.Rev.*, D57:2186–2206, 1998. doi: 10.1103/PhysRevD.57.2186.
- [452] Massimo Giovannini. Hypermagnetic knots, Chern-Simons waves and the baryon asymmetry. *Phys.Rev.*, D61:063502, 2000. doi: 10.1103/PhysRevD.61.063502.
- [453] Robert Foot, Girish C. Joshi, and H. Lew. Gauged Baryon and Lepton Numbers. *Phys.Rev.*, D40:2487–2489, 1989. doi: 10.1103/PhysRevD.40.2487.
- [454] Christopher D. Carone and Hitoshi Murayama. Possible light $U(1)$ gauge boson coupled to baryon number. *Phys.Rev.Lett.*, 74:3122–3125, 1995. doi: 10.1103/PhysRevLett.74.3122.

- [455] Christopher D. Carone and Hitoshi Murayama. Realistic models with a light U(1) gauge boson coupled to baryon number. *Phys.Rev.*, D52:484–493, 1995. doi: 10.1103/PhysRevD.52.484.
- [456] Timothy R. Dulaney, Pavel Fileviez Perez, and Mark B. Wise. Dark Matter, Baryon Asymmetry, and Spontaneous B and L Breaking. *Phys.Rev.*, D83:023520, 2011. doi: 10.1103/PhysRevD.83.023520.
- [457] Pavel Fileviez Perez and Mark B. Wise. Baryon and lepton number as local gauge symmetries. *Phys.Rev.*, D82:011901, 2010. doi: 10.1103/PhysRevD.82.079901,10.1103/PhysRevD.82.011901.
- [458] Pavel Fileviez Perez and Mark B. Wise. Low Energy Supersymmetry with Baryon and Lepton Number Gauged. *Phys.Rev.*, D84:055015, 2011. doi: 10.1103/PhysRevD.84.055015.
- [459] Pavel Fileviez Perez and Mark B. Wise. Breaking Local Baryon and Lepton Number at the TeV Scale. *JHEP*, 1108:068, 2011. doi: 10.1007/JHEP08(2011)068.
- [460] Pedro Schwaller, Tim M. P. Tait, and Roberto Vega-Morales. Dark Matter and Vectorlike Leptons from Gauged Lepton Number. *Phys. Rev.*, D88(3):035001, 2013. doi: 10.1103/PhysRevD.88.035001.
- [461] L. Parker. Particle creation in expanding universes. *Phys.Rev.Lett.*, 21:562–564, 1968. doi: 10.1103/PhysRevLett.21.562.
- [462] Leonard Parker. Quantized fields and particle creation in expanding universes. 1. *Phys.Rev.*, 183:1057–1068, 1969. doi: 10.1103/PhysRev.183.1057.
- [463] Oystein Elgaroy, Steen Hannestad, and Troels Haugboelle. Observational constraints on particle production during inflation. *JCAP*, 0309:008, 2003. doi: 10.1088/1475-7516/2003/09/008.
- [464] Neil Barnaby and Zhiqi Huang. Particle Production During Inflation: Observational Constraints and Signatures. *Phys.Rev.*, D80:126018, 2009. doi: 10.1103/PhysRevD.80.126018.
- [465] David H. Lyth. The curvature perturbation in a box. *JCAP*, 0712:016, 2007. doi: 10.1088/1475-7516/2007/12/016.
- [466] Marcela Carena, Alejandro Daleo, Bogdan A. Dobrescu, and Timothy M.P. Tait. Z' gauge bosons at the Tevatron. *Phys.Rev.*, D70:093009, 2004. doi: 10.1103/PhysRevD.70.093009.
- [467] Jonathan L. Rosner. Prominent decay modes of a leptophobic Z' . *Phys.Lett.*, B387:113–117, 1996. doi: 10.1016/0370-2693(96)01022-2.

- [468] Howard Georgi and Sheldon L. Glashow. Decays of a leptophobic gauge boson. *Phys.Lett.*, B387:341–345, 1996. doi: 10.1016/0370-2693(96)00997-5.
- [469] Elena Accomando, Alexander Belyaev, Luca Fedeli, Stephen F. King, and Claire Shepherd-Themistocleous. Z' physics with early LHC data. *Phys.Rev.*, D83:075012, 2011. doi: 10.1103/PhysRevD.83.075012.
- [470] Patrick J. Fox, Jia Liu, David Tucker-Smith, and Neal Weiner. An Effective Z' . *Phys.Rev.*, D84:115006, 2011. doi: 10.1103/PhysRevD.84.115006.
- [471] Bogdan A. Dobrescu and Felix Yu. Coupling-mass mapping of dijet peak searches. *Phys.Rev.*, D88(3):035021, 2013. doi: 10.1103/PhysRevD.88.035021,10.1103/PhysRevD.90.079901.
- [472] Pavel Fileviez Perez, Sebastian Ohmer, and Hiren H. Patel. Minimal Theory for Lepto-Baryons. *Phys.Lett.*, B735:283–287, 2014. doi: 10.1016/j.physletb.2014.06.057.
- [473] G. Aad et al. Search for new phenomena in the dijet mass distribution using $p - p$ collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys.Rev.*, D91(5):052007, 2015. doi: 10.1103/PhysRevD.91.052007.
- [474] Georges Aad et al. Search for high-mass dilepton resonances in pp collisions at 8 TeV with the ATLAS detector. *Phys.Rev.*, D90(5):052005, 2014. doi: 10.1103/PhysRevD.90.052005.
- [475] Julian Heeck. Unbroken B-L Symmetry. *Phys.Lett.*, B739:256–262, 2014. doi: 10.1016/j.physletb.2014.10.067.
- [476] Bogdan A. Dobrescu. Leptophobic Boson Signals with Leptons, Jets and Missing Energy. arXiv:1506.04435, 2015.
- [477] V. Khachatryan et al. Search for resonances and quantum black holes using dijet mass spectra in proton-proton collisions at $\sqrt{s} = 8$ TeV. *Phys.Rev.*, D91(5):052009, 2015. doi: 10.1103/PhysRevD.91.052009.
- [478] S. Rostam Zadeh and S. S. Gousheh. Effects of the $U_Y(1)$ Chern-Simons term and its baryonic contribution on matter asymmetries and hypermagnetic fields. *Phys. Rev.*, D95(5):056001, 2017. doi: 10.1103/PhysRevD.95.056001.
- [479] Andreas Albrecht, Paul J. Steinhardt, Michael S. Turner, and Frank Wilczek. Reheating an Inflationary Universe. *Phys. Rev. Lett.*, 48:1437, 1982. doi: 10.1103/PhysRevLett.48.1437.
- [480] Y. Shtanov, Jennie H. Traschen, and Robert H. Brandenberger. Universe reheating after inflation. *Phys. Rev.*, D51:5438–5455, 1995. doi: 10.1103/PhysRevD.51.5438.

- [481] Alexandre Dolgov and Katherine Freese. Calculation of particle production by Nambu Goldstone bosons with application to inflation, reheating and baryogenesis. *Phys. Rev.*, D51:2693–2702, 1995. doi: 10.1103/PhysRevD.51.2693.
- [482] Lev A. Kofman. The Origin of matter in the universe: Reheating after inflation. arXiv:9605155, 1996.
- [483] Lev Kofman, Andrei D. Linde, and Alexei A. Starobinsky. Towards the theory of reheating after inflation. *Phys. Rev.*, D56:3258–3295, 1997. doi: 10.1103/PhysRevD.56.3258.
- [484] Daniel J.H. Chung, Edward W. Kolb, and Antonio Riotto. Production of massive particles during reheating. *Phys.Rev.*, D60:063504, 1999. doi: 10.1103/PhysRevD.60.063504.
- [485] Sacha Davidson and Subir Sarkar. Thermalization after inflation. *JHEP*, 11:012, 2000. doi: 10.1088/1126-6708/2000/11/012.
- [486] Edward W. Kolb, Alessio Notari, and Antonio Riotto. On the reheating stage after inflation. *Phys.Rev.*, D68:123505, 2003. doi: 10.1103/PhysRevD.68.123505.
- [487] Bruce A. Bassett, Shinji Tsujikawa, and David Wands. Inflation dynamics and reheating. *Rev. Mod. Phys.*, 78:537–589, 2006. doi: 10.1103/RevModPhys.78.537.
- [488] Rouzbeh Allahverdi, Robert Brandenberger, Francis-Yan Cyr-Racine, and Anupam Mazumdar. Reheating in Inflationary Cosmology: Theory and Applications. *Ann. Rev. Nucl. Part. Sci.*, 60:27–51, 2010. doi: 10.1146/annurev.nucl.012809.104511.
- [489] Fumihiro Takayama. Extremely Long-Lived Charged Massive Particles as A Probe for Reheating of the Universe. *Phys. Rev.*, D77:116003, 2008. doi: 10.1103/PhysRevD.77.116003.
- [490] Jerome Martin and Christophe Ringeval. First CMB Constraints on the Inflationary Reheating Temperature. *Phys. Rev.*, D82:023511, 2010. doi: 10.1103/PhysRevD.82.023511.
- [491] Liang Dai, Marc Kamionkowski, and Junpu Wang. Reheating constraints to inflationary models. *Phys. Rev. Lett.*, 113:041302, 2014. doi: 10.1103/PhysRevLett.113.041302.
- [492] Gian Francesco Giudice, Edward W. Kolb, and Antonio Riotto. Largest temperature of the radiation era and its cosmological implications. *Phys.Rev.*, D64:023508, 2001. doi: 10.1103/PhysRevD.64.023508.
- [493] Alexandre Dolgov, Katherine Freese, Raghavan Rangarajan, and Mark Srednicki. Baryogenesis during reheating in natural inflation and comments on spontaneous baryogenesis. *Phys. Rev.*, D56:6155–6165, 1997. doi: 10.1103/PhysRevD.56.6155.
- [494] G. Magnano, M. Ferraris, and M. Francaviglia. Nonlinear gravitational Lagrangians. *Gen. Rel. Grav.*, 19:465, 1987. doi: 10.1007/BF00760651.

- [495] D. S. Gorbunov and A. A. Tokareva. Inflation and reheating in the Starobinsky model with conformal Higgs Field. *Phys. Part. Nucl. Lett.*, 10:633–636, 2013. doi: 10.1134/S1547477113070030.
- [496] John Ellis, Marcos A. G. Garcia, Dimitri V. Nanopoulos, and Keith A. Olive. Calculations of Inflaton Decays and Reheating: with Applications to No-Scale Inflation Models. *JCAP*, 1507(07):050, 2015. doi: 10.1088/1475-7516/2015/07/050.
- [497] Andrew G. Cohen and David B. Kaplan. Spontaneous baryogenesis. *Nucl. Phys.*, B308:913–928, 1988. doi: 10.1016/0550-3213(88)90134-4.
- [498] M. Yoshimura. Towards resolution of hierarchy problems in a cosmological context. *Phys. Lett.*, B608:183–188, 2005. doi: 10.1016/j.physletb.2005.01.025.
- [499] Kazuharu Bamba and Motohiko Yoshimura. Curvaton scenario in the presence of two dilatons coupled to the scalar curvature. *Prog. Theor. Phys.*, 115:269–308, 2006. doi: 10.1143/PTP.115.269.
- [500] Azusa Minamizaki and Akio Sugamoto. Can the Baryon Number Density and the Cosmological Constant be interrelated? *Phys. Lett.*, B659:656–660, 2008. doi: 10.1016/j.physletb.2007.11.052.
- [501] P. Reimann, R. Bartussek, R. Häußler, and P. Hänggi. Brownian motors driven by temperature oscillations. *Physics Letters A*, 215:26–31, February 1996. doi: 10.1016/0375-9601(96)00222-8.
- [502] Tatsu Takeuchi, Azusa Minamizaki, and Akio Sugamoto. Ratchet Model of Baryogenesis. In *Strong coupling gauge theories in LHC era. Proceedings, International Workshop, SCGT 09, Nagoya, Japan, December 8-11, 2009*, pages 378–384, 2011. doi: 10.1142/9789814329521_0041.
- [503] Andrew G. Cohen and David B. Kaplan. Thermodynamic Generation of the Baryon Asymmetry. *Phys. Lett.*, B199:251–258, 1987. doi: 10.1016/0370-2693(87)91369-4.
- [504] N. F. Pedersen, O. H. Soerensen, B. Dueholm, and J. Mygind. Half-Harmonic Parametric Oscillations in Josephson Junctions. *J. Low Temp. Phys.*, 38:1–23, 1980. doi: 10.1007/BF00115266.
- [505] D. D’Humieres, M. R. Beasley, B. A. Huberman, and A. Libchaber. Chaotic states and routes to chaos in the forced pendulum. *Phys. Rev.*, A26:3483–3496, 1982. doi: 10.1103/PhysRevA.26.3483.
- [506] Jeffrey A. Harvey and Michael S. Turner. Cosmological baryon and lepton number in the presence of electroweak fermion number violation. *Phys. Rev.*, D42:3344–3349, 1990. doi: 10.1103/PhysRevD.42.3344.

- [507] Tsutomu Yanagida. Horizontal Symmetry and Masses of Neutrinos. *Conf. Proc.*, C7902131:95–99, 1979.
- [508] Tsutomu Yanagida. Horizontal Symmetry and Masses of Neutrinos. *Prog. Theor. Phys.*, 64:1103, 1980. doi: 10.1143/PTP.64.1103.
- [509] Pierre Ramond. The Family Group in Grand Unified Theories. In *International Symposium on Fundamentals of Quantum Theory and Quantum Field Theory Palm Coast, Florida, February 25-March 2, 1979*, pages 265–280, 1979.
- [510] M. Fukugita and T. Yanagida. Baryogenesis Without Grand Unification. *Phys. Lett.*, B174:45–47, 1986. doi: 10.1016/0370-2693(86)91126-3.
- [511] T. Moroi, H. Murayama, and Masahiro Yamaguchi. Cosmological constraints on the light stable gravitino. *Phys. Lett.*, B303:289–294, 1993. doi: 10.1016/0370-2693(93)91434-O.
- [512] M. Bolz, A. Brandenburg, and W. Buchmuller. Thermal production of gravitinos. *Nucl. Phys.*, B606:518–544, 2001. doi: 10.1016/S0550-3213(01)00132-8,10.1016/j.nuclphysb.2007.09.020. [Erratum: *Nucl. Phys.*B790,336(2008)].
- [513] Masahiro Kawasaki, Kazunori Kohri, and Takeo Moroi. Big-Bang nucleosynthesis and hadronic decay of long-lived massive particles. *Phys. Rev.*, D71:083502, 2005. doi: 10.1103/PhysRevD.71.083502.
- [514] Matteo Viel, Julien Lesgourgues, Martin G. Haehnelt, Sabino Matarrese, and Antonio Riotto. Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with WMAP and the Lyman-alpha forest. *Phys. Rev.*, D71:063534, 2005. doi: 10.1103/PhysRevD.71.063534.
- [515] Fuminobu Takahashi and Masaki Yamada. Spontaneous Baryogenesis from Asymmetric Inflaton. *Phys. Lett.*, B756:216–220, 2016. doi: 10.1016/j.physletb.2016.03.020.
- [516] Julien Lesgourgues and Sergio Pastor. Cosmological implications of a relic neutrino asymmetry. *Phys. Rev.*, D60:103521, 1999. doi: 10.1103/PhysRevD.60.103521.
- [517] Nicole F. Bell, Raymond R. Volkas, and Yvonne Y. Y. Wong. Relic neutrino asymmetry evolution from first principles. *Phys. Rev.*, D59:113001, 1999. doi: 10.1103/PhysRevD.59.113001.
- [518] Raymond R. Volkas and Yvonne Y. Y. Wong. Further studies on relic neutrino asymmetry generation. 1. The adiabatic Boltzmann limit, nonadiabatic evolution, and the classical harmonic oscillator analog of the quantum kinetic equations. *Phys. Rev.*, D62:093024, 2000. doi: 10.1103/PhysRevD.62.093024.

- [519] Keith S. M. Lee, Raymond R. Volkas, and Yvonne Y. Y. Wong. Further studies on relic neutrino asymmetry generation. 2. A Rigorous treatment of repopulation in the adiabatic limit. *Phys. Rev.*, D62:093025, 2000. doi: 10.1103/PhysRevD.62.093025.
- [520] John March-Russell, Hitoshi Murayama, and Antonio Riotto. The Small observed baryon asymmetry from a large lepton asymmetry. *JHEP*, 11:015, 1999. doi: 10.1088/1126-6708/1999/11/015.
- [521] John McDonald. Naturally large cosmological neutrino asymmetries in the MSSM. *Phys. Rev. Lett.*, 84:4798–4801, 2000. doi: 10.1103/PhysRevLett.84.4798.
- [522] Dominik J. Schwarz and Maik Stuke. Lepton asymmetry and the cosmic QCD transition. *JCAP*, 0911:025, 2009. doi: 10.1088/1475-7516/2009/11/025,10.1088/1475-7516/2010/10/E01. [Erratum: JCAP1010,E01(2010)].
- [523] V. B. Semikoz, D. D. Sokoloff, and J. W. F. Valle. Is the baryon asymmetry of the Universe related to galactic magnetic fields? *Phys. Rev.*, D80:083510, 2009. doi: 10.1103/PhysRevD.80.083510.
- [524] Gabriela Barenboim and Wan-Il Park. A full picture of large lepton number asymmetries of the Universe. *JCAP*, 1704(04):048, 2017. doi: 10.1088/1475-7516/2017/04/048.
- [525] A. Yahil and G. Beaudet. Big-Bang Nucleosynthesis with Nonzero Lepton Numbers. *Astrophys. J.*, 206:26–29, 1976. doi: 10.1086/154352.
- [526] N. Terasawa and K. Sato. Constraints on Baryon and Lepton Number Asymmetries of the Early Universe from Primordial Nucleosynthesis. *Prog. Theor. Phys.*, 72:1262–1265, 1984. doi: 10.1143/PTP.72.1262.
- [527] Nobuo Terasawa and Katsuhiko Sato. Lepton and Baryon Number Asymmetry of the Universe and Primordial Nucleosynthesis. *Prog. Theor. Phys.*, 80:468, 1988. doi: 10.1143/PTP.80.468.
- [528] Kazunori Kohri, M. Kawasaki, and Katsuhiko Sato. Big bang nucleosynthesis and lepton number asymmetry in the universe. *Astrophys. J.*, 490:72–75, 1997. doi: 10.1086/512793.
- [529] William H. Kinney and Antonio Riotto. Measuring the cosmological lepton asymmetry through the CMB anisotropy. *Phys. Rev. Lett.*, 83:3366–3369, 1999. doi: 10.1103/PhysRevLett.83.3366.
- [530] Sergio Pastor and Julien Lesgourgues. Relic neutrino asymmetry, CMB and large scale structure. *Nucl. Phys. Proc. Suppl.*, 81:47–51, 2000. doi: 10.1016/S0920-5632(99)00857-9.
- [531] Julien Lesgourgues, Sergio Pastor, and Simon Prunet. Cosmological measurement of neutrino mass in the presence of leptonic asymmetry. *Phys. Rev.*, D62:023001, 2000. doi: 10.1103/PhysRevD.62.023001.

- [532] Julien Lesgourgues and Marco Peloso. Remarks on the Boomerang results, the baryon density, and the leptonic asymmetry. *Phys. Rev.*, D62:081301, 2000. doi: 10.1103/PhysRevD.62.081301.
- [533] Julien Lesgourgues and Andrew R. Liddle. The Lepton asymmetry: The Last chance for a critical-density cosmology? *Mon. Not. Roy. Astron. Soc.*, 327:1307, 2001. doi: 10.1046/j.1365-8711.2001.04833.x.
- [534] Vimal Simha and Gary Steigman. Constraining The Universal Lepton Asymmetry. *JCAP*, 0808:011, 2008. doi: 10.1088/1475-7516/2008/08/011.
- [535] Gianpiero Mangano, Gennaro Miele, Sergio Pastor, Ofelia Pisanti, and Srdjan Sarikas. Constraining the cosmic radiation density due to lepton number with Big Bang Nucleosynthesis. *JCAP*, 1103:035, 2011. doi: 10.1088/1475-7516/2011/03/035.
- [536] Gianpiero Mangano, Gennaro Miele, Sergio Pastor, Ofelia Pisanti, and Srdjan Sarikas. Updated BBN bounds on the cosmological lepton asymmetry for non-zero θ_{13} . *Phys. Lett.*, B708:1–5, 2012. doi: 10.1016/j.physletb.2012.01.015.
- [537] Emanuele Castorina, Urbano Franca, Massimiliano Lattanzi, Julien Lesgourgues, Gianpiero Mangano, Alessandro Melchiorri, and Sergio Pastor. Cosmological lepton asymmetry with a nonzero mixing angle θ_{13} . *Phys. Rev.*, D86:023517, 2012. doi: 10.1103/PhysRevD.86.023517.
- [538] Isabel M. Oldengott and Dominik J. Schwarz. Improved constraints on lepton asymmetry from the cosmic microwave background. arXiv:1706.01705, 2017.
- [539] A. D. Dolgov, S. H. Hansen, S. Pastor, S. T. Petcov, G. G. Raffelt, and D. V. Semikoz. Cosmological bounds on neutrino degeneracy improved by flavor oscillations. *Nucl. Phys.*, B632:363–382, 2002. doi: 10.1016/S0550-3213(02)00274-2.
- [540] Yvonne Y. Y. Wong. Analytical treatment of neutrino asymmetry equilibration from flavor oscillations in the early universe. *Phys. Rev.*, D66:025015, 2002. doi: 10.1103/PhysRevD.66.025015.
- [541] Kevork N. Abazajian, John F. Beacom, and Nicole F. Bell. Stringent constraints on cosmological neutrino anti-neutrino asymmetries from synchronized flavor transformation. *Phys. Rev.*, D66:013008, 2002. doi: 10.1103/PhysRevD.66.013008.
- [542] Gabriela Barenboim, William H. Kinney, and Wan-Il Park. Resurrection of large lepton number asymmetries from neutrino flavor oscillations. *Phys. Rev.*, D95(4):043506, 2017. doi: 10.1103/PhysRevD.95.043506.
- [543] Jeffrey A. Harvey and Edward W. Kolb. Grand Unified Theories and the Lepton Number of the Universe. *Phys. Rev.*, D24:2090, 1981. doi: 10.1103/PhysRevD.24.2090.

- [544] Paul Langacker, Gino Segre, and Sanjeev Soni. Majorana Neutrinos, Nucleosynthesis, and the Lepton Asymmetry of the Universe. *Phys. Rev.*, D26:3425, 1982. doi: 10.1103/PhysRevD.26.3425.
- [545] Jiang Liu and Gino Segre. Baryon asymmetry of the universe and large lepton asymmetries. *Phys. Lett.*, B338:259–262, 1994. doi: 10.1016/0370-2693(94)91375-7.
- [546] Robert Foot, Mark J. Thomson, and R. R. Volkas. Large neutrino asymmetries from neutrino oscillations. *Phys. Rev.*, D53:R5349–R5353, 1996. doi: 10.1103/PhysRevD.53.R5349.
- [547] Robert Foot and R. R. Volkas. Studies of neutrino asymmetries generated by ordinary sterile neutrino oscillations in the early universe and implications for big bang nucleosynthesis bounds. *Phys. Rev.*, D55:5147–5176, 1997. doi: 10.1103/PhysRevD.55.5147.
- [548] Alberto Casas, Wai Yan Cheng, and Graciela Gelmini. Generation of large lepton asymmetries. *Nucl. Phys.*, B538:297–308, 1999. doi: 10.1016/S0550-3213(98)00606-3.
- [549] Borut Bajc, Antonio Riotto, and Goran Senjanovic. Large lepton number of the universe and the fate of topological defects. *Phys. Rev. Lett.*, 81:1355–1358, 1998. doi: 10.1103/PhysRevLett.81.1355.
- [550] A. D. Dolgov, S. H. Hansen, S. Pastor, and D. V. Semikoz. Neutrino oscillations in the early universe: How large lepton asymmetry can be generated? *Astropart. Phys.*, 14:79–90, 2000. doi: 10.1016/S0927-6505(00)00111-0.
- [551] Antti Sorri. Physical origin of ‘chaoticity’ of neutrino asymmetry. *Phys. Lett.*, B477:201–207, 2000. doi: 10.1016/S0370-2693(00)00203-3.
- [552] R. Buras and D. V. Semikoz. Lepton asymmetry creation in the early universe. *Astropart. Phys.*, 17:245–261, 2002. doi: 10.1016/S0927-6505(01)00155-4.
- [553] R. Buras and D. V. Semikoz. Maximum lepton asymmetry from active sterile neutrino oscillations in the early universe. *Phys. Rev.*, D64:017302, 2001. doi: 10.1103/PhysRevD.64.017302.
- [554] Kimmo Kainulainen and Antti Sorri. Oscillation induced neutrino asymmetry growth in the early universe. *JHEP*, 02:020, 2002. doi: 10.1088/1126-6708/2002/02/020.
- [555] Masahide Yamaguchi. Generation of cosmological large lepton asymmetry from a rolling scalar field. *Phys. Rev.*, D68:063507, 2003. doi: 10.1103/PhysRevD.68.063507.
- [556] Luis Bento and Francisco C. Santos. Neutrino helicity asymmetries in leptogenesis. *Phys. Rev.*, D71:096001, 2005. doi: 10.1103/PhysRevD.71.096001.

- [557] Arthur Lue, Li-Min Wang, and Marc Kamionkowski. Cosmological signature of new parity violating interactions. *Phys. Rev. Lett.*, 83:1506–1509, 1999. doi: 10.1103/PhysRevLett.83.1506.
- [558] M. Novello, Herman J. Mosquera Cuesta, and V. A. De Lorenci. Birefringence of gravitational waves. In *Recent developments in theoretical and experimental general relativity, gravitation and relativistic field theories. Proceedings, 9th Marcel Grossmann Meeting, MG'9, Rome, Italy, July 2-8, 2000. Pts. A-C*, pages 1092–1095, 2000.
- [559] Stephon Alexander and Jerome Martin. Birefringent gravitational waves and the consistency check of inflation. *Phys. Rev.*, D71:063526, 2005. doi: 10.1103/PhysRevD.71.063526.
- [560] Stephon H. S. Alexander. Inflationary Birefringence and Baryogenesis. *Int. J. Mod. Phys.*, D25(11):1640013, 2016. doi: 10.1142/S0218271816400137.
- [561] R. Jackiw and S.Y. Pi. Chern-Simons modification of general relativity. *Phys.Rev.*, D68:104012, 2003. doi: 10.1103/PhysRevD.68.104012.
- [562] Stephon Alexander and Nicolas Yunes. Chern-Simons Modified General Relativity. *Phys.Rept.*, 480:1–55, 2009. doi: 10.1016/j.physrep.2009.07.002.
- [563] Kiyotomo Ichiki, Masahide Yamaguchi, and Jun'Ichi Yokoyama. Lepton asymmetry in the primordial gravitational wave spectrum. *Phys. Rev.*, D75:084017, 2007. doi: 10.1103/PhysRevD.75.084017.
- [564] Parampreet Singh and Banibrata Mukhopadhyay. Gravitationally induced neutrino asymmetry. *Mod. Phys. Lett.*, A18:779–785, 2003. doi: 10.1142/S0217732303009691.
- [565] Ansgar Denner, H. Eck, O. Hahn, and J. Kublbeck. Feynman rules for fermion number violating interactions. *Nucl. Phys.*, B387:467–481, 1992. doi: 10.1016/0550-3213(92)90169-C.
- [566] Daniel Burns and Apostolos Pilaftsis. Matter Quantum Corrections to the Graviton Self-Energy and the Newtonian Potential. *Phys. Rev.*, D91(6):064047, 2015. doi: 10.1103/PhysRevD.91.064047.
- [567] Damiano Anselmi. A Note on the dimensional regularization of the standard model coupled with quantum gravity. *Phys. Lett.*, B596:90–95, 2004. doi: 10.1016/j.physletb.2004.06.089.
- [568] M. Gomes, T. Mariz, J. R. Nascimento, E. Passos, A. Yu. Petrov, and A. J. da Silva. On the ambiguities in the effective action in Lorentz-violating gravity. *Phys. Rev.*, D78:025029, 2008. doi: 10.1103/PhysRevD.78.025029.

- [569] A. N. Redlich and L. C. R. Wijewardhana. Induced Chern-simons Terms at High Temperatures and Finite Densities. *Phys. Rev. Lett.*, 54:970, 1985. doi: 10.1103/PhysRevLett.54.970.
- [570] James G. McCarthy and Andy Wilkins. Induced Chern-Simons terms. *Phys. Rev.*, D58:085007, 1998. doi: 10.1103/PhysRevD.58.085007.
- [571] M. Joyce and Mikhail E. Shaposhnikov. Primordial magnetic fields, right-handed electrons, and the Abelian anomaly. *Phys. Rev. Lett.*, 79:1193–1196, 1997. doi: 10.1103/PhysRevLett.79.1193.
- [572] Roman Jackiw and V. Alan Kostelecky. Radiatively induced Lorentz and CPT violation in electrodynamics. *Phys. Rev. Lett.*, 82:3572–3575, 1999. doi: 10.1103/PhysRevLett.82.3572.
- [573] T. Mariz, J.R. Nascimento, E. Passos, and R.F. Ribeiro. Chern-Simons - like action induced radiatively in general relativity. *Phys.Rev.*, D70:024014, 2004. doi: 10.1103/PhysRevD.70.024014.
- [574] Maxim Dvornikov and Victor B. Semikoz. Instability of magnetic fields in electroweak plasma driven by neutrino asymmetries. *JCAP*, 1405:002, 2014. doi: 10.1088/1475-7516/2014/05/002.
- [575] Yukinao Akamatsu and Naoki Yamamoto. Chiral Plasma Instabilities. *Phys. Rev. Lett.*, 111:052002, 2013. doi: 10.1103/PhysRevLett.111.052002.
- [576] J. C. C. Felipe, A. R. Vieira, A. L. Cherchiglia, A. P. Baêta Scarpelli, and Marcos Sampaio. Arbitrariness in the gravitational Chern-Simons-like term induced radiatively. *Phys. Rev.*, D89(10):105034, 2014. doi: 10.1103/PhysRevD.89.105034.
- [577] Mohamed M. Anber and Eray Sabancilar. Chiral Gravitational Waves from Chiral Fermions. *Phys. Rev.*, D96(2):023501, 2017. doi: 10.1103/PhysRevD.96.023501.
- [578] Tristan L. Smith, Adrienne L. Erickcek, Robert R. Caldwell, and Marc Kamionkowski. The Effects of Chern-Simons gravity on bodies orbiting the Earth. *Phys.Rev.*, D77:024015, 2008. doi: 10.1103/PhysRevD.77.024015.
- [579] Nicolas Yunes and David N. Spergel. Double Binary Pulsar Test of Dynamical Chern-Simons Modified Gravity. *Phys.Rev.*, D80:042004, 2009. doi: 10.1103/PhysRevD.80.042004.
- [580] Yacine Ali-Haïmoud. Revisiting the double-binary-pulsar probe of non-dynamical Chern-Simons gravity. *Phys.Rev.*, D83:124050, 2011. doi: 10.1103/PhysRevD.83.124050.

-
- [581] Sergei Dyda, Eanna E. Flanagan, and Marc Kamionkowski. Vacuum Instability in Chern-Simons Gravity. *Phys.Rev.*, D86:124031, 2012. doi: 10.1103/PhysRevD.86.124031.
- [582] Stephon Alexander, Lee Samuel Finn, and Nicolas Yunes. A Gravitational-wave probe of effective quantum gravity. *Phys. Rev.*, D78:066005, 2008. doi: 10.1103/PhysRevD.78.066005.
- [583] Sean M. Carroll, Mark Hoffman, and Mark Trodden. Can the dark energy equation of state parameter w be less than -1 ? *Phys. Rev.*, D68:023509, 2003. doi: 10.1103/PhysRevD.68.023509.
- [584] James M. Cline, Sangyong Jeon, and Guy D. Moore. The Phantom menaced: Constraints on low-energy effective ghosts. *Phys. Rev.*, D70:043543, 2004. doi: 10.1103/PhysRevD.70.043543.