

Axion misalignment driven to the hilltop

Raymond T. Co,^a Eric Gonzalez^a and Keisuke Harigaya^b

^a*Leinweber Center for Theoretical Physics, Department of Physics, University of Michigan, 450 Church Street, Ann Arbor, Michigan 48104, U.S.A.*

^b*School of Natural Sciences, Institute for Advanced Study, 1 Einstein Drive, Princeton, NJ 08540, U.S.A.*

E-mail: rtco@umich.edu, ericgz@umich.edu, keisukeharigaya@ias.edu

ABSTRACT: The QCD axion serves as a well-motivated dark matter candidate and the misalignment mechanism is known to reproduce the observed abundance with a decay constant $f_a \simeq \mathcal{O}(10^{12})$ GeV for a misalignment angle $\theta_{\text{mis}} \simeq \mathcal{O}(1)$. While $f_a \ll 10^{12}$ GeV is of great experimental interest, the misalignment mechanism requires the axion to be very close to the hilltop, i.e. $\theta_{\text{mis}} \simeq \pi$. This particular choice of θ_{mis} has been understood as fine-tuning the initial condition. We offer a dynamical explanation for $\theta_{\text{mis}} \simeq \pi$ in a class of models. The axion dynamically relaxes to the minimum of the potential by virtue of an enhanced mass in the early universe. This minimum is subsequently converted to a hilltop because the CP phase of the theory shifts by π when one contribution becomes subdominant to another with an opposite sign. We demonstrate explicit and viable examples in supersymmetric models where the higher dimensional Higgs coupling with the inflaton naturally achieves both criteria. Associated phenomenology includes a strikingly sharp prediction of $3 \times 10^9 \text{ GeV} \lesssim f_a \lesssim 10^{10} \text{ GeV}$ and the absence of isocurvature perturbation.

KEYWORDS: Cosmology of Theories beyond the SM, Confinement, Supersymmetric Gauge Theory, Supersymmetric Standard Model

ARXIV EPRINT: [1812.11192](https://arxiv.org/abs/1812.11192)

Contents

1	Introduction	1
2	Axion misalignment & early relaxation	4
3	Dynamical axion misalignment production at the hilltop	5
3.1	Axion mass during inflation	6
3.2	Shifted axion potential	8
4	Cosmological evolution	9
4.1	Minimal models	9
4.2	Extended models	12
5	Conclusion	15

1 Introduction

Measurements of the neutron electric dipole moment indicate an unnaturally small value of the CP-violating QCD θ parameter [1, 2], which is known as the strong CP problem [3]. Shortly after recognizing this discrepancy, the Peccei-Quinn (PQ) mechanism [4, 5] was developed as a resolution; an anomalous $U(1)_{\text{PQ}}$ symmetry is spontaneously broken at a scale f_a and the resulting pseudo Nambu-Goldstone mode, called the axion a , dynamically relaxes $\bar{\theta} = \theta - \langle a/f_a \rangle$ to a vanishing value consistent with experiments. The value of f_a plays a critical role in observables related to the axion so its precise determination, theoretically and experimentally, is crucial.

A model with a weak scale decay constant was initially proposed [6, 7] but immediately ruled out by laboratory searches. Today, supernovae cooling is the most competitive lower bound giving $f_a \gtrsim 10^8$ GeV [8–12]. Since the axion is very light and stable on cosmological time scales, one can imagine a scenario where its relic abundance accounts for the observed dark matter (DM) abundance $\Omega_{\text{DM}} h^2 = 0.12$. A thermal axion relic abundance is too hot and scarce to be consistent with cold DM. Two non-thermal production mechanisms are commonly considered. The relic abundance from the misalignment mechanism [13–15], namely coherent oscillations due to an initial axion field value $\theta_{\text{mis}} f_a$, is

$$\Omega_{\text{mis}} h^2 \simeq 0.12 \theta_{\text{mis}}^2 \mathcal{F}(\theta_{\text{mis}}) \left(\frac{f_a}{5 \times 10^{11} \text{ GeV}} \right)^{7/6}, \quad (1.1)$$

where $\mathcal{F}(\theta_{\text{mis}})$ is the anharmonicity factor. With the natural assumption of $\mathcal{O}(1)$ initial misalignment, $f_a = 10^{11}$ – 10^{12} GeV is compatible with the observed DM abundance. If the

PQ symmetry is broken after inflation and the domain wall number is unity, the abundance of axions emitted from the string-domain wall network is [16–18]

$$\Omega_{\text{string,DW}} h^2 \simeq 0.04 - 0.3 \left(\frac{f_a}{10^{11} \text{ GeV}} \right)^{7/6}. \quad (1.2)$$

The decay constant $f_a \sim 10^{11}$ GeV reproduces the DM abundance.¹

As ongoing axion experiments are about to reach sensitivity required to probe small decay constants of $10^8 \text{ GeV} < f_a < 10^{12} \text{ GeV}$ [22–32], exploring the theoretical landscape pertaining to small f_a is important. Some studies have been successful in allowing small f_a in a natural setting, such as parametric resonance from a PQ symmetry breaking field [33] and decays of quasi-stable domain walls [17, 34–36]. The misalignment mechanism can reproduce the observed DM abundance for $f_a \ll 10^{12}$ GeV if θ_{mis} is taken sufficiently close to π [37–41], where the anharmonicity factor $\mathcal{F}(\theta_{\text{mis}})$ becomes important.

In this study, we propose a scenario which dynamically predicts $\theta_{\text{mis}} \simeq \pi$ and thus small f_a in the context of axion DM from the misalignment mechanism. It is commonly assumed that no misalignment angles are special in the early universe, and $\theta_{\text{mis}} \simeq \pi$ requires a fine-tuned initial condition. This is not the case given two conditions are met: 1) the axion field dynamically relaxes to the minimum of the potential in the early universe and 2) the model possesses a non-trivial prediction between the minima of the axion potential in the early and today’s epochs. We refer to the axion relaxation with the fulfillment of these requirements as Dynamical Axion Misalignment Production (DAMP). We study DAMP by the dynamics of the Higgs fields during inflation. The mechanism follows from suspending the assumption that axion’s late-time dynamics is agnostic to inflationary dynamics. To be concrete, we study the Minimal Supersymmetric Standard Model (MSSM). The Higgs fields H_u and H_d in general couple to the inflaton potential energy via higher dimensional operators, which lead to so-called Hubble induced masses. The Higgs fields can acquire a large field value in the early universe by virtue of the Hubble induced mass. This large field value gives large quark masses, which enhance the confinement scale to Λ'_{QCD} during inflation. Since m_a is proportional to Λ'_{QCD} , this raises the axion mass to allow for earlier relaxation to the minimum. Note that we need to assume the Higgs fields are not charged under PQ symmetry; otherwise, the decay constant will be as high as the Higgs VEV and suppress the axion mass. For context, early studies [42–44] have made use of moduli fields to raise the QCD confinement scale $\Lambda_{\text{QCD}} \rightarrow \Lambda'_{\text{QCD}}$ during inflation. This avoids fine-tuning problems that arise under the assumption of an $\mathcal{O}(1)$ initial misalignment with large values $f_a > 10^{12}$ GeV. Later studies [45, 46] used Higgs fields as the moduli fields and refined the scope of the mechanism to reduce isocurvature perturbations for models with large inflation scales, which comes at the cost of an inability to suppress the axion abundance. This loss of abundance predictability is because no assumptions are made about the evolution of the axion minimum through inflation. In the MSSM for example,

¹The abundance is estimated assuming a scaling law, with the uncertainty given by that of the spectrum of axions. Recent studies suggest that the number of strings per horizon may increase logarithmically in time [19–21]. If this is actually the case, the abundance may be larger than eq. (1.2).

we have [44]

$$\theta_{\text{eff}} = \theta_{\text{QCD}} + \arg(\det \lambda_u \lambda_d) + 3 \arg(m_{\tilde{g}}) + 3 \arg(B\mu), \quad (1.3)$$

where λ_u, λ_d are the Yukawa coupling matrices, $m_{\tilde{g}}$ denotes the gluino mass, and $B\mu$ is the soft breaking mass for Higgs scalars. Although a large Λ'_{QCD} can help fulfill the first DAMP criterion, we should also explain how a Hubble induced mass fits in with eq. (1.3) to fulfill the second criterion.

The Kähler potential can give rise to a Hubble induced $B\mu$ term. If the argument of the term is different from the vacuum $B\mu$ term by π and dominates, a shift of π relative to the vacuum value is induced in the axion potential.² The difference of π in the arguments can be understood by the (approximate) CP symmetry of the theory, such that the $B\mu$ terms are real and the difference of π is simply the opposite signs of the terms. An approximate CP symmetry is also invoked in ref. [43], where a relaxation to $\theta_{\text{mis}} \simeq 0$ is considered. Note that the shift of π in the axion potential occurs only if the number of generations is odd.

A large m_a allows the axion field to relax to the bottom of the potential during inflation, and a π shifted axion potential means this minimum coincides with today's hilltop. Without additional particles beyond the MSSM, Λ'_{QCD} and consequently m_a cannot be arbitrarily large; we find $m_a \lesssim 10 \text{ TeV}$. Thus, in the minimal scenario, we consider TeV scales for Hubble during inflation H_I to allow for the relaxation of the axion misalignment during inflation. We also explore non-minimal models where m_a and thus H_I can be larger. Relaxing the axion arbitrarily close to today's hilltop may cause overproduction of axion DM, but we find that the running of Yukawa terms in the Standard Model (SM) gives a sufficient CP phase change $\mathcal{O}(10^{-16})$ to avoid the scenario [48, 49]. An exciting implication of this mechanism is that f_a is fixed to roughly $3 \times 10^9 \text{ GeV}$ by the observed DM abundance and CP-violating phase renormalization in the theory. We impose CP symmetry in the Higgs and inflaton sectors. Additional CP violation (CPV) of up to $\mathcal{O}(10^{-4})$ only induces $\mathcal{O}(1)$ changes in the prediction of f_a . In summary, by the inflationary dynamics of the Higgs fields as well as the (approximate) CP symmetry, we can fulfill both criteria of a DAMP scenario; in particular in this paper we explore the case where the inflationary minimum is shifted by π from today's minimum, which is referred to as DAMP $_{\pi}$.

We now elaborate on the approximate CP symmetry. Although the $\mathcal{O}(1)$ amount of CPV measured in the SM must be generated in the theory, a small CPV in the extended sectors can be a consequence of the suppressed couplings with the source of CP violation. Such hierarchical couplings can result from the protection of additional symmetries or the geometric separation in the extra dimensions. Additionally, any quantum corrections that attempt to transfer $\mathcal{O}(1)$ CPV from the SM to the extended sectors are automatically small. The reason is that the CP phase of the Yukawa couplings only becomes physical when all three generations are involved, suggesting that the interactions are suppressed by small Yukawa couplings, mixing among generations, and higher loop factors.³ With this

²If the arguments are the same, we may dynamically relax the axion to today's minimum during inflation as discussed in a separate paper [47].

³Even though the CP symmetry is a solution to the strong CP problem alternative to the axion, the $\mathcal{O}(1)$ CPV in the Yukawa sector may unacceptably modify the θ term. For models that avoid such consequences, refer to refs. [50–53].

CP structure, the implications for the extended sectors are as follows. The (approximate) CP symmetry treats the CP-odd and CP-even moduli differently in such a way that the moduli affecting the axion minimum can be stabilized at the CP-conserving points. The smallness of the CP phases is also guaranteed in the masses of any additional colored particles that we introduce in the non-minimal models. Crucially, a CP symmetry only ensures all relevant parameters are real but does not forbid the change of signs throughout the evolution; this is exactly what can give rise to a shift of π in the axion potential.

In section 2 we briefly review the axion misalignment mechanism and the role of the anharmonicity factor. We also discuss how the amount of CPV in a theory can be connected to the axion abundance in a DAMP $_{\pi}$ model. In section 3 we show how a Hubble induced mass for the Higgs in the early universe can induce an axion mass enhancement and a phase shift of π in the axion potential, fulfilling the DAMP $_{\pi}$ criteria. In section 4 we discuss both a set of minimal models with the cosmology fully evaluated, and extended models with a larger viable parameter space and a simplified discussion of the post-inflationary cosmology. Finally, in section 5 we summarize and discuss the possible phenomenological implications of this model as well as future directions.

2 Axion misalignment & early relaxation

We first review the axion misalignment mechanism. The equation of motion and energy density of axions are given by

$$\ddot{\theta}_a + 3H\dot{\theta}_a = -m_a^2 \sin \theta_a \tag{2.1}$$

$$\rho_a = \frac{1}{2} (m_a^2 a^2 + \dot{a}^2) \tag{2.2}$$

where $\theta_a \equiv a/f_a$ parametrizes the axion field value a and H is the Hubble expansion rate. In principle, all relevant parameters in eqs. (2.1) and (2.2) can vary throughout cosmological evolutions but we focus on the case where the PQ symmetry is already broken during inflation. The misalignment contribution to the axion DM abundance is as follows. In the conventional setup where m_a is assumed negligible compared to the Hubble parameter during inflation, the axion field value is practically frozen due to a large Hubble friction term in eq. (2.1) with the solution approximated by

$$\theta_0 \simeq \theta_i e^{-N_e m_a^2 / 3H_I^2} \quad \text{for } m_a \ll H_I, \tag{2.3}$$

with θ_0 (θ_i) the angle at the end (onset) of inflation, unless the number of e-folding is exceedingly large $N_e \sim (H_I/m_a)^2$ as pointed out by refs. [54–56]. As a result of inflation, the misalignment angle takes a random but uniform value θ_{mis} in the observable universe. Around the QCD phase transition, the axion acquires a mass from the QCD non-perturbative effects and starts to oscillate, when $3H \simeq m_a$, from $a_{\text{mis}} = \theta_{\text{mis}} f_a$ towards the minimum today. Without fine-tuning, θ_{mis} is expected to be order unity. The coherent oscillations of axions contribute to the cold dark matter abundance

$$\Omega_a h^2 = 0.12 \langle \theta_{\text{mis}}^2 \mathcal{F}(\theta_{\text{mis}}) \rangle \left(\frac{f_a}{5 \times 10^{11} \text{ GeV}} \right)^{7/6}, \tag{2.4}$$

where $\mathcal{F}(\theta_{\text{mis}}) \simeq 1$ for $\theta_{\text{mis}} \ll \pi$ and, for $\theta_{\text{mis}} \gtrsim 0.9\pi$, is analytically approximated by [38]

$$\mathcal{F}(\theta_{\text{mis}}) \simeq \frac{16\sqrt{2}}{\pi^3} \left[\ln \left(\frac{1}{1 - \theta_{\text{mis}}/\pi} \right) \right]^{7/6}. \quad (2.5)$$

Several numerical studies have been devoted to the determination of $\mathcal{F}(\theta_{\text{mis}})$ [37, 39, 40, 57] but DAMP $_{\pi}$ calls for a dedicated study for the extreme limit of $\pi - \theta_{\text{mis}} \ll 1$. The exponents in eqs. (2.4) and (2.5) assume the topological susceptibility of QCD given by the dilute instanton gas approximation (see the lattice results in refs. [58–62]) but our results are insensitive to this uncertainty.

We now discuss how our framework, by relaxing the above assumptions, makes a prediction for f_a using the DM abundance and the CP-violating phase $\delta\theta_{\text{CP}}$ of the theory. The axion mass arises from QCD dynamics. There is no a priori reason that the axion mass during inflation is given by the exact same QCD effect observed today. In fact, there are numerous scenarios where the axion is enhanced in the early universe, e.g. a large QCD confinement scale [42–46, 63], explicit PQ breaking [64–67], and magnetic monopoles [68, 69]. If the axion mass is larger than the Hubble scale during inflation, the axion starts oscillations and is rapidly relaxed towards the minimum,

$$\theta_0 \simeq \theta_i e^{-3N_e/2} \quad \text{for } m_a \gg H_I. \quad (2.6)$$

Additionally, if the CP phase of the model has a phase shift of π after inflation, as explained in section 1 and elaborated in section 3, the location of this minimum is then converted into the maximum of the potential, making the effective misalignment angle $\theta_{\text{mis}} \simeq \pi$. There is however a limit on how close θ_{mis} can be to π because the quantum correction to the θ parameter from the CP violation in the SM Yukawa couplings is $\delta\theta_{\text{CP}} \sim 10^{-16}$ [48, 49] and the running of the Yukawa couplings necessarily induces a phase shift of similar order between the inflationary and low energy scales. This small deviation from the hilltop $\delta\theta = \pi - \theta_{\text{mis}}$ allows for the prediction of f_a due to the anharmonic effects. In the limit $\theta_{\text{mis}} \rightarrow \pi$, $\mathcal{F}(\theta_{\text{mis}})$ and thus $\Omega_a h^2$ are only logarithmic dependent on $\delta\theta$ so one can predict f_a in terms of the deviation $\delta\theta$ by requiring DM abundance using eqs. (2.4) and (2.5)

$$f_a \simeq 2.4 \times 10^9 \text{ GeV} \left(\frac{\Omega h^2}{0.11} \right)^{6/7} \left(1 + 0.026 \ln \left(\frac{\delta\theta}{10^{-16}} \right) \right), \quad (2.7)$$

where we assume $\left| \ln \left(\frac{\delta\theta}{10^{-16}} \right) \right| \ll \ln \left(\frac{\pi}{\delta\theta_{\text{CP}}} \right) \simeq 38$ or equivalently $\delta\theta \ll 1$. This is the striking feature of the anharmonic effect — the value of f_a necessary for the DM abundance has an exceptionally mild logarithmic dependence on the CP phase shift as long as it is much less than unity. The sharp prediction of f_a is illustrated in figure 1 using the analytic approximations in eqs. (2.4) and (2.5). The prediction on the decay constant only changes by $\mathcal{O}(1)$ factor even if CPV of $\mathcal{O}(10^{-4})$ is added.

3 Dynamical axion misalignment production at the hilltop

We would like to show that allowing H_u and H_d to acquire large VEVs during inflation can lead to a DAMP $_{\pi}$ scenario. To guide the reader, we first restate the conditions under which




Figure 1. Insensitivity of f_a to very small shifts from the hilltop.

the DAMP model is applicable: 1) the axion field dynamically relaxes to the minimum of the potential in the early universe and 2) the model possesses a non-trivial prediction between the minima of the axion potential in the early and today’s epochs. Throughout our discussion of DAMP $_{\pi}$ models, we have in mind a minimal model as a proof of principle and extended models to further explore viable parameter space. Generically, we can include inflaton-Higgs dynamics with the effective operators suppressed by the cutoff scale M in the Kähler potential

$$\Delta K = \frac{|X|^2}{M^2} \left(|H_u|^2 + |H_d|^2 - (H_u H_d + \text{c.c.}) - \frac{|H_u|^2 |H_d|^2}{M^2} - \frac{|H_u|^4}{M^2} - \frac{|H_d|^4}{M^2} \right), \quad (3.1)$$

where X is the chiral field whose F -term provides the inflaton potential energy. We omit $\mathcal{O}(1)$ coupling constants here and hereafter. For illustration purposes, we only show lower dimensional operators relevant for the following discussion. Higher dimensional operators do not change the discussion.

During inflation, the inflaton F -term gives the Higgs fields Hubble induced terms,

$$\Delta V = cH_I^2 \left(-|H_u|^2 - |H_d|^2 + (H_u H_d + \text{c.c.}) + \frac{|H_u|^2 |H_d|^2}{M^2} + \frac{|H_u|^4}{M^2} + \frac{|H_d|^4}{M^2} \right), \quad (3.2)$$

where $c = (M_{\text{Pl}}/M)^2$ and H_I is the Hubble scale during inflation. We assume that the Hubble induced mass terms are negative. They push the Higgs fields in the D -flat direction $|H_u| = |H_d|$ up to the cutoff scale M , and as we will see in the following sections the large Higgs VEVs realize DAMP $_{\pi}$.

3.1 Axion mass during inflation

Together with the effective terms from the Kähler potential in eq. (3.1), the MSSM Higgs potential reads

$$\begin{aligned} V_{\text{Higgs}} = & (|\mu|^2 + m_{H_u}^2 - cH_I^2) |H_u|^2 + (|\mu|^2 + m_{H_d}^2 - cH_I^2) |H_d|^2 - (B\mu - cH_I^2) (H_u H_d + \text{c.c.}) \\ & + \frac{g^2 + g'^2}{8} (|H_u|^2 - |H_d|^2)^2 + \frac{g^2}{2} |H_u H_d^*|^2 + \frac{cH_I^2}{M^2} (|H_u|^2 |H_d|^2 + |H_u|^4 + |H_d|^4). \end{aligned} \quad (3.3)$$

We assume that the Higgs sector is nearly CP symmetric, which is anyway required from the limits on the electric dipole moment for TeV scale supersymmetry. See refs. [70] and [71] for the latest measurement and its implication to supersymmetric theories, respectively. We also assume a CP symmetry in the inflaton-Higgs coupling. The Higgs fields break $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$ by both the early universe VEV and today's VEV. Parameterizing the Higgs field space in terms of a radial mode $\phi \equiv |H_u| = |H_d|$ along the D -flat direction and an angular mode $\xi = \arg(H_u H_d)$, which is the relative phase of the Higgs fields, allows us to write

$$V \simeq (m_{\text{SUSY}}^2 - cH_I^2)\phi^2 - (B\mu - cH_I^2)\cos(\xi)\phi^2 + \frac{cH_I^2}{M^2}\phi^4, \quad (3.4)$$

where we have taken $m_{H_u} \sim m_{H_d} \sim \mu \sim m_{\text{SUSY}}$. The phases of $H_u H_d$ are chosen so that $\xi = 0$ in the vacuum today. The radial mode, for $\sqrt{c}H_I \gtrsim m_{\text{SUSY}}$, acquires a large VEV of order $\phi_i \sim M$. This is clearly seen from minimizing the potential. Assuming that the sign of the Hubble induced $B\mu$ term is opposite to the vacuum one as shown in eq. (3.4), the phase initially obtains a value during inflation of $\xi = \pi$, while today's value is $\xi = 0$. We discuss the implication of the phase shift in the next subsection and focus this subsection on the large radial direction.⁴

The large VEV ϕ_i gives quarks very large masses during inflation. In the MSSM, the 1-loop renormalization group equation (RGE) is

$$\mu_r \frac{d}{d\mu_r} \frac{8\pi^2}{g^2} = 3N - F, \quad (3.5)$$

where μ_r is the renormalization scale, $N = 3$ is the gauge group index, and F is the number of active fermions in the theory. Solving the RGE from the TeV scale up to the scale ϕ_i , and from the scale down while pretending that all quarks are above the scale where the gauge coupling diverges, we obtain the fiducial dynamical scale

$$\Lambda_{\text{fid}} = 10^7 \text{ GeV} \left(\frac{\phi_i}{10^{16} \text{ GeV}} \right)^{2/3} \left(\frac{\tan\beta}{1} \right)^{1/3}. \quad (3.6)$$

This is the physical dynamic scale Λ'_{QCD} if all quarks (including the KSVZ quarks [72, 73]) are above the scale. If some quarks are below the scale, the physical dynamical scale Λ'_{QCD} is given by

$$\Lambda_{\text{fid}} = \Lambda'_{\text{QCD}} \times \prod_{m_q < \Lambda'_{\text{QCD}}} \left(\frac{m_q}{\Lambda'_{\text{QCD}}} \right)^{1/9}. \quad (3.7)$$

The axion mass vanishes when the gluino is massless since strong dynamics gives the mass dominantly to the R-axion. The axion mass is hence given by

$$m_a \simeq \frac{1}{4\pi} \frac{m_{\tilde{g}}^{1/2} \Lambda_{\text{fid}}^{3/2}}{f_a}, \quad (3.8)$$

⁴In the extended model discussed below, the sign flip of the $B\mu$ is not necessary. The Hubble induced $B\mu$ term is not necessary as long as the vacuum one is larger than H_I^2 .

where we assume that the gluino mass is below the physical dynamical scale and that the large Higgs VEV does not break the PQ symmetry. We include the factor of 4π expected from the naive dimensional analysis [74–77]. Here $m_{\tilde{g}}$ is the RGE invariant one, $m_{\tilde{g},\text{phys}}/g^2$. The holomorphy of the gauge coupling guarantees that we may use the fiducial dynamical scale to evaluate the axion mass. Physically, the suppression of the fiducial dynamical scale in comparison with the physical dynamical scale takes into account the suppression of the axion mass by light quarks. For the minimal setup where the dynamical scale is raised solely by large Higgs VEVs as in eq. (3.6),

$$m_a \simeq 30 \text{ GeV} \left(\frac{m_{\tilde{g}}}{\text{TeV}} \right)^{1/2} \left(\frac{\Lambda_{\text{fid}}}{10^7 \text{ GeV}} \right)^{3/2} \left(\frac{3 \times 10^9 \text{ GeV}}{f_a} \right). \quad (3.9)$$

We may raise the dynamical scale further by introducing additional particles. One possibility is to introduce a moduli field whose field value controls the gauge coupling [42–46, 63], and assume that the moduli field value during inflation raises the gauge coupling constant. Another possibility is to introduce additional $\text{SU}(3)_c$ charged fields and assume that their masses are large during inflation as considered in ref. [45]. A field whose field value controls the masses of the additional particles can be regarded as a moduli field. For N_Ψ pairs of $\text{SU}(3)_c$ fundamental chiral fields with a mass M_Ψ and $M_{\Psi,I}$ in the vacuum and during inflation respectively, the dynamical scale is given by

$$\Lambda_{\text{fid}} = 10^7 \text{ GeV} \left(\frac{M_{\Psi,I}}{M_\Psi} \right)^{N_\Psi/9} \left(\frac{\phi_i}{10^{16} \text{ GeV}} \right)^{2/3} \left(\frac{\tan\beta}{1} \right)^{1/3}. \quad (3.10)$$

To achieve the second requirement of the DAMP scenario, CPV phases in $M_{\Psi,I}$ and M_Ψ should be absent. Instead of flipping the sign of the $B\mu$ term, we may flip the sign of the masses of Ψ to achieve DAMP $_\pi$. We will see later in section 4 that this dynamical scale cannot be arbitrarily large because of the backreaction of strong dynamics to the Higgs as well as the PQ sector.

Combining eqs. (3.8) and (3.10) we find that for appropriate values of H_I , the early universe axion mass is large enough for relaxation of the axion field to its minimum. Since the largeness of the dynamical scale Λ'_{QCD} depends on the VEV of ϕ , the decay of the inflaton and proceeding relaxation of ϕ to today's VEV means that the post-inflationary cosmology is non-trivial. Prior to exploring this complex cosmology, however, we turn our attention to the relative π phase shift of the axion potential.

3.2 Shifted axion potential

Another consequence of a large Higgs VEV during inflation from the Kähler potential in eq. (3.1) is that the relative phase between the Higgs fields is shifted by π as can be seen explicitly in eq. (3.4). The shift of ξ also shifts the minimum of the axion potential: eq. (1.3) shows a direct connection between the $B\mu$ term and the axion misalignment angle minimum θ_{eff} . Once the inflaton decays or its energy is redshifted and the Hubble induced terms become subdominant, minimization of the potential is achieved for $\xi = 0$. In the extended model discussed in the previous section, the sign flip of the masses of extra quarks Ψ can achieve a similar situation.

The phase shift of the axion potential is not exactly π because of the $\mathcal{O}(1)$ renormalization of Yukawa couplings. Since the CPV from Yukawa couplings manifests as $\mathcal{O}(10^{-16})$ shifts in the axion potential [48, 49], running these couplings from the early large Higgs VEVs to the electroweak scale necessarily induces an $\mathcal{O}(10^{-16})$ shift in the axion potential. We may also add small CPV to the $B\mu$ terms to induce further shift. Even if the shift is as large as $\mathcal{O}(10^{-4})$, the prediction of f_a changes only by an $\mathcal{O}(1)$ factor.

To summarize, a Kähler potential such as the one in eq. (3.1) gives Higgs fields Hubble induced masses, and the D -flat potential in eq. (3.4) is minimized at a large Higgs VEV with an opposite phase from today. The π shifted Higgs phase ξ along with the Yukawa coupling renormalization induce a shift in the axion potential by $\pi - \mathcal{O}(10^{-16})$. This sets the scene for a DAMP $_{\pi}$ scenario where the DM abundance is given by eq. (2.4) and the value of f_a can be predicted using the anharmonicity factor eq. (2.5).

4 Cosmological evolution

In section 3 we demonstrated that, in the early universe, both a large axion mass and a phase shift of π are possible due to a large Higgs VEV with an opposite phase from today. The remaining question is whether there exists a viable cosmology with a consistent evolution between the two periods without spoiling predictions. We first explore the inflationary and post-inflationary constraints in the minimal model, and then later comment on the broader parameter space allowed by extended models.

4.1 Minimal models

The first consistency check we should perform is to ensure the axion mass is larger than Hubble friction during inflation. As extensively noted, for a given cutoff scale M , the large Higgs VEV $\phi_i \simeq M$ determines Λ'_{QCD} and the axion mass during inflation is enhanced. The suppression of the angle by early relaxation can be approximated by eqs. (2.3) and (2.6). As a benchmark point, we require the suppression factor to be $\theta_0/\theta_i = 10^{-4}$ or smaller during the number of e-foldings of 60. This gives an upper bound on the value of H_I , which is shown in the blue regions of figure 2 with the left (right) panel for $M = M_{\text{GUT}} \equiv 2 \times 10^{16}$ GeV (M_{Pl}) respectively. The blue contours are also shown for $\theta_0/\theta_i = 10^{-16}$. The orange regions reflect a lower bound on the value of H_I from requiring $cH_I^2 > \max(m_{\text{SUSY}}^2, B\mu)$ necessary for obtaining a large Higgs VEV and the phase shift, respectively.

One needs to carefully consider the evolution of the axion potential after inflation ends. There must be a transition of the value of ξ from the inflationary minimum toward today's minimum. This transition necessarily induces the transition of the minimum from π to 0 in the axion potential. If this transition occurs at a time when the enhanced axion mass is still comparable to or larger than Hubble, the misalignment angle could relax to a value very different from π . To understand this constraint, we turn to the post-inflationary evolution of the Higgs fields.

The Hubble induced mass terms which stabilize ϕ at a large VEV are tied to the inflaton energy density. If the sign of the Hubble induced terms remains the same after inflation,




Figure 2. Parameter space for the inflationary Hubble scale H_I and reheate temperature T_R given $f_a = 3 \times 10^9$ GeV, $m_{\tilde{g}} = m_{\text{SUSY}} = \text{TeV}$, $B\mu = m_{\text{SUSY}}^2 / \tan\beta$, $\tan\beta = 50$, $N_e = 60$, and $\phi_i = M$. The left (right) panel is for the cutoff scale $M = M_{\text{GUT}}$ (M_{Pl}) respectively.

ϕ continues to be trapped around M . This means the radial and angular directions of the Higgs fields do not oscillate until the Hubble induced terms become subdominant to the MSSM soft terms of the corresponding mode as the inflaton energy density redshifts and/or decays.

A second possibility for this evolution is that the sign of the Hubble induced mass flips after inflation (except for the $H_u H_d$ term.) This may occur in two-field inflation models. For example, with $K = (-c_Z |Z|^2 + c_{\bar{Z}} |\bar{Z}|^2) |\phi|^2$ with $c_Z > c_{\bar{Z}}$ and $W = m_Z Z \bar{Z}$, we assume that the scalar component of Z acquires a large field value and drives inflation. It is \bar{Z} whose F -term, $F_{\bar{Z}}^2 = m_Z^2 \phi_{\bar{Z}}^2$, is non-zero. During inflation, the kinetic energy of ϕ_Z is much smaller than its potential energy, i.e. $|\partial_\mu Z|^2 \ll F_{\bar{Z}}^2$, and thus the Hubble induced mass for ϕ is negative. As inflation ends, Z 's potential and kinetic energies become comparable but, since $c_Z > c_{\bar{Z}}$, the sign of the Hubble induced mass for the Higgs radial mode flips to positive. Consequently, ϕ is no longer trapped at a large VEV but oscillates towards the origin immediately after inflation. The early onset of radial oscillations helps because a longer period of redshifting in ϕ suppresses Λ'_{QCD} , which leads to the desired post-inflationary suppression of the axion mass.

It is only necessary to track the ratio m_a/H between the onset of angular oscillations at $cH^2 = B\mu$ and thermalization of the Higgs fields. During this period, the Higgs phase ξ can evolve and with it comes the shift in the axion potential. If the axion mass is subdominant to Hubble friction, however, the axion field is overdamped and remains agnostic to this evolution. When the Higgs is finally thermalized, its energy density is depleted and the field is quickly set to the minimum today, removing the axion mass enhancement. As a result, to preserve the prediction of the axion misalignment angle, m_a/H needs to stay under unity during this period.

Thermalization of the Higgs is mediated by scattering with gluons via a loop-suppressed operator [78, 79]

$$\Gamma_h = \frac{B}{16\pi^2} \frac{T^2}{\phi}, \quad (4.1)$$

with $B \simeq 10^{-2}$ and ϕ identified as the oscillation amplitude. Interestingly, due to the scaling properties during a matter-dominated era, the Higgs scattering generates a radiation energy density that is constant in time, whose contribution to thermal bath's temperature is

$$T_h = \left(\frac{30}{\pi^2 g_*(T_h)} \frac{B}{16\pi^2} \right)^{1/2} \left(\frac{m_\phi^2 \phi_i}{H_I} \right)^{1/2}, \quad (4.2)$$

with ϕ_i as the field value of the Higgs at the end of inflation. This radiation persists throughout the evolution until the Higgs fields are thermalized at $H \simeq \Gamma_h$. This radiation is important because in some cases it can dominate over the radiation produced from the inflaton decay and cause a period of a constant temperature in the cosmological evolution. This has the effect of maintaining the finite temperature suppression to the axion mass

$$m_a(T) = \frac{1}{4\pi} \frac{m_{\tilde{g}}^{1/2} \Lambda_{\text{fid}}^{3/2}}{f_a} \left(\frac{\Lambda'_{\text{QCD}}}{T} \right)^n, \quad (4.3)$$

where $n = 3$ ($n = 0$) for $T > \Lambda'_{\text{QCD}}$ ($T < \Lambda'_{\text{QCD}}$). The temperature dependence is determined by the contribution from the gauge multiplets, while the contribution from chiral multiplets vanishes because of the cancellation between the RGE contribution and the fermion mass suppression. In the extended models, Λ'_{QCD} may be different from the estimate in eq. (3.7) because of the backreaction from strong QCD dynamics. We note that the value of m_a evolves after inflation not only due to this temperature suppression, but also its explicit dependence on $\Lambda_{\text{fid}} \propto \phi^{2/3}$, which evolves as the Higgs oscillation amplitude redshifts until the Higgs fields thermalize and settle into today's vacuum. Including the decrease in Λ'_{QCD} from the redshift in $\phi \propto H^{-\frac{1}{1+w}}$, with the equation of state of the total energy density w , the axion mass scales like $m_a|_{\text{MD}} \propto H^{\frac{3+2n}{3}}/T^n$ during the early matter-dominated era ($w = 0$) and $m_a|_{\text{RD}} \propto H^{\frac{3+2n}{4}}/T^n$ during the radiation-dominated era ($w = 1/3$) after reheating. These considerations of the Higgs oscillations and axion mass suppression are taken into account when determining the post-inflationary constraint in the regions shaded in dark gray (light gray enclosed by the dashed contour) in figure 2 assuming that the radial direction starts oscillation at $\sqrt{c}H \simeq m_{\text{SUSY}}$ (right after inflation) respectively. This constraint is milder in the left panel because $M = M_{\text{GUT}}$ starts out with a smaller axion mass during inflation than $M = M_{\text{Pl}}$ so m_a/H is more likely to be less than unity during the transition period. In fact, these gray regions disappear for $M \lesssim 10^{16}$ GeV opening up regions of low T_R even though the blue constraint becomes stronger.

Another requirement of DAMP_π comes from avoiding PQ symmetry restoration. In a thermal environment, the PQ breaking field saxion P acquires a thermal mass $y^2 T^2 P^2$ because of the Yukawa coupling $yPQ\bar{Q}$ with the PQ quarks Q, \bar{Q} . This thermal mass can be relevant at large temperatures and stabilize P at a vanishing value to restore PQ

symmetry. This can be easily prevented if the symmetry breaking is enforced by the following superpotential,

$$\Delta W_P = \lambda S (P\bar{P} - f_a^2). \tag{4.4}$$

The F -term of S stabilizes P and \bar{P} in the moduli space $P\bar{P} = f_a^2$ to break PQ. The coupling constant λ may be as large as 4π in strongly coupled models [80, 81]. To ensure that PQ is not thermally restored after inflation where the maximum temperature achieved during reheating is $T_{\text{max}} \simeq (H_I M_{\text{Pl}} T_R^2)^{1/4}$,⁵ the thermal mass must be less than $\lambda f_a < 4\pi f_a$ at this time, giving an upper bound on the Yukawa coupling

$$y \lesssim 1.5 \left(\frac{m_{P,\bar{P}}}{f_a} \right) \left(\frac{f_a}{3 \times 10^9 \text{ GeV}} \right) \left(\frac{\text{TeV}}{H_I} \right)^{1/4} \left(\frac{10^{10} \text{ GeV}}{T_R} \right)^{1/2}, \tag{4.5}$$

which can be easily satisfied with $y \lesssim \mathcal{O}(1)$ in the allowed parameter space of figure 2. The constraint is stronger if the mass of the PQ symmetry breaking field is only as large as m_{SUSY} , but we do not pursue this issue further.

A runaway potential for P is generated by strong dynamics via $W_{\text{eff}} \simeq (1/4\pi)^2 \cdot \Lambda_{\text{fid}}^3 (P/f_a)^{1/3}$, which introduces an effective saxion mass of order $(1/4\pi)^2 \Lambda_{\text{fid}}^3 / f_a^2$. Keeping in mind the dependence of Λ_{fid} on both the cutoff scale M and $\tan \beta$ from eq. (3.10), we find this effective saxion mass for the parameters in figure 2 to be of order 10^2 GeV (10^6 GeV) for $M = M_{\text{GUT}}$ (M_{Pl}) respectively. For values of H_I in the allowed parameter space, a Hubble induced mass for P is not always large enough to stabilize P . For simplicity, we stabilize P by superpotential terms $W = m_P P Y + m_{\bar{P}} \bar{P} \bar{Y}$. The F -terms of Y and \bar{Y} give large masses $m_P \sim m_{\bar{P}}$ to P and \bar{P} . These masses can be as large as $4\pi f_a$ without destroying the moduli space $P\bar{P} = f_a^2$ and are large enough to stabilize P against the runaway potential. The quarks Q and \bar{Q} have a large mass yP , which allows us to neglect the effects of their Hubble induced masses in the minimal models. We will see in section 4.2 that larger values of Λ'_{QCD} and H_I will modify the dynamics of these fields.

Finally, the purple regions in figure 2 are excluded by energy conservation which restricts the reheat temperature T_R to a maximum value dictated by the energy in the inflaton, $\rho_I \simeq M_{\text{Pl}}^2 H_I^2 \gtrsim T_R^4$. In the red regions, the axion starts to oscillate from the hilltop towards today's minimum during a matter-dominated era by the inflaton, in which case reheating produces entropy, dilutes the axion abundance, and spoils the prediction of a small f_a . Even though the minimal model has proven to provide a viable cosmology for DAMP $_{\pi}$, we explore extended models in the following subsection to further broaden the parameter space.

4.2 Extended models

The most stringent constraint in the minimal model is a relatively low upper bound on H_I due to difficulties in enhancing the axion mass during inflation. As shown in eq. (3.10), we can enhance Λ'_{QCD} , and consequently m_a , by introducing additional matter content. As we raise the value of Λ'_{QCD} , the values of fields in the PQ sector may be shifted from the one

⁵The actual maximal temperature is smaller after taking into account the efficiency of the thermalization [82].

in the vacuum and impact the evaluation of the axion mass. The field value of the Higgs may be also affected.

We consider a simple model where a PQ symmetry breaking field P couples to KSVZ quarks $Q\bar{Q}$ by a Yukawa coupling y [72, 73]. We need to reliably evaluate the VEVs of both P and $Q\bar{Q}$, as both are PQ charged and the decay constant during inflation f_I is given by the larger of P and $(Q\bar{Q})^{1/2}$. The superpotential of P and $Q\bar{Q}$ is

$$W = \frac{1}{(4\pi)^3} \frac{\tilde{\Lambda}^4}{(Q\bar{Q})^{1/2}} + yPQ\bar{Q}, \tag{4.6}$$

where the first term is the non-perturbative Affleck-Dine-Seiberg superpotential [83]. The scale $\tilde{\Lambda}$ is related with Λ_{fid} via

$$\tilde{\Lambda} = \Lambda_{\text{fid}} \left(\frac{\Lambda_{\text{fid}}}{yf_a} \right)^{1/8}. \tag{4.7}$$

The relation can be obtained by comparing the effective superpotential after integrating out $Q\bar{Q}$ from eq. (4.6) and the effective potential

$$W_{\text{eff}} \simeq \frac{1}{(4\pi)^2} \Lambda_{\text{fid}}^3 \left(\frac{P}{f_a} \right)^{1/3}, \tag{4.8}$$

where the effect of a P field value different from f_a is included. Note that the potential of P from strong dynamics exhibits a runaway behavior, $\partial W_{\text{eff}}/\partial P \propto P^{-2/3}$. For this reason, we introduce a higher dimensional term $|P|^6/M^2$ to further stabilize P at a large field value which can come from a superpotential term $(\chi/M)P^3$, where χ is a chiral field. We also consider the Hubble induced mass of P and $Q\bar{Q}$. Explicitly we take

$$\Delta V \simeq cH_I^2|Q|^2 + cH_I^2|\bar{Q}|^2 - cH_I^2|P|^2 + \frac{|P|^6}{M^2}. \tag{4.9}$$

We stress that these terms are used in our analysis but they are not the only possible extensions to DAMP $_{\pi}$. The sign of the Hubble induced mass of Q and \bar{Q} is taken to be positive to ensure that Q is not destabilized by the Hubble induced mass. We study a negative Hubble induced mass for P for the following reason. We find that $Q > P$ and hence f_I is dominated by Q for this choice of the signs. If the Hubble induced mass of P is positive instead, the field value of P becomes smaller and makes the field value of Q larger because of the smaller mass of Q . This increases f_I while decreasing the dynamical scale and suppressing the axion mass.

As the Hubble induced mass of Q breaks the supersymmetry of the QCD sector, the axion mass may non-trivially depends on the parameters. When yP is larger than Q , $Q\bar{Q}$ can be integrated out at the mass threshold yP and the effective potential is given by eq. (4.8). The axion mass is given by

$$m_a \simeq \frac{1}{4\pi} \frac{m_{\tilde{g}}^{1/2} \Lambda_{\text{fid}}^{3/2}}{f_I} \left(\frac{P}{f_a} \right)^{1/6}. \tag{4.10}$$




Figure 3. Parameter space for the inflationary Hubble scale H_I and the fiducial confinement scale Λ_{fid} defined in eq. (3.7) given $f_a = 3 \times 10^9$ GeV, $m_{\tilde{g}} = m_{\text{SUSY}} = \text{TeV}$, and $\phi_i = M$. The left (right) panel is for the cutoff scale $M = M_{\text{GUT}}$ (M_{Pl}) respectively.

If Q is larger than yP , the theory below the mass threshold Q is a supersymmetric, pure $SU(2)$ gauge theory with a dynamical scale $\tilde{\Lambda}(\tilde{\Lambda}/Q)^{1/3}$. The axion mass is then given by

$$m_a \simeq \frac{1}{4\pi} \frac{m_{\tilde{g}}^{1/2} \tilde{\Lambda}^{3/2}}{f_I} \left(\frac{\tilde{\Lambda}}{4\pi Q} \right)^{1/2} = \frac{1}{4\pi} \frac{m_{\tilde{g}}^{1/2} \Lambda_{\text{fid}}^{3/2}}{f_I} \left(\frac{\Lambda_{\text{fid}}^3}{16\pi^2 y f_a Q^2} \right)^{1/4}. \quad (4.11)$$

Note that the two formulae agree with each other when the field value of Q is determined by the F -term condition of $Q\bar{Q}$ from the superpotential in eq. (4.6).

Strong dynamics also affects the Higgs. The effective superpotential of ϕ is given by

$$W \simeq \frac{\Lambda_{\text{eff}}^3}{16\pi^2} \left(\frac{\phi}{M} \right)^2, \quad \Lambda_{\text{eff}}^3 = \begin{cases} \Lambda_{\text{fid}}^3 \left(\frac{P}{f_a} \right)^{1/3} & yP > Q \\ \Lambda_{\text{fid}}^3 \left(\frac{\Lambda_{\text{fid}}^3}{16\pi^2 y f_a Q^2} \right)^{1/2} & yP < Q. \end{cases} \quad (4.12)$$

This gives the Higgs a mass $\simeq \Lambda_{\text{eff}}^3/(16\pi^2 M^2)$, which should be smaller than $\sqrt{c}H_I$.

By computing and comparing the axion mass to H_I , we put an upper bound on the allowed values of H_I such that DAMP $_{\pi}$'s first criterion is fulfilled during inflation, which is shown in figure 3. In deriving the blue-shaded region, we integrate out $Q\bar{Q}$, obtain the scalar potential of P from the effective potential eq. (4.8), add the potential of P in eq. (4.9), determine the field value of P during inflation, and compute the axion mass. This corresponds to the case where Q is actually determined by the F -term condition $\partial W/\partial Q = 0$. The gray contours show the constraint using the full potential described above. As y becomes larger the constraints approach to the blue-shaded region. An additional constraint shown in the red regions arises because strong dynamics drives the Higgs to the origin.

In computing the field values of P and Q , we treat Q as a canonically normalized field. This is a good approximation if yP or Q is above the dynamical scale. We find that in the allowed parameter space, either yP or Q is no smaller than one order of magnitude below the dynamical scale so we expect the approximation gives a good order of magnitude estimate.

We now discuss the post-inflationary evolution and constraints similar to section 4.1. Although this study is comprehensive in evaluating the inflationary constraints, the thermal masses for P and Q dramatically complicate PQ dynamics during and after reheating. Nonetheless, we are able to identify a large allowed parameter space in the H_I, T_R plane in the following way. There exists a wide region where the Higgs fields thermalize before the Higgs angular mode ξ begins to oscillate so that Λ'_{QCD} is quickly set to today's value Λ_{QCD} before ξ has a chance to evolve and shift the axion potential. Physically, this means that the axion potential turns off before the location of its minimum shifts. Therefore, the prediction of the misalignment angle is automatically preserved without the need to track the post-inflationary evolution of P and $Q\bar{Q}$. This is the case when the Higgs scattering rate in eq. (4.1) equals the Hubble rate before cH^2 drops below $B\mu$. This region is described by an allowed window of T_R for a given H_I

$$T_R \gtrsim 10^8 \text{ GeV} \left(\frac{M}{M_{\text{GUT}}} \right)^{3/2} \left(\frac{B\mu}{\text{TeV}} \right)^{3/4} \left(\frac{\phi_i}{M_{\text{GUT}}} \right) \left(\frac{10^5 \text{ GeV}}{H_I} \right), \quad (4.13)$$

$$T_R \lesssim 6 \times 10^{11} \text{ GeV} \left(\frac{M_{\text{GUT}}}{M} \right)^{3/2} \left(\frac{\text{TeV}}{B\mu} \right)^{3/4} \left(\frac{H_I}{10^5 \text{ GeV}} \right)^2 \left(\frac{M_{\text{GUT}}}{\phi_i} \right)^2, \quad (4.14)$$

where the two distinct formulae come from thermalization during the matter- and radiation-dominated epochs respectively. This window becomes wider as H_I increases so as long as

$$H_I > 6 \text{ TeV} \left(\frac{M}{M_{\text{GUT}}} \right) \left(\frac{B\mu}{\text{TeV}} \right)^{1/2} \left(\frac{\phi_i}{M_{\text{GUT}}} \right) \quad (4.15)$$

a consistent range of T_R exists. The upper bound on H_I ultimately enters from the inflationary constraint shown in the blue regions of figure 3.

We now comment on one plausible extension to further open up the parameter space with higher H_I . The gluino mass affects the axion mass as in eq. (3.8) and is assumed to stay invariant between inflation and today. If $m_{\tilde{g}}$ is also larger during inflation, the axion mass and the upper bound on H_I can be raised by as much as $(\Lambda_{\text{fid}}/m_{\tilde{g}})^{1/2}$, making high scale inflation easily compatible with DAMP $_{\pi}$.

With respect to PQ restoration in the extended models, we can assume the same PQ breaking mechanism as in section 4.1 so the constraint in eq. (4.5) applies equally here.

While the post-inflationary constraints are not fully evaluated, these extended models have been shown capable of fulfilling the criteria of DAMP $_{\pi}$ while extending the allowed parameter space to much higher H_I than in section 4.1.

5 Conclusion

It has been widely known that the misalignment mechanism can source axion dark matter in the early universe with a decay constant $f_a \simeq \mathcal{O}(10^{12})$ GeV. For $f_a \ll 10^{12}$ GeV as is of

interest to many experimental searches, the observed DM abundance can be obtained if the misalignment angle θ_{mis} is taken sufficiently close to π , where the anharmonic effect becomes important. In particular, when the axion is very close to the hilltop of the potential, the onset of oscillations is delayed so the axion abundance is less redshifted and thus more enhanced. As demonstrated in figure 1 and eq. (2.7), $f_a \simeq 10^{10}$ GeV corresponds to $\delta\theta \equiv \pi - \theta_{\text{mis}} \simeq \mathcal{O}(10^{-3})$, while $f_a \simeq 4 \times 10^9$ GeV already requires $\delta\theta \simeq \mathcal{O}(10^{-9})$. Such a small $\delta\theta$ has generically been understood as fine-tuning of the initial condition. In this paper, we offer an explanation to this small $\delta\theta$ using axion dynamics in the early universe.

We point out that a class of models violates the canonical assumption that the axion field is overdamped by Hubble friction and takes a random value during inflation. Instead, there exist numerous possibilities wherein the axion is large compared to Hubble during inflation and thus relaxes to the minimum of the potential. We refer to this mechanism as Dynamical Axion Misalignment Production (DAMP). Additionally, if the model possesses an approximate CP symmetry, then the axion potential may receive a phase shift of π because the nearly real parameters for setting the axion minimum can flip the sign between inflation and the QCD phase transition. This shift converts the potential minimum into a maximum and explains why the axion is very close to the hilltop — a mechanism we dub DAMP_π .

We explicitly construct models for DAMP_π , where the higher dimensional coupling between the Higgs in the MSSM and the inflaton gives rise to a large axion mass and the phase shift of the axion potential. Specifically, a negative Hubble induced mass drives the Higgs to a large field value that enhances the quark masses, which in turn raise the QCD scale. The axion is larger than usual due to stronger QCD dynamics. Lastly, a Hubble induced $B\mu$ term that carries an opposite sign from that of the MSSM necessarily induces a shift in the axion potential by π . Together, renormalization from the SM Yukawa couplings and any additional CP violating phases in the model can provide the desired finite phase shift between $\mathcal{O}(10^{-16}\text{--}10^{-3})$. This example works only if the number of generations is odd.

Strikingly, due to the anharmonic effects of the axion potential, the prediction of f_a from the DM abundance has an extraordinarily mild logarithmic dependence on $\delta\theta \ll 1$. Therefore, DAMP_π makes a rather sharp prediction of $3 \times 10^9 \text{ GeV} \lesssim f_a \lesssim 10^{10} \text{ GeV}$. Other phenomenological features of DAMP_π are as follows. Due to early relaxation, the fluctuation of the axion field is exponentially damped and hence dark matter isocurvature perturbations are suppressed. The upper bound on H_I from isocurvature perturbations does not apply.

The impact of this anharmonicity on the structure formation has been investigated in the literature. Refs. [37, 39] study this numerically and show that isocurvature perturbation modes whose wavelengths are larger than the horizon size are enhanced by anharmonic effects. Refs. [84, 85] numerically study the anharmonicity effects on the growth of structures arising from the large fluctuations in an inhomogeneous background, i.e. in the context of post-inflationary PQ symmetry breaking. Finally, ref. [86] has shown that in a quasi-homogeneous region, parametric resonance can be important for amplifying fluctuations, and this effect is monotonically enhanced for larger misalignment angles. While the anharmonicity effect may stimulate structure formation, the aforementioned works are not

directly applicable to this model as the isocurvature perturbations are suppressed and the PQ symmetry is already broken during inflation.

The axions from the misalignment mechanism are necessarily cold — a feature to distinguish from other non-thermal production mechanisms. It is also potentially interesting to study the imprints of maximal CP violation on the QCD phase transition as well as Big Bang nucleosynthesis.

Acknowledgments

The authors thank Asimina Arvanitaki, Joshua W. Foster, Andrew J. Long, Aaron Pierce, Benjamin R. Safdi, and Ken van Tilburg for discussions. K.H. thanks the Leinweber Center for Theoretical Physics for the warm hospitality he received during his visit when most of this work was done. The work was supported in part by the DoE Early Career Grant DE-SC0019225 (R.C.) and the DoE grant DE-SC0009988 (K.H.).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] R.J. Crewther, P. Di Vecchia, G. Veneziano and E. Witten, *Chiral Estimate of the Electric Dipole Moment of the Neutron in Quantum Chromodynamics*, *Phys. Lett.* **B 88** (1979) 123 [*Erratum ibid.* **B 91** (1980) 487] [[INSPIRE](#)].
- [2] C.A. Baker et al., *An Improved experimental limit on the electric dipole moment of the neutron*, *Phys. Rev. Lett.* **97** (2006) 131801 [[hep-ex/0602020](#)] [[INSPIRE](#)].
- [3] G. 't Hooft, *Symmetry Breaking Through Bell-Jackiw Anomalies*, *Phys. Rev. Lett.* **37** (1976) 8 [[INSPIRE](#)].
- [4] R.D. Peccei and H.R. Quinn, *CP Conservation in the Presence of Instantons*, *Phys. Rev. Lett.* **38** (1977) 1440 [[INSPIRE](#)].
- [5] R.D. Peccei and H.R. Quinn, *Constraints Imposed by CP Conservation in the Presence of Instantons*, *Phys. Rev.* **D 16** (1977) 1791 [[INSPIRE](#)].
- [6] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40** (1978) 223 [[INSPIRE](#)].
- [7] F. Wilczek, *Problem of Strong P and T Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40** (1978) 279 [[INSPIRE](#)].
- [8] J.R. Ellis and K.A. Olive, *Constraints on Light Particles From Supernova SN 1987A*, *Phys. Lett.* **B 193** (1987) 525 [[INSPIRE](#)].
- [9] G. Raffelt and D. Seckel, *Bounds on Exotic Particle Interactions from SN1987A*, *Phys. Rev. Lett.* **60** (1988) 1793 [[INSPIRE](#)].
- [10] M.S. Turner, *Axions from SN1987A*, *Phys. Rev. Lett.* **60** (1988) 1797 [[INSPIRE](#)].
- [11] R. Mayle, J.R. Wilson, J.R. Ellis, K.A. Olive, D.N. Schramm and G. Steigman, *Constraints on Axions from SN 1987A*, *Phys. Lett.* **B 203** (1988) 188 [[INSPIRE](#)].

- [12] G.G. Raffelt, *Astrophysical axion bounds*, in proceedings of the *Joint ILLIAS-CAST-CERN Axion Training at CERN*, Geneva, Switzerland, 30 November–2 December 2005, *Lect. Notes Phys.* **741** (2008) 51 [[hep-ph/0611350](#)] [[INSPIRE](#)].
- [13] J. Preskill, M.B. Wise and F. Wilczek, *Cosmology of the Invisible Axion*, *Phys. Lett. B* **120** (1983) 127 [[INSPIRE](#)].
- [14] L.F. Abbott and P. Sikivie, *A Cosmological Bound on the Invisible Axion*, *Phys. Lett. B* **120** (1983) 133 [[INSPIRE](#)].
- [15] M. Dine and W. Fischler, *The Not So Harmless Axion*, *Phys. Lett. B* **120** (1983) 137 [[INSPIRE](#)].
- [16] R.L. Davis, *Cosmic Axions from Cosmic Strings*, *Phys. Lett. B* **180** (1986) 225 [[INSPIRE](#)].
- [17] M. Kawasaki, K. Saikawa and T. Sekiguchi, *Axion dark matter from topological defects*, *Phys. Rev. D* **91** (2015) 065014 [[arXiv:1412.0789](#)] [[INSPIRE](#)].
- [18] V.B. Klaer and G.D. Moore, *The dark-matter axion mass*, *JCAP* **11** (2017) 049 [[arXiv:1708.07521](#)] [[INSPIRE](#)].
- [19] M. Gorghetto, E. Hardy and G. Villadoro, *Axions from Strings: the Attractive Solution*, *JHEP* **07** (2018) 151 [[arXiv:1806.04677](#)] [[INSPIRE](#)].
- [20] M. Kawasaki, T. Sekiguchi, M. Yamaguchi and J. Yokoyama, *Long-term dynamics of cosmological axion strings*, *Prog. Theor. Exp. Phys.* **2018** (2018) 091E01 [[arXiv:1806.05566](#)] [[INSPIRE](#)].
- [21] C.J.A.P. Martins, *Scaling properties of cosmological axion strings*, *Phys. Lett. B* **788** (2019) 147 [[arXiv:1811.12678](#)] [[INSPIRE](#)].
- [22] J.K. Vogel et al., *IAXO — The International Axion Observatory*, in proceedings of the *8th Patras Workshop on Axions, WIMPs and WISPs (AXION-WIMP 2012)*, Chicago, Illinois, U.S.A., 18–22 July 2012, [FERMILAB-PUB-13-699-A](#) (2013) [[arXiv:1302.3273](#)] [[INSPIRE](#)] and online pdf version at <http://lss.fnal.gov/archive/2013/pub/fermilab-pub-13-699-a.pdf>.
- [23] E. Armengaud et al., *Conceptual Design of the International Axion Observatory (IAXO)*, *2014 JINST* **9** T05002 [[arXiv:1401.3233](#)] [[INSPIRE](#)].
- [24] TASTE collaboration, *Towards a medium-scale axion helioscope and haloscope*, *2017 JINST* **12** P11019 [[arXiv:1706.09378](#)] [[INSPIRE](#)].
- [25] G. Rybka, A. Wagner, A. Brill, K. Ramos, R. Percival and K. Patel, *Search for dark matter axions with the Orpheus experiment*, *Phys. Rev. D* **91** (2015) 011701 [[arXiv:1403.3121](#)] [[INSPIRE](#)].
- [26] MADMAX Working Group, *Dielectric Haloscopes: A New Way to Detect Axion Dark Matter*, *Phys. Rev. Lett.* **118** (2017) 091801 [[arXiv:1611.05865](#)] [[INSPIRE](#)].
- [27] A. Arvanitaki and A.A. Geraci, *Resonantly Detecting Axion-Mediated Forces with Nuclear Magnetic Resonance*, *Phys. Rev. Lett.* **113** (2014) 161801 [[arXiv:1403.1290](#)] [[INSPIRE](#)].
- [28] ARIADNE collaboration, *Progress on the ARIADNE axion experiment*, in proceedings of the *2nd Workshop on Microwave Cavities and Detectors for Axion Research*, Livermore, California, U.S.A., 10–13 January 2017, *Springer Proc. Phys.* **211** (2018) 151 [[arXiv:1710.05413](#)] [[INSPIRE](#)].
- [29] P. Sikivie, *Axion Dark Matter Detection using Atomic Transitions*, *Phys. Rev. Lett.* **113** (2014) 201301 [[arXiv:1409.2806](#)] [[INSPIRE](#)].

- [30] A. Arvanitaki, S. Dimopoulos and K. Van Tilburg, *Resonant absorption of bosonic dark matter in molecules*, *Phys. Rev. X* **8** (2018) 041001 [[arXiv:1709.05354](#)] [[INSPIRE](#)].
- [31] M. Baryakhtar, J. Huang and R. Lasenby, *Axion and hidden photon dark matter detection with multilayer optical haloscopes*, *Phys. Rev. D* **98** (2018) 035006 [[arXiv:1803.11455](#)] [[INSPIRE](#)].
- [32] ADMX collaboration, *A Search for Invisible Axion Dark Matter with the Axion Dark Matter Experiment*, *Phys. Rev. Lett.* **120** (2018) 151301 [[arXiv:1804.05750](#)] [[INSPIRE](#)].
- [33] R.T. Co, L.J. Hall and K. Harigaya, *QCD Axion Dark Matter with a Small Decay Constant*, *Phys. Rev. Lett.* **120** (2018) 211602 [[arXiv:1711.10486](#)] [[INSPIRE](#)].
- [34] T. Hiramatsu, M. Kawasaki and K. Saikawa, *Evolution of String-Wall Networks and Axionic Domain Wall Problem*, *JCAP* **08** (2011) 030 [[arXiv:1012.4558](#)] [[INSPIRE](#)].
- [35] T. Hiramatsu, M. Kawasaki, K. Saikawa and T. Sekiguchi, *Axion cosmology with long-lived domain walls*, *JCAP* **01** (2013) 001 [[arXiv:1207.3166](#)] [[INSPIRE](#)].
- [36] K. Harigaya and M. Kawasaki, *QCD axion dark matter from long-lived domain walls during matter domination*, *Phys. Lett. B* **782** (2018) 1 [[arXiv:1802.00579](#)] [[INSPIRE](#)].
- [37] M.S. Turner, *Cosmic and Local Mass Density of Invisible Axions*, *Phys. Rev. D* **33** (1986) 889 [[INSPIRE](#)].
- [38] D.H. Lyth, *Axions and inflation: Sitting in the vacuum*, *Phys. Rev. D* **45** (1992) 3394 [[INSPIRE](#)].
- [39] K. Strobl and T.J. Weiler, *Anharmonic evolution of the cosmic axion density spectrum*, *Phys. Rev. D* **50** (1994) 7690 [[astro-ph/9405028](#)] [[INSPIRE](#)].
- [40] K.J. Bae, J.-H. Huh and J.E. Kim, *Update of axion CDM energy*, *JCAP* **09** (2008) 005 [[arXiv:0806.0497](#)] [[INSPIRE](#)].
- [41] L. Visinelli and P. Gondolo, *Dark Matter Axions Revisited*, *Phys. Rev. D* **80** (2009) 035024 [[arXiv:0903.4377](#)] [[INSPIRE](#)].
- [42] G.R. Dvali, *Removing the cosmological bound on the axion scale*, [hep-ph/9505253](#) [[INSPIRE](#)].
- [43] T. Banks and M. Dine, *The Cosmology of string theoretic axions*, *Nucl. Phys. B* **505** (1997) 445 [[hep-th/9608197](#)] [[INSPIRE](#)].
- [44] K. Choi, H.B. Kim and J.E. Kim, *Axion cosmology with a stronger QCD in the early universe*, *Nucl. Phys. B* **490** (1997) 349 [[hep-ph/9606372](#)] [[INSPIRE](#)].
- [45] K.S. Jeong and F. Takahashi, *Suppressing Isocurvature Perturbations of QCD Axion Dark Matter*, *Phys. Lett. B* **727** (2013) 448 [[arXiv:1304.8131](#)] [[INSPIRE](#)].
- [46] K. Choi, E.J. Chun, S.H. Im and K.S. Jeong, *Diluting the inflationary axion fluctuation by a stronger QCD in the early Universe*, *Phys. Lett. B* **750** (2015) 26 [[arXiv:1505.00306](#)] [[INSPIRE](#)].
- [47] R.T. Co, E. Gonzalez and K. Harigaya, *Axion Misalignment Driven to the Bottom*, *JHEP* **05** (2019) 162 [[arXiv:1812.11186](#)] [[INSPIRE](#)].
- [48] J.R. Ellis and M.K. Gaillard, *Strong and Weak CP-violation*, *Nucl. Phys. B* **150** (1979) 141 [[INSPIRE](#)].
- [49] I.B. Khriplovich and A.R. Zhitnitsky, *What Is the Value of the Neutron Electric Dipole Moment in the Kobayashi-Maskawa Model?*, *Phys. Lett. B* **109** (1982) 490 [[INSPIRE](#)].

- [50] A.E. Nelson, *Naturally Weak CP-violation*, *Phys. Lett. B* **136** (1984) 387 [INSPIRE].
- [51] S.M. Barr, *Solving the Strong CP Problem Without the Peccei-Quinn Symmetry*, *Phys. Rev. Lett.* **53** (1984) 329 [INSPIRE].
- [52] L. Bento, G.C. Branco and P.A. Parada, *A Minimal model with natural suppression of strong CP-violation*, *Phys. Lett. B* **267** (1991) 95 [INSPIRE].
- [53] G. Hiller and M. Schmaltz, *Solving the Strong CP Problem with Supersymmetry*, *Phys. Lett. B* **514** (2001) 263 [hep-ph/0105254] [INSPIRE].
- [54] S. Dimopoulos and L.J. Hall, *Inflation and Invisible Axions*, *Phys. Rev. Lett.* **60** (1988) 1899 [INSPIRE].
- [55] P.W. Graham and A. Scherlis, *Stochastic axion scenario*, *Phys. Rev. D* **98** (2018) 035017 [arXiv:1805.07362] [INSPIRE].
- [56] F. Takahashi, W. Yin and A.H. Guth, *QCD axion window and low-scale inflation*, *Phys. Rev. D* **98** (2018) 015042 [arXiv:1805.08763] [INSPIRE].
- [57] O. Wantz and E.P.S. Shellard, *Axion Cosmology Revisited*, *Phys. Rev. D* **82** (2010) 123508 [arXiv:0910.1066] [INSPIRE].
- [58] P. Petreczky, H.-P. Schadler and S. Sharma, *The topological susceptibility in finite temperature QCD and axion cosmology*, *Phys. Lett. B* **762** (2016) 498 [arXiv:1606.03145] [INSPIRE].
- [59] S. Borsányi et al., *Calculation of the axion mass based on high-temperature lattice quantum chromodynamics*, *Nature* **539** (2016) 69 [arXiv:1606.07494] [INSPIRE].
- [60] F. Burger, E.-M. Ilgenfritz, M.P. Lombardo and A. Trunin, *Chiral observables and topology in hot QCD with two families of quarks*, *Phys. Rev. D* **98** (2018) 094501 [arXiv:1805.06001] [INSPIRE].
- [61] C. Bonati, M. D’Elia, G. Martinelli, F. Negro, F. Sanfilippo and A. Todaro, *Topology in full QCD at high temperature: a multicanonical approach*, *JHEP* **11** (2018) 170 [arXiv:1807.07954] [INSPIRE].
- [62] M. Gorghetto and G. Villadoro, *Topological Susceptibility and QCD Axion Mass: QED and NNLO corrections*, *JHEP* **03** (2019) 033 [arXiv:1812.01008] [INSPIRE].
- [63] S. Ipek and T.M.P. Tait, *Early Cosmological Period of QCD Confinement*, *Phys. Rev. Lett.* **122** (2019) 112001 [arXiv:1811.00559] [INSPIRE].
- [64] T. Higaki, K.S. Jeong and F. Takahashi, *Solving the Tension between High-Scale Inflation and Axion Isocurvature Perturbations*, *Phys. Lett. B* **734** (2014) 21 [arXiv:1403.4186] [INSPIRE].
- [65] M. Kawasaki, M. Yamada and T.T. Yanagida, *Cosmologically safe QCD axion as a present from extra dimension*, *Phys. Lett. B* **750** (2015) 12 [arXiv:1506.05214] [INSPIRE].
- [66] F. Takahashi and M. Yamada, *Strongly broken Peccei-Quinn symmetry in the early Universe*, *JCAP* **10** (2015) 010 [arXiv:1507.06387] [INSPIRE].
- [67] J. Kearney, N. Orlofsky and A. Pierce, *High-Scale Axions without Isocurvature from Inflationary Dynamics*, *Phys. Rev. D* **93** (2016) 095026 [arXiv:1601.03049] [INSPIRE].
- [68] M. Kawasaki, F. Takahashi and M. Yamada, *Suppressing the QCD Axion Abundance by Hidden Monopoles*, *Phys. Lett. B* **753** (2016) 677 [arXiv:1511.05030] [INSPIRE].

- [69] Y. Nomura, S. Rajendran and F. Sanches, *Axion Isocurvature and Magnetic Monopoles*, *Phys. Rev. Lett.* **116** (2016) 141803 [[arXiv:1511.06347](#)] [[INSPIRE](#)].
- [70] ACME collaboration, *Improved limit on the electric dipole moment of the electron*, *Nature* **562** (2018) 355 [[INSPIRE](#)].
- [71] C. Cesarotti, Q. Lu, Y. Nakai, A. Parikh and M. Reece, *Interpreting the Electron EDM Constraint*, [arXiv:1810.07736](#) [[INSPIRE](#)].
- [72] J.E. Kim, *Weak Interaction Singlet and Strong CP Invariance*, *Phys. Rev. Lett.* **43** (1979) 103 [[INSPIRE](#)].
- [73] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Can Confinement Ensure Natural CP Invariance of Strong Interactions?*, *Nucl. Phys. B* **166** (1980) 493 [[INSPIRE](#)].
- [74] A. Manohar and H. Georgi, *Chiral Quarks and the Nonrelativistic Quark Model*, *Nucl. Phys. B* **234** (1984) 189 [[INSPIRE](#)].
- [75] H. Georgi and L. Randall, *Flavor Conserving CP-violation in Invisible Axion Models*, *Nucl. Phys. B* **276** (1986) 241 [[INSPIRE](#)].
- [76] M.A. Luty, *Naive dimensional analysis and supersymmetry*, *Phys. Rev. D* **57** (1998) 1531 [[hep-ph/9706235](#)] [[INSPIRE](#)].
- [77] A.G. Cohen, D.B. Kaplan and A.E. Nelson, *Counting 4π 's in strongly coupled supersymmetry*, *Phys. Lett. B* **412** (1997) 301 [[hep-ph/9706275](#)] [[INSPIRE](#)].
- [78] D. Bödeker, *Moduli decay in the hot early Universe*, *JCAP* **06** (2006) 027 [[hep-ph/0605030](#)] [[INSPIRE](#)].
- [79] K. Mukaida and K. Nakayama, *Dynamics of oscillating scalar field in thermal environment*, *JCAP* **01** (2013) 017 [[arXiv:1208.3399](#)] [[INSPIRE](#)].
- [80] K. Harigaya, M. Ibe, K. Schmitz and T.T. Yanagida, *Peccei-Quinn Symmetry from Dynamical Supersymmetry Breaking*, *Phys. Rev. D* **92** (2015) 075003 [[arXiv:1505.07388](#)] [[INSPIRE](#)].
- [81] K. Harigaya and J. Leedom, *Unified Models of the QCD Axion and Supersymmetry Breaking*, *Nucl. Phys. B* **921** (2017) 507 [[arXiv:1702.00401](#)] [[INSPIRE](#)].
- [82] K. Harigaya and K. Mukaida, *Thermalization after/during Reheating*, *JHEP* **05** (2014) 006 [[arXiv:1312.3097](#)] [[INSPIRE](#)].
- [83] I. Affleck, M. Dine and N. Seiberg, *Dynamical Supersymmetry Breaking in Supersymmetric QCD*, *Nucl. Phys. B* **241** (1984) 493 [[INSPIRE](#)].
- [84] E.W. Kolb and I.I. Tkachev, *Nonlinear axion dynamics and formation of cosmological pseudosolitons*, *Phys. Rev. D* **49** (1994) 5040 [[astro-ph/9311037](#)] [[INSPIRE](#)].
- [85] E.W. Kolb and I.I. Tkachev, *Axion miniclusters and Bose stars*, *Phys. Rev. Lett.* **71** (1993) 3051 [[hep-ph/9303313](#)] [[INSPIRE](#)].
- [86] P.B. Greene, L. Kofman and A.A. Starobinsky, *Sine-Gordon parametric resonance*, *Nucl. Phys. B* **543** (1999) 423 [[hep-ph/9808477](#)] [[INSPIRE](#)].