



# Fermi National Accelerator Laboratory

FERMILAB-PUB-91/354-T

## A Determination of the Strong Coupling Constant from the Charmonium Spectrum

Aida X. El-Khadra, George Hockney,  
Andreas S. Kronfeld, and Paul B. Mackenzie

*Theoretical Physics Group  
Fermi National Accelerator Laboratory  
P.O. Box 500  
Batavia, IL 60510*

### ABSTRACT

Lattice gauge theory techniques have recently achieved sufficient accuracy to permit a determination of the strong coupling constant from the 1P-1S splitting in the Charmonium system, with all systematic errors estimated quantitatively. The present result is  $\alpha_{\overline{MS}}(5 \text{ GeV}) = 0.174 \pm 0.012$ , or equivalently,  $\Lambda_{\overline{MS}}^{(4)} = 160^{+47}_{-37} \text{ MeV}$ .



A central task in understanding quantum chromodynamics (QCD) is the determination of its coupling constant,  $g^2$ . The Review of Particle Properties quotes values for  $\alpha_s(5 \text{ GeV}) \equiv g^2/4\pi$  in the range 0.18–0.22.[1] Recent LEP measurements yield values in the range 0.20–0.24.[2] Most perturbative determinations of  $g^2$  contain nonperturbative contaminations which become small only at high energies. On the other hand, high energy determinations yield  $g^2$  at lower energies only imprecisely. Lattice gauge theory calculations provide a nonperturbative means of determining the strong coupling constant from low energy quantities.

In principle, any lattice calculation of a mass or energy  $E$  allows a determination of the strong coupling constant. The lattice calculation yields the dimensionless quantity  $aE$ , where  $a$  is the lattice spacing which is determined by comparing  $aE$  with the experimentally measured value for  $E$ . The bare lattice coupling constant  $g_0^2$  at scale  $a$  may then be converted into one of the more familiar definitions of the coupling constant using known perturbative results.[3, 4] In practice, most existing lattice calculations contain systematic errors which are difficult to analyze quantitatively. Consider for example the obvious case of the proton mass. Lattice calculations have not yet been done with quark masses as light as their physical values. Chiral perturbation theory calculations[5] indicate that at pion masses of around 400 MeV, where lattice calculations are often done, the proton mass is reduced by a term (of order  $m_\pi^3$ ) of around 100% of its physical, light pion value. Similarly, the most accurate lattice calculations to date have been done ignoring the effects of sea quarks. Some chiral quark model calculations[6] estimate that the proton mass may be altered as much as 30% by the effects of the strange quarks in the sea, let alone the light quarks. Whether or not these calculations are quantitatively reliable, the point is that the approximation of ignoring the sea quarks (the “quenched” approximation) introduces potentially large systematic errors for the light hadrons which are difficult to analyze and control.

Heavy quark systems offer the best opportunity for determining the strong coupling constant with present day lattice calculations.[7] For these systems no extrapolation to light valence quark masses is necessary, and errors arising from the omission of sea quarks and also from the finiteness of the lattice spacing may be systematically analyzed and quantitatively estimated with some input from phenomenology, as we discuss below.

The cleanest quantity in heavy quark systems from which to extract the strong coupling constant is the splitting between the spin averaged masses of the 1S and 1P states. This splitting is insensitive to errors in spin dependent interactions which are induced by the finite size of the lattice spacing. It is also known to be quite insensitive to any errors in the definition of the quark mass, since the 1P–1S splittings in the  $\psi$  and  $\Upsilon$  systems are almost identical. Higher order finite lattice spacing errors such as those resulting in effective  $p^4$  interactions and four Fermi interactions may be easily analyzed perturbatively using potential models and Coulomb gauge lattice wave functions, and if necessary removed by corrections to the lattice action.

We have calculated this splitting using standard Monte Carlo techniques at three lattice spacings (or equivalently, three values of  $\beta \equiv 6/g_0^2$ ). The smallest lattice spacing used corresponds to  $\beta = 6.1$  on lattice volumes of  $24^4$  with 25 gauge configurations separated by 8000 pseudo-heat bath sweeps. The larger lattice spacings correspond to  $\beta = 5.9$  on volumes of  $16^4$  and  $\beta = 5.7$  on volumes of  $12^3 \times 24$ , each with 25 configurations separated by 2000 pseudo-heat bath sweeps.

The Wilson action for fermions was used with the  $O(a)$  correction term,  $-\frac{i}{2}c\bar{\psi}\Sigma_{\mu\nu}F_{\mu\nu}\psi$ , added.[8] (Addition of this operator to the action suffices to correct  $O(a)$  errors for Wilson fermions.) The coefficient  $c$  was set to 1.4 rather than its tree level value  $c = 1$  on the basis of mean field theory estimates of the higher order corrections. For nonrelativistic fermions, it contributes mainly an additional  $\sigma \cdot B$  interaction to the quarks. The spin averaged level splitting on which this paper is based is expected to be very insensitive to this correction. On the other hand, such spin splittings as the  $\psi - \eta_c$  splitting which we have also investigated are very sensitive to it.[9]

The lattice spacing for each value of  $\beta$  was obtained by calculating in lattice units the difference of the spin averaged mass of the 1S states (the  $J/\psi$  and the  $\eta_c$ ) and the mass of the recently discovered spin singlet 1P state (the  $h_c$ )[10], and then comparing with the experimentally measured splitting,  $M_{h_c} - (3M_\psi + M_{\eta_c})/4 = 458.6 \pm 0.4$  MeV. Coulomb gauge wave functions were used to create and destroy the meson states to reduce errors from excited states.

From the lattice spacing  $a$  and the bare lattice coupling constant  $g_0^2$ , the  $\overline{MS}$  coupling at scale  $\pi/a$  may be obtained using the one loop perturbative




Figure 1: The wave function of the  $\psi$  meson.

formula  $g_{\overline{MS}}^{-2}(\frac{\pi}{a}) = g_0^{-2}(1 - \frac{1}{3}g^2) + 0.025$ . [3, 4] The background field calculation of this correction [4] shows that it is dominated by the term in parentheses, which is the perturbative expectation value of the plaquette  $\langle Tr U_P \rangle$ . We have therefore substituted the known nonperturbative value of the plaquette at each  $\beta$  to correct for higher order effects in the relation between the bare and  $\overline{MS}$  coupling constants [11], using

$$\frac{1}{g_{\overline{MS}}^2(\frac{\pi}{a})} = \frac{1}{g_0^2} \langle Tr U_P \rangle_{MC} + 0.025. \quad (1)$$

(The values for  $\langle Tr U_P \rangle_{MC}$  are 0.549, 0.582, and 0.605 at  $\beta = 5.7, 5.9$ , and  $6.1$ , respectively.) This yields an additional correction to the  $\overline{MS}$  coupling constant (11% of the final value) which is much smaller than the one loop correction (about 44%) but not negligible.

Figure 1 shows the Coulomb gauge wave function of the  $\psi$  meson calculated on the  $24^4$ ,  $\beta = 6.1$  lattices. It is probably not controversial that finite volume errors are negligible for the lattice sizes used. To check this, a low statistics calculation was done at  $\beta = 6.1$  on a volume of  $16^4$  which yielded a value of the 1P-1S




Figure 2:  $\Lambda^{(0)}_{\overline{MS}}$  as a function of  $a^2$ .

splitting 10–20% larger than that on the  $24^4$  lattices. Figure 1 shows that the wave function of the  $\psi$  is about a factor of five smaller at a distance of 12 lattice spacings (half way across the  $24^4$  lattice) than it is at 8 lattice spacings (half way across the  $16^4$  lattice). Assuming that finite volume errors fall roughly as the square of the wave function halfway across the lattice leads to the conclusion that the errors on the  $24^4$  lattice are under a percent in the spin splitting and therefore in  $\Lambda_{\overline{MS}}$ . This implies errors in  $\alpha$  of a fraction of a percent.

We have already noted that the spin averaged 1P–1S splitting is very insensitive to the  $O(a)$  errors of the Wilson quark action. Further, we have used a corrected action which minimizes these errors. It is therefore to be expected that the most important finite lattice spacing errors remaining will be of order  $a^2$ . To test for the size of these, the calculation was performed at three lattice spacings. From the 1P–1S splitting, we obtained  $a^{-1} = 1.15(8)$  GeV,  $1.78(9)$  GeV, and  $2.43(15)$  GeV at  $\beta = 5.7$ ,  $5.9$ , and  $6.1$  respectively. The errors in parentheses

are statistical only. Using Equation (1) and the parameterization for  $\alpha_s$  of the Particle Data Group, values for  $\Lambda_{\overline{MS}}^{(0)}$  were obtained for each lattice spacing. In Figure 2 the results are plotted as a function of  $a^2$ . The value of  $\Lambda_{\overline{MS}}^{(0)} = 234$  MeV after extrapolation to  $a^2 = 0$  is about 4% larger than its value at  $\beta = 6.1$ . The statistical errors are not small enough to distinguish between possible functional forms for the finite  $a$  errors (e. g.,  $a$ ,  $a^2$ , or  $a^4$ ), and we rely on theoretical prejudice in making the extrapolation in  $a^2$ . We therefore take the 4% difference between the extrapolated value and the value at  $\beta = 6.1$  as a contribution to the systematic uncertainty. The next step in removing this small contribution to the uncertainty will be the evaluation of the known discretization errors present in the lattice action with the Coulomb gauge wave functions obtained from the lattice calculations, to verify or improve the functional form of the extrapolation in Figure 2.

The final and largest source of uncertainty arises from the conversion from the zero light quark running coupling constant of the lattice calculation to the four quark running coupling of the real world. It is more convenient to discuss this uncertainty in terms of  $g^2$  rather than  $\Lambda$ . The adjustment of the bare parameters of the lattice action to reproduce experimental physics is really the adjustment of the parameters in the effective action at the physics momentum scale to be correct. Since the  $\beta$  function of the quenched (zero light quark) lattice theory is slightly too large, the short distance coupling constant of the quenched lattice theory must be slightly too small. It is this discrepancy which must finally be estimated. In perturbation theory, requiring that the effective actions of the lattice and the real world match at the physics scale  $\mu_1$  requires approximately that the running coupling constants match at this scale. The required difference in the running couplings at scale  $\mu_2$  is then given by

$$\Delta g^{-2} = \int_{\mu_1}^{\mu_2} d \ln q^2 (\beta_0^{(n_f)} - \beta_0^{(0)} + (\beta_1^{(n_f)} - \beta_1^{(0)})g^2 + \dots), \quad (2)$$

where  $\beta_0^{(n_f)} = (11 - 2/3 n_f)/(16\pi^2)$ , and  $\beta_1^{(n_f)} = (102 - 38/3 n_f)/(16\pi^2)^2$ . From potential models we know that the typical momentum transfer in  $J/\psi$  mesons is around  $\mu_1 \approx 400$  MeV. Most of the required correction arises from the large  $\mu$  region where perturbation theory is valid, and is therefore reliably given by Equation (2). Integrating the correction from  $\mu_2 = 5$  GeV down to 750 MeV, the scale of typical quark momentum in the  $J/\psi$ , and making the appropriate

change from the four quark to the three quark  $\beta$  function at the charmed quark mass, gives a contribution to the correction of  $\Delta g^{-2} = -0.080$ . ( $g_V$ , the coupling constant defined from the heavy quark potential, was used in the integration.) This is certainly an underestimate of the correction, since the  $\beta$  function is quite convergent in this region and 750 MeV is the largest momentum scale in the systems we are studying. If we take  $\mu_1 \approx 400$  MeV as the best guess for the appropriate matching scale, we obtain an additional correction of  $\Delta g^{-2} = -0.060$ . This large contribution from a small region of integration arises from the fact that 400 MeV is near the fictitious pole in the two-loop perturbative coupling constant, causing  $g_V^2$  to begin to blow up near the end of the integration region. (At  $\mu_1 \approx 400$  MeV, we obtained  $g_V^2 = 45$ .) Although the true contribution of the low energy region to the coupling constant correction cannot be reliably estimated perturbatively, it is certainly not divergent, so this is almost certainly an overestimate of the true correction. Since the effects of light quarks on the potential are relatively mild both at short distances (known from the perturbative  $\beta$  functions) and at large distances (known from comparing the string tensions obtained from Regge phenomenology and from quenched lattice calculations), it is implausible that their effects in the intermediate region are overwhelmingly severe. We therefore take these two estimates as upper and lower bounds on the true correction, giving  $\Delta g^{-2} = -0.110 \pm 0.030$ . The approximate size of this correction, 24%, clearly makes sense, since if the comparison scale 5 GeV is replaced by an asymptotically large value  $\mu_2$ , the obtained correction must approach  $(\beta_0^{(n_f)} - \beta_0^{(0)})/\beta_0^{(0)} = (2/3)(n_f/11) \approx 24\%$  for  $n_f = 4$ . Over the next few years, this largest source of uncertainty will be eliminated by the inclusion of sea quarks in lattice calculations. In the short term, it may be clarified and, we hope, reduced, with the use of potential models and a study of the static potentials of the quenched and unquenched lattice theories.

Our final result is

$$\alpha_{\overline{MS}}(5 \text{ GeV}) = 0.174 \pm 0.012. \quad (3)$$

This corresponds to  $\Lambda_{\overline{MS}}^{(4)} = 160_{-37}^{+47}$  MeV, using the parameterization of the Particle Data Group. The known corrections and uncertainties are summarized in Table 1. Each correction  $\epsilon_i$  is defined to mean that the corrected coupling  $\alpha_i$  is given by  $\alpha_i = (1 + \epsilon_i)\alpha_{i-1}$ . The uncertainty is dominated by the uncertainty in

Source	Correction	Uncertainty
$g_0^2 \rightarrow g_{\overline{MS}}^2$ (One loop, Ref. [3, 4])	44%	-
$g_0^2 \rightarrow g_{\overline{MS}}^2$ (Monte Carlo correction, Ref. [11])	11%	-
$g^{(0)} \rightarrow g^{(4)}$	24%	6.6%
Statistics	-	2%
Finite lattice spacing	1%	1%
Finite volume	-	-

Table 1: Summary of corrections and uncertainties in the determination of  $\alpha_{\overline{MS}}^{(4)}$ . Note that all of the corrections have the same sign, raising the obtained value of  $\alpha$ .

relating the zero quark coupling constant of the lattice calculation to the four quark coupling constant of the real world.

Over the next few years, using Monte Carlo simulations including the effects of sea quarks, it will be possible to eliminate the errors dominant in the present calculation arising from the correction in Equation 2. The stated uncertainty does not include the 5 – 10% uncertainty intrinsic to any second order perturbative QCD calculation. This arises in this calculation solely from the introduction of Equation 1 to convert to the  $\overline{MS}$  definition of the coupling constant for the purposes of comparison with existing results. It can be eliminated by the adoption of a physical definition of the strong coupling constant such as  $g_V$ , the coupling defined from the static potential, for the purposes of comparison. Such a coupling constant may be determined nonperturbatively directly from the static potential on the lattice (and of course computed to as many loops as desired for the purposes of short distance analyses).

Recently, a similar calculation has been performed for both the  $\psi$  and  $\Upsilon$  systems using the nonrelativistic formulation of lattice fermions[12]. For the  $\Upsilon$  system, the systematic errors and corrections are quite different from the ones reported here. The results are compatible.

Extrapolating to the mass of the  $Z$ , we obtain  $\alpha_{\overline{MS}}(M_Z) = 0.105 \pm 0.004$ . This is a little more than  $2\sigma$  below the combined results from LEP:  $\alpha(M_Z) = 0.119 \pm 0.006$ [2].



## Acknowledgements

We thank Peter Lepage and Estia Eichten for helpful conversations. This calculation was performed on the Fermilab lattice supercomputer, ACPMAPS. We would like to thank our colleagues in the Fermilab Computer Research and Development Group for their collaboration. Fermilab is operated by Universities Research Association, Inc. under contract with the U.S. Department of Energy.

## References

- [1] The Particle Data Group, Phys. Lett. B239 (1990).
- [2] H. Wachsmuth, paper given at Particles and Fields '91, Vancouver, Canada, Aug. 18-22 1991, CERN preprint PPE/91-145.
- [3] A. Hasenfratz and P. Hasenfratz, Phys. Lett. B93 (1980) 165.
- [4] R. Dashen and D. Gross, Phys. Rev. D23 (1981) 2340.
- [5] J. Gasser and H. Leutwyler, Phys. Repts. 87 (1982) 77, and references therein.
- [6] D. B. Kaplan and A. Manohar, Nuc. Phys. B310 (1988) 527.
- [7] The importance of heavy quark systems for lattice gauge theory has been emphasized by G. P. Lepage and B. A. Thacker, in *Field Theory on the Lattice* A. Billoire et al., editors, Nuc. Phys. B (Proc. Suppl.) 4 (1988) 199.
- [8] B. Sheikholeslami and R. Wohlert, Nuc. Phys. B259 (1985) 572.
- [9] Talks given by P. B. Mackenzie and A. X. El-Khadra at *Lattice '91*, Tsukuba, Japan, Nov. 5-9, 1991, Fermilab preprints CONF-92/09-T and CONF-92/10-T.
- [10]  $M_{h_c} = 3526.15 \pm 0.15$  MeV, J. Peoples, private communication.
- [11] G. P. Lepage and P. B. Mackenzie, Fermilab preprint 91/355-T.
- [12] C. T. H. Davies, B. A. Thacker and G. P. Lepage, to be published; talk by G. P. Lepage at *Lattice '91*, Tsukuba, Japan, Nov. 5-9, 1991.