

Time-dependent correlation functions for a bimodal Bose–Hubbard model

Zapiski Nauchnykh Seminarov POMI, 2015, **433**, 65–77

References

- [1] G. Milburn, J. Corney, E. Wright, D. Walls, “Quantum dynamics of an atomic Bose–Einstein condensate in a double-well potential”, *Phys. Rev. A*, **55** (1997), 4318 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#)
- [2] D. Witthaut, F. Trimborn, S. Wimberger, “Dissipation induced coherence of a two-mode Bose–Einstein condensate”, *Phys. Rev. Lett.*, **101** (2008), 200402 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#)
- [3] E. Boukobza, M. Chuchem, D. Cohen, A. Vardi, “Phase-diffusion dynamics in weakly coupled Bose–Einstein condensates”, *Phys. Rev. Lett.*, **102** (2009), 180403 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#) [LIBRARY.RU](#)
- [4] T. Pudlik, H. Hennig, D. Witthaut, D. Campbell, “Dynamics of entanglement in a dissipative Bose–Hubbard dimer”, *Phys. Rev. A*, **88** (2013), 063606 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#)
- [5] F. Trimborn, D. Witthaut, V. Kegel, H. J. Korsch, “Nonlinear Landau-Zener tunneling in quantum phase space”, *New J. Phys.*, **12** (2010), 053010 [crossref](#) [WEB OF SCIENCE™](#) [LIBRARY.RU](#)
- [6] I. Tikhonenkov, M. G. Moore, A. Vardi, “Robust sub-shot-noise measurement via Rabi-Josephson oscillations in bimodal Bose–Einstein condensates”, *Phys. Rev. A*, **83** (2011), 063628 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#) [LIBRARY.RU](#)
- [7] M. Chuchem, K. Smith-Mannschott, M. Hiller, T. Kottos, A. Vardi, D. Cohen, “Quantum dynamics in the bosonic Josephson junction”, *Phys. Rev. A*, **82** (2010), 053617 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#) [LIBRARY.RU](#)
- [8] V. Giovannetti, S. Lloyd, L. Maccone, “Quantum-Enhanced Measurements: Beating the Standard Quantum Limit”, *Science*, **306** (2004), 1330 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#)
- [9] D. Jaksch, H.-J. Briegel, J. Cirac, C. Gardiner, P. Zoller, “Entanglement of Atoms via Cold Controlled Collisions”, *Phys. Rev. Lett.*, **82** (1999), 1975 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#)
- [10] L. D. Faddeev, “Quantum completely integrable models of field theory”, *Sov. Sci. Rev. Math. C*, **1** (1980), 107–160; *40 Years in Mathematical Physics*, World Sci. Ser. 20th Century Math., **2**, World Sci., Singapore, 1995, 187–235 [crossref](#) [MathSciNet](#)
- [11] P. P. Kulish, E. K. Sklyanin, “Quantum spectral transform method. Recent developments”, *Lecture Notes Phys.*, **151**, 1982, 61–119 [crossref](#) [MathSciNet](#) [Zentralblatt MATH](#)
- [12] V. Z. Enol’skii, V. B. Kuznetsov, M. Salerno, “On the quantum inverse scattering method for the DST dimer”, *Phys. D*, **68** (1993), 138 [crossref](#) [MathSciNet](#) [Zentralblatt MATH](#) [WEB OF SCIENCE™](#)
- [13] J. Links, K. Hibberd, “Bethe Ansatz Solutions of the Bose–Hubbard Dimer”, *SIGMA*, **2** (2006), Paper 095 [Math-Net.Ru](#) [crossref](#) [MathSciNet](#) [Zentralblatt MATH](#)
- [14] R. Orús, S. Dusuel, J. Vidal, “Equivalence of Critical Scaling Laws for Many-Body Entanglement in the Lipkin–Meshkov–Glick Model”, *Phys. Rev. Lett.*, **101** (2008), 025701 [crossref](#) [ads*](#) [WEB OF SCIENCE™](#)
- [15] N. M. Bogoliubov, R. K. Bullough, J. Timonen, “Exact solution of generalised Tavis–Cummings models in quantum optics”, *J. Phys. A*, **29** (1996), 6305 [crossref](#) [MathSciNet](#) [Zentralblatt MATH](#) [ads*](#) [WEB OF SCIENCE™](#)
- [16] N. M. Bogolyubov, P. P. Kulish, “Tochno-reshaemye modeli kvantovoi nelineinoi optiki”, *Zap. nauchn. sem. POMI*, **398**, 2012, 26 [Math-Net.Ru](#) [MathSciNet](#)
- [17] N. M. Bogolyubov, A. G. Izergin, V. E. Korepin, *Korrelatsionnye funktsii integriruemyykh sistem i kvantovyi metod obratnoi zadachi*, Nauka, Moskva, 1992 [MathSciNet](#)

- [18] V. E. Korepin, N. M. Bogoliubov, A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press, Cambridge, 1993 [MathSciNet](#) [ZentralMATH](#)
- [19] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Oxford University Press, Oxford, 1995 [MathSciNet](#) [ZentralMATH](#)
- [20] N. A. Slavnov, “Vychislenie skalyarnykh proizvedenii volnovykh funktsii i formfaktorov v ramkakh algebraicheskogo anzatsa Bete”, *Teor. Matem. Fiz.*, **79**:2 (1989), 232 [Math-Net.Ru](#) [MathSciNet](#)
- [21] N. Kitanine, J. M. Maillet, V. Terras, “Form factors of the XXZ Heisenberg spin-1/2 finite chain”, *Nucl. Phys. B*, **516** (1999), 647 [crossref](#) [MathSciNet](#) [ads*](#)
- [22] V. E. Korepin, “Calculation of norms of Bethe wave functions”, *Comm. Math. Phys.*, **86** (1982), 391 [crossref](#) [MathSciNet](#) [ZentralMATH](#) [ads*](#) [WEB OF SCIENCE™](#)
- [23] M. Hillery, M. Zubairy, “Entanglement Conditions for Two-Mode States”, *Phys. Rev. Lett.*, **96** (2006), 050503 [crossref](#) [MathSciNet](#) [ads*](#) [WEB OF SCIENCE™](#)
- [24] Q. He, M. Reid, T. Vaughan, C. Gross, M. Oberthaler, P. Drummond, “Einstein–Podolsky–Rosen Entanglement Strategies in Two-Well Bose–Einstein Condensates”, *Phys. Rev. Lett.*, **106** (2011), 120405 [crossref](#) [WEB OF SCIENCE™](#)