


Towards integrability for $\text{AdS}_3/\text{CFT}_2$

Alessandro Sfondrini



2014

Cover: scattering of two wave-packets in $1 + 1$ dimensions.

Printed by Ipskamp Drukkers, Enschede.

ISBN: 978-94-6259-086-1.

Towards integrability for $\text{AdS}_3/\text{CFT}_2$

Op weg naar integreerbaarheid voor $\text{AdS}_3/\text{CFT}_2$

(met een samenvatting in het Nederlands)

Proefschrift

ter verkrijging van de graad van doctor aan de Universiteit Utrecht
op gezag van de rector magnificus, prof.dr. G.J. van der Zwaan,
ingevolge het besluit van het college voor promoties in het openbaar
te verdedigen op maandag 28 april 2014 des middags te 4.15 uur

door

Alessandro Sfondrini

geboren op 2 december 1987

te Como, Italië

Promotor: Prof.dr. G.E. Arutyunov

This thesis was accomplished with financial support from the Netherlands Organization for Scientific Research (NWO) under the VICI grant 680-47-602.

*Io stimo più il trovare un vero, benché di cosa leggiera,
che l' disputar lungamente delle massime questioni
senza conseguir verità nissuna.*

— Galileo

Author's contributions

This thesis is based on the following research by the author:

- [S1] R. Borsato, O. Ohlsson Sax, A. Sfondrini and B. Stefański, Jr.,
“All-loop worldsheet S matrix for $AdS_3 \times S^3 \times T^4$ ”,
arxiv:1403.4543,
- [S2] R. Borsato, O. Ohlsson Sax, A. Sfondrini, B. Stefański, Jr. and A. Torrielli,
“Dressing phases of AdS_3/CFT_2 ”,
Phys.Rev. D88, 066004 (2013), arxiv:1306.2512,
- [S3] R. Borsato, O. Ohlsson Sax, A. Sfondrini, B. Stefański, Jr. and A. Torrielli,
“The all-loop integrable spin-chain for strings on $AdS_3 \times S^3 \times T^4$: the massive sector”,
JHEP 1308, 043 (2013), arxiv:1303.5995,
- [S4] R. Borsato, O. Ohlsson Sax and A. Sfondrini,
“All-loop Bethe ansatz equations for AdS_3/CFT_2 ”,
JHEP 1304, 116 (2013), arxiv:1212.0505,
- [S5] R. Borsato, O. Ohlsson Sax and A. Sfondrini,
“A dynamic $\mathfrak{su}(1|1)^2$ S -matrix for AdS_3/CFT_2 ”,
JHEP 1304, 113 (2013), arxiv:1211.5119,

The author has also contributed to the following papers:

- [S6] R. Gurau, V. Rivasseau and A. Sfondrini,
“Renormalization: an advanced overview”,
arxiv:1401.5003,
- [S7] A. Sfondrini,
“An introduction to universality and renormalization group techniques”,
PoS ModaveVIII, 005 (2012), arxiv:1210.2262,
- [S8] G. Arutyunov, S. Frolov and A. Sfondrini,
“Exceptional operators in $\mathcal{N} = 4$ super Yang-Mills”,
JHEP 1209, 006 (2012), arxiv:1205.6660,
- [S9] A. Sfondrini and S. J. van Tongeren,
“Lifting asymptotic degeneracies with the Mirror TBA”,
JHEP 1109, 050 (2011), arxiv:1106.3909,
- [S10] A. Sfondrini and T. A. Koslowski,
“Functional renormalization of noncommutative scalar field theory”,
Int.J.Mod.Phys. A26, 4009 (2011), arxiv:1006.5145.
- [S11] F. Cardin, G. D. Marco and A. Sfondrini,
“Finite reduction and Morse index estimates for mechanical systems”,
Nonlinear Differential Equations and Applications 18, 557 (2011), arxiv:1004.1133.
- [S12] A. Sfondrini, J. Cerrillo, N. Schuch and J. I. Cirac,
“Simulating 2- and 3-dim. frustrated quantum systems with string-bond states”,
Phys. Rev. B81, 214426 (2010), arxiv:0908.4036.

Contents

1	Introduction	1
1.1	Gravity theories and holography	3
1.2	The string non-linear σ model	9
1.3	Aim and content of this thesis	16
2	The non-linear σ model and its symmetries	19
2.1	Bosonic strings in light-cone gauge	20
2.2	The $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ supercoset	24
2.3	Massive modes in light-cone gauge	29
2.4	Symmetry algebra and its representation	35
2.5	Chapter summary	41
3	The all-loop integrable S matrix	43
3.1	Classical and quantum integrability	43
3.2	Representations of $\mathfrak{su}(1 1)^2$ centrally extended	51
3.3	Finding the S matrix	57
3.4	Chapter summary	63
4	Crossing symmetry and dressing factors	65
4.1	Crossing symmetry	65
4.2	Solving the crossing equations	73
4.3	Poles of the S matrix	76
4.4	Chapter summary	82
5	The $\mathfrak{psu}(1,1 2)^2$ spin chain	83
5.1	The weakly-coupled spin chain	83
5.2	The dynamical spin chain	89
5.3	The spin-chain S matrix	95
5.4	Comparing with the worldsheet S matrix	99
5.5	Chapter summary	103

6	The all-loop Bethe ansatz equations	105
6.1	Bethe ansatz essentials	105
6.2	Nested Bethe ansatz	109
6.3	The $\mathfrak{psu}(1, 1 2)^2$ Bethe ansatz	117
6.4	Chapter summary	125
7	Comparison with perturbative calculations	127
7.1	Expansions of the dressing factors	127
7.2	Comparisons	131
7.3	Chapter summary	135
8	Recent developments and new directions	137
8.1	Massless modes in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$	137
8.2	Integrability for the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background	140
8.3	Mixed-flux backgrounds	146
8.4	Conclusions and outlook	149
A	Appendices	151
A.1	Generalities of the $\mathfrak{psu}(1, 1 2)$ superalgebra	151
A.2	Explicit expressions for the coset action	154
A.3	Quadratic charges in terms of the fields	157
A.4	Analytic properties of the dressing factors	159
A.5	Dualisation of the Bethe ansatz equations	163
	Samenvatting	167
	Curriculum vitae	171
	Acknowledgements	173
	References	175

1 Introduction

Quantum field theory (QFT) is a framework that originated from the necessity to formulate a quantum theory of electrodynamics, and which proved crucial in expanding our understanding of the world around us. We know that many essential features of strong, weak and electromagnetic interactions can be described using QFT, fitting together in the standard model (SM) of fundamental interactions. Moreover, QFTs can be employed to describe statistical ensembles of quantum particles, yielding a description of phenomena such as superconductivity, superfluidity, the quantum Hall effect, and so on.

There is even more: not only do we know that quantum field theory is the framework where many successful physical theories can be formulated, but it is in this framework that we can most easily understand, by means of the so-called renormalisation group, the existence of universality classes that allow us to describe a physical system through a relatively simple effective theory. For this point of view, QFTs are of crucial importance in that any relativistic quantum theory can be described like a QFT at low enough energy scales.

Despite the central role of QFT in modern Physics and its many successes, much is still to be understood. For instance, in realms where QFT is expected to be applicable, there are phenomena which we cannot currently explain satisfactorily, such as the mechanism behind high-temperature superconductivity, or the presence of a mass for SM neutrinos. Even in cases where we have a good microscopic formulation of a QFT, the strongly-coupled regime is hardly accessible to calculations; this is the case for quantum chromodynamics (QCD), where the details of the mechanism confining the quarks into hadrons still elude us. Indeed, for most cases of interest (and notably in four space-time dimensions) there is no rigorous mathematical construction of interacting QFTs at all—whether the coupling is small or large. We do have some insight into non-perturbative properties of QFTs (such as unitarity, invariance under symmetries, cluster decomposition, *etc.*) which can be a guiding principle in establishing properties such as the relation between spin and statistic, or offer a computational tool like unitarity cuts; still, our understanding of QFT is essentially perturbative, *i.e.* most of the computations are obtained from an asymptotic expansion in some parameter around a free theory.

Another very significant issue is how to include Gravity in such a picture. Of course

at sufficiently low energies QFT provides a good effective description of Gravity but it is not clear whether a fundamental theory of quantum Gravity could at all be a QFT. Even if several appealing proposals exist, none is completely satisfactory yet, so that many fundamental issues related to cosmology and to very high-energy Physics remain essentially unanswered.

There are several routes that can be explored to hopefully overcome these problems and improve our understanding of quantum theories. In some cases we can try to engineer specific QFTs, with special symmetries or particle content to better describe a certain phenomenon, in others it seems more appropriate—like it was done with string theory—to introduce a conceptually new framework. Further progress has been made in condensed matter and particle Physics by coming up with new computational techniques, which sometimes combine analytical tools with computer simulations. Finally, a somewhat less direct route is to focus on some special models that happen to be exactly solvable. The research contained in this thesis represents a step along this last route.

Clearly not all of the theories of our interest will be exactly solvable: believing so would be a step back towards a sort of Laplacian determinism. Nonetheless, they may offer insight in real-world models through universality, or when used as a theoretical laboratory. Exactly solvable models have always constituted an important testing and discovering ground for new ideas. In statistical mechanics, many subtleties on the thermodynamic limit were illustrated and sometimes understood by working on specific models. This is *e.g.* the case of the occurrence of phase transitions in the Lenz-Ising model and the role of boundary conditions in the six-vertex ice model. In hydrodynamics, the Korteweg-de Vries equation illustrated how a non-linear partial differential equation can admit stable, wave-like localised solutions—solitons. Integrable (quantum) spin-chains played and play an important role in condensed matter physics, as do one-dimensional gases of almost-free particles. QFTs in two space-time dimensions are indeed special. When they possess conformal symmetry, as it is the case for theories describing second-order phase transitions, they can be sometimes solved entirely by representation theory of the symmetry algebra. If they instead have well-defined asymptotic states, they have a chance of being exactly solvable owing to the factorisation of their S matrices—a mechanism highlighted by the Zamolodchikov brothers in 1978. In recent times, this was used to provide compelling evidence in favour of the holographic duality among $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) theory and type IIB superstring theory on the background spacetime $\text{AdS}_5 \times S^5$,¹ as we will briefly discuss later.

The subject of this thesis is precisely the investigation of some properties of certain superstring theories on $\text{AdS}_3 \times S^3 \times T^4$ and $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ backgrounds by treating them as two-dimensional QFTs and assuming that their S matrix does indeed factorise. Before

¹We will always denote the n -dimensional, negative-curvature maximally symmetric anti-de Sitter space as AdS_n , the n -dimensional sphere as S^n and the n -torus as T^n .

commencing that investigation, we would like to motivate the choice of such theories. This requires to discuss, at least qualitatively, some features of gravity theories and especially string theories on those backgrounds, with particular attention to the holographic duality.

1.1 Gravity theories and holography

General relativity provides a remarkably successful theory of classical gravity. However, promoting that to a quantum theory, or even fully understanding the classical theory and its physical solutions remains quite a formidable task. Significant progress in both directions has been made by restricting to lower dimensional models, using semiclassical techniques and focusing on specific solutions.

One of the first solutions to the equations of general relativity to be found were black holes, *i.e.* solutions presenting an event horizon such that an observer inside the horizon cannot communicate with the outside universe. It appears that black holes are not just a mathematical curiosity, but may indeed exist in our universe, representing the final stage of gravitational collapse. Since it is impossible to measure what lies inside a classical black hole, such an object is characterised in terms of a few charges: its mass M , angular momentum J and possibly electric charge. It was noted by Bekenstein [1], Bardeen, Carter and Hawking [2] that these charges obey relations reminiscent of thermodynamics. In particular, when considering a neutral stationary black hole and perturbing it slightly, one finds²

$$\delta M = \frac{\kappa}{8\pi} \delta A + \Omega \delta J, \quad \delta A \geq 0, \quad (1.1)$$

where κ is the surface gravity (which is constant on the horizon), A the horizon area, and Ω the angular velocity. By analogy with the first two laws of Thermodynamics, it appears that κ plays the role of temperature, and A of entropy. Furthermore, it was found by Hawking [3, 4] that semiclassically a black hole would indeed emit thermal radiation of temperature $T = \frac{1}{2\pi} \kappa$.

This seems to suggest that the entropy of a black hole is proportional to its area,

$$S = \frac{A}{4}, \quad (1.2)$$

in striking contradiction with entropy extensivity predicted by statistical mechanics. This may be considered not only as a characteristic of black hole solutions, but as a fundamental property of gravity [5], meaning that there is an upper bound to the amount of fundamental microscopic states that a given volume may contain, and the bound is saturated by black holes as in eq. (1.2). Bringing these ideas even further, 't Hooft advocated that the fundamental degrees of freedom in a D -dimensional theory containing gravity

²In units where $\hbar = G_N = c = 1$.




Figure 1.1: Two of the possible interactions of the string. The left panel depicts the emission of a closed string from two open ones, while the right panel features three closed strings. Notice that there is no localised interaction vertex.

live in $D - 1$ dimensions [6]. This would establish a relation between (quantum) gravity theories and quantum theories in one dimension less—the holographic duality³.

Since it requires working with gravity, finding explicit instances of such a duality is hard. However, in the simpler case of gravity in three spacetime dimensions with a negative cosmological constant, a remarkable result was found by Brown and Henneaux [8], who showed that the symmetry algebra of such a theory is given by two copies of the Virasoro algebra—the same algebra occurring in two-dimensional conformal quantum field theories (CFTs). This is particularly remarkable because the Virasoro algebra is infinite dimensional, and can be constraining enough to completely determine all of the observables of the CFT once the operator content and the central charges of the theory are established.

Gravity and holography from strings

Further progress in understanding holographic dualities has been made in the context of string theory [9] culminating in the AdS/CFT correspondence [10–12], see also ref. [13]. String theory is a quantum theory of interacting extended one-dimensional objects—strings. One may consider a theory of “closed” strings (that have the topology of a circle), or both “open” and “closed” strings (segments and circles). The strings propagate in a target space, spanning a two-dimensional surface called the worldsheet (in analogy with particles’ worldlines). Qualitatively,

$$\text{worldsheet} = \mathcal{W} \ni (\tau, \sigma) \mapsto X^\mu(\tau, \sigma) \in \mathcal{T} = \text{target space}, \quad (1.3)$$

³See also ref. [7] for a review of its early history and a more complete list of references.

where τ, σ are the time and the space coordinate on the worldsheet respectively, and \mathcal{T} is some space-time. The string can interact by joining and splitting and, as figure 1.1 shows, open strings can always yield close strings. In this framework, one finds out a few important facts:

- Excitations of the string can be thought of as particles in the target space \mathcal{T} . Requiring the spectrum of string excitations to include boson and fermions and be free of tachyons mandates the introduction of supersymmetry.
- Quantum-mechanical consistency of the theory yields a constraint on the dimension D of the target space \mathcal{T} . In a supersymmetric theory $D = 10$.
- The spectrum of any theory that contains closed strings presents a massless spin-two excitation—a candidate to be the graviton in the target space \mathcal{T} .
- Due to the strings being extended objects, string theories may be well-behaved at high energies—in fact better-behaved than a generic QFT.
- An appropriate low-energy limit of supersymmetric string theories yields supersymmetric QFTs of gravity—supergravities.

For these reasons, superstring theories can be regarded as quantum theories of gravity—even if not necessarily resembling real-world gravity. One can formulate theories where closed strings propagate in target spaces of the form $\mathcal{T} = \text{AdS}_{n+1} \times \mathcal{M}_{9-n}$, where \mathcal{M} is a manifold without boundary. The AdS/CFT correspondence is the statement that such theories are equivalent to certain QFTs, in the sense that in principle for each observable of the string theory there exists a QFT observable that takes precisely the same value (up to correctly identifying the coupling constants), and *vice versa*. Furthermore, such a QFT is in fact a conformal field theory in dimension n . In fact, AdS_{n+1} is a maximally symmetric negative-curvature space that can be realised as an embedded submanifold

$$-x_0^2 - x_1^2 + x_2^2 + \cdots + x_n^2 + x_{n+1}^2 = 1, \quad \vec{x} \in \mathbb{R}^{n+2}, \quad (1.4)$$

making manifest that its isometries form the pseudo-orthogonal group $SO(2, n)$ which is essentially the n -dimensional conformal group. This suggests a fascinating holographic picture where string theory on $\text{AdS}_{n+1} \times \mathcal{M}_{9-n}$ can be described in terms of a CFT on the n -dimensional flat boundary of AdS_{n+1} .

Of course the matching of symmetries is only a necessary condition for the duality to hold, and it is actually possible to put the duality on firmer grounds in a string theoretical framework. To this end one can show that the AdS geometry and the QFT description are two facets of the same stringy description. This is done by considering N solitonic objects called D-branes that can on the one hand deform the string background to an




Figure 1.2: Diagrammatic expansions in terms of the topology of the graphs. The first two panels exemplify the perturbation theory of Yang-Mills theories in terms of ribbon graphs. Panel (a) depicts a one-loop contribution to the two-point function that is planar, while the contribution in panel (b) cannot be drawn on a plane without self-intersections, and is subleading in a $1/N$ expansion. In panel (c), the genus expansion also appears naturally for closed strings, due to the interactions of figure 1.1b.

AdS geometry, and on the other hand naturally give rise to $SU(N)$ gauge theories, whose long-distance behaviour is described by a CFT.

A particularly interesting case is the one where N is *large* and the Yang-Mills coupling g_{YM} is scaled suitably. Already in 1973 it was noticed by 't Hooft [14] that one can formulate a diagrammatic expansion in $1/N$ while keeping fixed $\lambda = N g_{\text{YM}}^2$, in terms of the *topology* of the graphs. For instance, in the case of theories involving only adjoint $SU(N)$ fields, the diagrams involve double lines, yielding ribbons rather than lines for propagators. One can see that the leading terms come from planar diagrams, *i.e.* diagrams that can be drawn on a plane without self intersections, see figure 1.2. Diagrams that can be drawn with no intersection only on a surface of higher and higher genus will be more and more subleading in $1/N$, see figure 1.3. A similar genus expansion appears in string theory, where higher genus surfaces can be generated by joining and splitting of closed strings as in figure 1.2c. This suggests that when $N \rightarrow \infty$ the strings become free. This leaves still one parameter: the 't Hooft coupling λ in the gauge theory, or equivalently the string length (in units of the radius of AdS), ℓ_s/R . We would need a more detailed analysis to establish the precise relation between λ and ℓ_s/R . In general, one finds that the weakly coupled regime in one theory is the strongly coupled in the other⁴, *i.e.*

$$\lambda \ll 1 \Leftrightarrow \ell_s/R \gg 1 \quad \text{and} \quad \ell_s/R \ll 1 \Leftrightarrow \lambda \gg 1, \quad (1.5)$$

which is unsurprising since the perturbative string and gauge theories appear very different. Therefore, the holographic correspondence is a *weak-strong duality*. This is an

⁴Clearly in gauge theory the weakly coupled regime is $\lambda \ll 1$. For strings the non-trivial dynamics is due to the propagation in a curved space so that the weakly coupled regime is $\ell_s/R \ll 1$, when the target space is approximately flat.




Figure 1.3: The order in $1/N$ of a diagram is given by the genus of the simplest surface where it can be drawn without self-intersections. In panel (a), we can draw a ribbon graph on a sphere (genus zero), while in panel (b) we draw one on a torus (genus one). Note that these ribbon graphs have a large number of vertices and approximate well a smooth surface—a qualitative similarity between ribbon graphs and the string worldsheet.

appealing feature—it provides a new handle to manage strongly-coupled physics once the correspondence is established—but it makes it very hard to test whether the duality is truly there.

Let us be more precise. In the case of free strings, the natural observables are the string energy levels. These are the eigenvalues $\{E_j^{\text{t.s.}}\}$ of the generator $\mathbb{H}^{\text{t.s.}}$ of time-translations in the target space⁵. For some string states (such as the vacuum) supersymmetry may prescribe them to take a simple value. In general, however, they are a non-trivial function of ℓ_s/R , due to the fact that the free strings probe the curved $\text{AdS}_{n+1} \times \mathcal{M}_{9-n}$ geometry. In the large- N limit of the CFT, the leading observables are the two-point functions, whose form is constrained by conformal symmetry—schematically

$$\langle \mathcal{O}_j(x) \mathcal{O}_j(y) \rangle = \frac{1}{|x-y|^{2\Delta_j}}, \quad (1.6)$$

where Δ_j is the eigenvalue of the generator of dilatations \mathbb{D} acting on \mathcal{O}_j , and is in general a non-trivial function of λ . In fact, $\{\Delta_j\}$ is supposed to be dual to the string spectrum. Any perturbative calculation in the string theory will give $E_j^{\text{t.s.}}$ at $\ell_s/R \ll 1$, while in the CFT we would find Δ_j at $\lambda \ll 1$, making it impossible to compare the results in general. For this reason, significant effort has been invested in finding specific instances of the duality where exact results may be found for the whole spectrum.

Some instances of AdS/CFT

As we mentioned, the most well-studied case is the one of superstrings on $\text{AdS}_5 \times \text{S}^5$ and the dual $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) theory in four dimensions. This is nat-

⁵More precisely, this is when using so-called global coordinates for AdS.

ural since such string theory background preserves as much supersymmetry as possible, and it is interesting since the four-dimensional gauge theory, despite being conformal, may offer some insight on how to better understand QCD. Additionally, $\mathcal{N} = 4$ SYM can be easily studied in perturbation theory, which prompted the discovery of integrable structures at one loop [15] and beyond [16–18]. Similar structures appeared in classical [19–23] and quantum strings [24, 25]. Eventually, integrability was established as an all-loop feature of the planar $\text{AdS}_5/\text{CFT}_4$ duality [26–28], see also refs. [29, 30] for reviews and extensive lists of references. Shortly afterwards, the study was extended to strings on⁶ $\text{AdS}_4 \times \mathbb{CP}^3$ and the dual $\mathcal{N} = 6$ supersymmetric Chern-Simons theory in three dimensions [31] where integrable structures also appeared, see ref. [32] for a review.

Here we would like to focus on the maximally supersymmetric backgrounds involving AdS_3 spaces.⁷ Gravity on AdS_3 is a toy model, yet it retains interesting physical properties, such as black hole solutions as found by Bañados, Teitelboim and Zanelli (BTZ) [33]. Such BTZ solutions are locally isometric to empty AdS_3 and globally can be obtained by quotienting AdS_3 by a discrete group [34]. Therefore they are relatively simple, and can be used to investigate BH physics [35]. In fact, even if the “thermodynamics” of BHs is now well-established, there is still little understanding of the underlying “statistical mechanics”—if BHs have entropy, what are the microscopic states that give rise to it? It is much easier to answer this question in simpler settings, such as gravity on AdS_3 . There it was possible to match the area of a BTZ black hole of a given mass to, loosely speaking, the density of states of an appropriate energy in the dual CFT_2 , confirming the relation between area and entropy [36, 37]. To this end it was crucial to exploit the Virasoro symmetry of the CFT_2 [38]. Considering that BTZ-like geometries play a role also in higher-dimensional black hole solutions [39, 40], a deeper understanding of the $\text{AdS}_3/\text{CFT}_2$ duality is highly desirable.

Attempts in this directions were made already in Maldacena’s seminal paper [10]. In the case of type IIB strings on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, much progress can be made if one takes the background to be generated by a three-form flux of the Neveu-Schwarz-Neveu-Schwarz (NSNS) type, because then the worldsheet theory can be studied by CFT techniques [41–44]. The theory is dual to a CFT with chiral $(4, 4)$ extended supersymmetry on an orbifold of a product of T^4 tori [10, 41–46]. However this approach is much more difficult in presence of a Ramond-Ramond (RR) flux [47]. Furthermore, in the case of $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background, which has the same amount of supersymmetry, there is only limited insight on the dual CFT [48, 49]. On the other hand, it has recently been found that the string theory non-linear σ model (NLSM) is classically integrable for these

⁶Following standard conventions we denote by \mathbb{CP}^n the n -dimensional complex projective space, which can be regarded as a real quotient manifold $\text{S}^{2n+1}/U(1)$.

⁷These have 16 supersymmetries as opposed to the 32 of $\text{AdS}_5 \times \text{S}^5$ background and the 24 of $\text{AdS}_4 \times \mathbb{CP}^3$. At this stage, the requirement of extended supersymmetry can be taken as a technical one, likely to simplify the study.

backgrounds in presence of pure RR flux [50, 51], and even in presence of mixed RR and NSNS fluxes [52]. It is then natural to try and apply the S-matrix integrability approach that proved successful in $\text{AdS}_5 \times S^5$ and $\text{AdS}_4 \times \mathbb{CP}^3$. Such an investigation may also offer new insight on how integrability is related to the exact solvability in terms of Virasoro representations, and on the properties of integrable S matrices involving massless particles—which here arise due to the presence of flat directions in the target space [50].

This program is ambitious, and in this thesis we will describe the initial steps that have been undertaken to complete it. Our approach will be based on the S matrix of worldsheet excitations for the gauge-fixed string NLSM, as we will outline in the next section.

1.2 The string non-linear σ model

Let us consider a theory of closed (super)strings only. In absence of string interactions, the worldsheet of the string is a cylinder, and the classical string theory is defined by an action of the form

$$S_{\text{bos}} = -\frac{h}{2} \int_{\mathcal{W}} d\tau d\sigma \sqrt{|\gamma|} \gamma^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu G_{\mu\nu}(X), \quad (1.7)$$

for the bosons, which should be supplemented by fermionic terms—here we avoid doing so to keep the discussion simple. Here $h \approx R^2/\ell_s^2$ is a coupling constant, $\gamma_{\alpha\beta}$ is the metric on the worldsheet, X^μ can be thought of as coordinates in the target space, whose metric is $G_{\mu\nu}$. For a superstring theory, \mathcal{T} must be ten-dimensional, so that $\mu = 0, \dots, 9$.

It is easy to see that eq. (1.7) is invariant under reparametrisations of the worldsheet. As a result, to understand what the physical degrees of freedom of the theory are we need to fix this local invariance. We have enough freedom to take the worldsheet metric to be flat, $\gamma_{\alpha\beta} \rightarrow \eta_{\alpha\beta}$. Then, the action in (1.7) becomes the one of a NLSM, from a cylinder into a target space with (fixed) metric $G_{\mu\nu}$, subject to the additional constraints

$$\left. \frac{\delta S_{\text{bos}}}{\delta \gamma^{\alpha\beta}} \right|_{\gamma_{\alpha\beta}=\eta_{\alpha\beta}} = 0. \quad (1.8)$$

However, we have not exhausted the reparametrisation invariance of the string⁸, due to the fact that the action (1.7) also possesses a local invariance under Weyl rescaling. In fact, let us assume for the moment that the target space is flat, $G_{\mu\nu} = \eta_{\mu\nu}$. Then, if we introduce light-cone coordinates

$$X_\pm = X^0 \pm X^9, \quad (1.9)$$

⁸This residual invariance is somewhat similar to the one appearing, *e.g.*, when considering the covariant gauge in electrodynamics: there imposing $\partial_\mu A^\mu = 0$ leaves the possibility of making gauge transformation $A_\mu \rightarrow A_\mu + \partial_\mu \lambda$ provided that $\partial_\mu \partial^\mu \lambda = 0$. Only removing all gauge invariance allows to go from the four polarisations of A^μ to the two physical polarisations of the photon.

it is easy to see that the residual invariance is enough to set

$$X_+(\sigma, \tau) = x_+ + \tau. \quad (1.10)$$

In this way, one can use the constraint (1.8) to solve for X^- in terms of the remaining fields. This still leaves one non-linear constraint, the so-called *level-matching condition*, which enforces periodicity of the strings along σ and amounts to the vanishing of the worldsheet momentum \mathbb{P} .⁹

This gauge-fixed theory of free strings (or rather, a suitable supersymmetric version on certain curved backgrounds) is what we will be considering in this thesis. In the bosonic sector we are left with eight physical fields X^j defined on a cylinder. The physical Hilbert space will consist of the excitation of these eight fields subject to the level-matching condition, which is realised as a projection on the Hilbert space

$$\mathbb{P} \{ \{M_1\}, \dots, \{M_8\} \} = 0, \quad (1.11)$$

where $\{M_j\}$ are label the excitations of each field. A preferred basis is the one of eigenstates of the worldsheet Hamiltonian \mathbb{H} , that is the operator generating time evolution on the worldsheet in the sense of Stone's theorem. The spectrum we eventually want to compute is the one corresponding to time evolution *in the target space*. However, by light-cone gauge fixing we have related the worldsheet time τ to X_+ and hence to X^0 , that is the target-space time. From this it follows that the eigenvalues of $\mathbb{H}^{\text{t.s.}}$ are simply related to the ones of \mathbb{H} , and it will be enough to compute the latter.

It is worth recalling that the spectrum of physical states will organise itself into multiplets of the symmetry algebra of the theory. Even if we did not construct the string Hamiltonian, it is easy to see that in the flat case the physical fields X^j , $j = 1, \dots, 8$ in (1.7) are invariant under the action of $SO(8)$, which is the little group of massless particles in $D = 10$. This is the *manifest* symmetry of the theory in light-cone gauge. However, it is easy to see that the theory contains infinitely many states that can be interpreted as massive particles in $D = 10$. Since the little group of such excitation is $SO(9)$, it follows that the theory should enjoy a larger symmetry, and that the $\mathfrak{so}(8)$ Lie algebra¹⁰ multiplets should arrange themselves into irreducible representations of $\mathfrak{so}(9)$. This illustrates how in general the manifest symmetries of the light-cone gauge-fixed theory form a subalgebra of the whole symmetry algebra, which in fact for flat space should be the full $\mathfrak{so}(1,9)$.

In practice, for curved supersymmetric backgrounds there will be several complications: the action will involve fermions and non-linear terms. An useful approach is to rewrite the action (or part of it) as a coset action of a suitable supergroup, following what Metsaev and Tseytlin did in the case of $\text{AdS}_5 \times S^5$ [53]. Still, the gauge fixed Hamiltonian

⁹Here for simplicity we only consider string configurations with no winding.

¹⁰We will denote Lie algebras and superalgebras in Gothic letters.

will be highly non-linear [54–56], so that only perturbative quantisation is possible, see also ref. [29]. A way around this complication—valid as long as we can identify asymptotic states—is to find a (hopefully unique) S matrix that preserves the symmetries of the theory, and use that to find the spectrum, in the spirit of refs. [57, 58].

The decompactification limit and the worldsheet S matrix

In the light-cone gauge, the worldsheet is no longer invariant under rescalings. In fact, a more careful analysis would reveal that the size ℓ of the worldsheet is fixed in terms of the momentum conjugated to the light-cone coordinate X_- . It is interesting to consider the limit $\ell \rightarrow \infty$, whereby the worldsheet cylinder decompactifies into a plane. In this case, we are dealing with a two-dimensional QFT with well-defined asymptotic states. In particular, the spectrum can be described in terms of M -particle states on the worldsheet created by raising operators from a vacuum,

$$|p_1, \dots, p_n\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in}, \text{out})} = a_{\alpha_1}^{\dagger(\text{in}, \text{out})}(p_1) \cdots a_{\alpha_n}^{\dagger(\text{in}, \text{out})}(p_n) |0\rangle, \quad (1.12)$$

where the in and out raising operators $a^{\dagger(\text{in}, \text{out})}$ satisfy canonical commutation relations with the in and out lowering operators $a^{(\text{in}, \text{out})}$. If we want to consider physical states, we will have to impose the level matching condition,

$$\mathbb{P} |p_1, \dots, p_n\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in}, \text{out})} = (p_1 + \cdots + p_n) |p_1, \dots, p_n\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in}, \text{out})} = 0. \quad (1.13)$$

We will call a state satisfying eq. (1.13) “on-shell”, as opposed to a generic (off-shell) state. In either case, the action of the Hamiltonian is very simple

$$\begin{aligned} \mathbb{H}^{(\text{in}, \text{out})} |p_1, \dots, p_n\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in}, \text{out})} &= E(\{\alpha_j, p_j\}) |p_1, \dots, p_n\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in}, \text{out})}, \\ E(\{\alpha_i, p_i\}) &= \sum_{j=1}^M \omega_{\alpha_j}(p_j), \end{aligned} \quad (1.14)$$

where $\omega_{\alpha}(p)$ is the dispersion relation. In the case of a relativistic theory we should find

$$\omega_{\alpha}(p) = \sqrt{m_{\alpha}^2 + p^2}. \quad (1.15)$$

However, for AdS background it has been found that light-cone gauge fixing breaks the relativistic invariance on the worldsheet so that the dispersion relation takes a lattice-like form [56, 59]

$$\omega_{\alpha}(p) = \sqrt{m_{\alpha}^2 + 4h^2 \sin^2 \frac{p}{2}}, \quad (1.16)$$

where h is the coupling constant.

Given that the two sets of raising and lowering operators $\{a^{\dagger(\text{in})}, a^{(\text{in})}\}$ and $\{a^{\dagger(\text{out})}, a^{(\text{out})}\}$ both satisfy canonical commutation relations, by virtue of the Stone-von Neumann theorem they must be related by a unitary operator \mathbb{S} satisfying

$$\mathbb{S}^{\dagger} \mathbb{S} = \mathbb{S} \mathbb{S}^{\dagger} = \mathbb{I}, \quad \mathbb{S} |0\rangle = |0\rangle, \quad a^{\dagger(\text{in})} = \mathbb{S} a^{\dagger(\text{out})} \mathbb{S}^{\dagger}, \quad a^{(\text{in})} = \mathbb{S} a^{(\text{out})} \mathbb{S}^{\dagger}, \quad (1.17)$$

from which one immediately finds that \mathbb{S} is the familiar S matrix that relates in- and out-states,

$$|p_1, \dots, p_n\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in})} = \mathbb{S} |\tilde{p}_1, \dots, \tilde{p}_M\rangle_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_M}^{(\text{out})}. \quad (1.18)$$

In practice, finding the S matrix is hard, and it can be done perturbatively only at one- or two-loop order for the models of our interest. We will circumvent this problem by dealing with theories whose S matrix can be determined by the symmetries of the theory—that is, *integrable theories*.¹¹

Equipped with the S matrix and dispersion relation, and remembering that we are dealing with a QFT in 1+1 dimensions, we can start looking at some observables. Let us prepare a one-particle state $|p\rangle_\alpha^{(\text{in})}$ of definite momentum and flavour. In absence of external fields such a stable asymptotic state will satisfy $|p\rangle_\alpha^{(\text{in})} = |p\rangle_\alpha^{(\text{out})}$ and its energy will be

$$E = \omega_\alpha(p), \quad (1.19)$$

so that the spectrum is continuous. If, however, we take into account the fact that the worldsheet is a cylinder of size ℓ , we should also impose the spatial periodicity of the wave-function, which amounts to the quantisation condition for the momentum

$$e^{ip\ell} = 1, \quad (1.20)$$

resulting in a *discrete* spectrum. Let us now take a state $|p, q\rangle_{\alpha, \alpha}^{(\text{in})}$ consisting of two particles of the same flavour α , with momenta $p > q$ so that they are asymptotically well separated. Let us also assume that they scatter elastically without producing any other particle—which is generally not the case in a QFT. Then the corresponding out-state will contain again two particles of the same flavour and momenta p, q . If we now impose periodicity, however, we have to account for the fact that each particle underwent a phase shift due to the scattering, so that we have

$$e^{ip\ell} S(p, q) = 1, \quad e^{iq\ell} S(q, p) = 1, \quad (1.21)$$

where $S(p, q)$ is the diagonal matrix element for the flavour α , $S_{\alpha\alpha}^{\alpha\alpha}(p, q)$. Inserting the solutions for p, q into the dispersion relations (1.14) will yield again a discrete energy spectrum. Since we are interested in the *physical* spectrum, we must impose the level matching condition (1.13), finding that there are no non-trivial one-particle states in the on shell theory, and that for two particles one must have $q = -p$. If one were able to follow a similar recipe for any number of particles of arbitrary flavours, then he would have a description of the string spectrum. It turns out that this is possible provided that \mathbb{S} “factorises”, *i.e.* provided that a M -body scattering event can be understood as a sequence of two-body ones. Again, this is a typical feature of integrable theories. Then,

¹¹ We will discuss at length the notion of integrability in chapter 3.

the M -particle analogue of eq. (1.21) are the *Bethe-Yang equations* (BY equations), that are schematically of the form

$$e^{i p_k \ell} \prod_{j \neq k}^M S(p_k, p_j) = 1 \quad \text{for} \quad k = 1, \dots, M, \quad (1.22)$$

where $S(p, q)$ is a suitable S-matrix element. Equations of this type were first found in the context of QFTs by Yang [60], inspired by the ansatz that Bethe proposed in the context of quantum spin chains [61].

A spin-chain picture

The appearance of the periodic dispersion relation (1.16) and the fact that the spectrum can be described in terms equations of the Bethe ansatz type strongly hint that the underlying theory may have an alternative description in terms of a discrete model, perhaps of a quantum spin chain.

It has been long known that some NLSMs are equivalent to certain quantum spin chains, see *e.g.* chapter 5 in ref. [62]. For instance, a coset model similar to the one emerging from string theory but with target space S^2 is equivalent to the long wavelength limit of the Heisenberg spin chain. The Heisenberg chain is perhaps the prototype of a quantum spin chain, defined on ℓ sites periodically identified and with Hamiltonian

$$\mathbb{H} = \mu \sum_{j=1}^{\ell} \left(\frac{1}{4} - \vec{S}_j \cdot \vec{S}_{j+1} \right), \quad (1.23)$$

where \vec{S}_j is a spin at the j th site and μ is a coupling constant.

The sphere S^2 may be part of the target space for our string NLSM, *e.g.* in the case of $\text{AdS}_3 \times S^3 \times \mathcal{M}_4$. Therefore the spectrum of the Heisenberg chain could somehow be part of the one that we wish to compute. For several integrable strings backgrounds this is actually the case [63, 64], and for instance in the case of $\text{AdS}_5/\text{CFT}_4$ the first evidence of integrability was discovered in 4-dimensional $\mathcal{N} = 4$ SYM theory precisely in form of the Heisenberg spin chain Hamiltonian [15]. Let us briefly review that picture.

The operator dual to the string Hamiltonian is proportional to the generator of dilatations \mathbb{D} in the CFT. String states are dual to local operators which in the large N limit take the form *e.g.*

$$\mathcal{O} = \text{tr} [Z Z \partial^\mu X \partial_\mu Z \psi \cdots \bar{\psi} Z], \quad (1.24)$$

where $X, Z, \psi, \bar{\psi}$ are some of the (bosonic and fermionic) fundamental fields appearing in the CFT_n Lagrangian, which are all evaluated at the same spacetime point, and the trace ensures gauge invariance. The spectral problem now reads

$$\mathbb{D} \mathcal{O} = \Delta \mathcal{O}. \quad (1.25)$$

Each of the fields that compose \mathcal{O} are identified by their transformation properties under the n -dimensional superconformal algebra. In fact, we can think of them as of “spins” of that algebra, so that \mathcal{O} becomes a state of the *periodic* (due to the trace) spin chain.

In particular, in the case of $\mathcal{N} = 4$ SYM, if we restrict to two $\mathfrak{su}(2)$ -charged scalar fields X, Z , we have that an operator \mathcal{O} is equivalent to a state $|\Psi\rangle$

$$\mathcal{O} = \text{tr}[Z Z X Z X \cdots X Z] \quad \longleftrightarrow \quad |\Psi\rangle = |\downarrow\downarrow\uparrow\downarrow\uparrow \cdots \uparrow\downarrow\rangle, \quad (1.26)$$

where the arrows indicate $\mathfrak{su}(2)$ spins, $S^3|\downarrow\rangle = -\frac{1}{2}|\downarrow\rangle$ and $S^3|\uparrow\rangle = +\frac{1}{2}|\uparrow\rangle$. The breakthrough of ref. [15] was realising that at 1-loop in the weakly coupled CFT, the dilatation operator \mathbb{D} coincides with \mathbb{H} of (1.23) up to suitably identifying the coupling constants. This is particularly remarkable because \mathbb{H} enjoys a large number of symmetries that make the explicit solution of the spectral problem possible in terms of a Bethe ansatz.

As it turns out, the whole spectral problem of $\mathcal{N} = 4$ SYM can be related to a spin chain for the superconformal algebra, and solved by considering an asymptotic *spin-chain S matrix* [27, 65], in a procedure that strongly resembles the one described for the string worldsheet theory. The first step is to consider the limit of a long spin chain, $\ell \rightarrow \infty$ and effectively decompactify the chain¹². We then consider collective excitations of definite momentum, called *spin waves* or *magnons* of the form *e.g.*

$$|p\rangle = \sum_{j=1}^{\infty} e^{ipj} |\downarrow\downarrow\downarrow\downarrow \cdots \downarrow\downarrow\uparrow\downarrow\downarrow \cdots\rangle, \quad (1.27)$$

where the overturned spin sits at the j th site. Notice that we have implicitly picked a *vacuum state* of downward spins. This operation is akin to the light-cone gauge fixing and breaks the manifest symmetry of the model. Remarkably, it still may happen that the S matrix corresponding to the scattering of two magnons is uniquely fixed by the residual symmetries, and the multiparticle scattering “factorises” in the sense alluded to earlier. In that case, we can proceed to write down *asymptotic* Bethe ansatz equations of the form (1.22) by requiring that M magnons live on a periodic spin chain of finite length ℓ . Finally, requiring that the states are invariant under cyclic permutations precisely reproduces the level-matching constraint (1.13), as we will see in chapter 6.

The above picture can be seen to hold for several instances of AdS/CFT. However, in the cases where a Lagrangian description of the CFT cannot be efficiently used, as it happens in AdS₃/CFT₂, it is less clear how the spin chain emerges and how to relate it to the CFT observables. Still, if such a description exists it may be used as a tool to investigate the CFT side of the duality, as well as to give an alternative albeit similar way

¹²In simpler cases such as the Heisenberg chain we might not need to first consider the $\ell \rightarrow \infty$ limit, and the Bethe ansatz would be exact for any ℓ . However, one peculiarity of the spin chains arising from AdS/CFT is that the Hamiltonian may couple states of different length—in the case of $\mathcal{N} = 4$ SYM this can be seen from the presence of a Yukawa interaction in the SYM Lagrangian. Then the Bethe ansatz description is only asymptotic, *i.e.* valid for $\ell \rightarrow \infty$.




Figure 1.4: Two processes involving exchange of real or virtual particles that wrap around the worldsheet cylinder, and therefore are not captured by the Bethe-Yang equations.

to solve the spectral problem. One may speculate that a spin-chain description is always possible simply by considering a discretisation of the string worldsheet [66, 67]. However it is unknown whether such a discretisation is natural for any general string theory in light-cone gauge on AdS. In fact, very recently for AdS₃/CFT₂ there appeared evidence that this may not be the case [67], as we will describe in the last chapter of this thesis.

The spectrum of the finite-size theory

The Bethe-Yang equations (1.22) do *not* describe the spectrum of the finite-size AdS/CFT duality [68]. They rather describe the spectrum of a QFT defined on a plane, after periodic identification of its spatial direction. To appreciate the difference between the two cases, let us consider the processes of figure 1.4. There we depict the propagation of (possibly virtual) particles wrapping the worldsheet cylinder, which cannot be accounted for by the S matrix derived in the decompactified theory. In terms of the spin chain, a similar effect arises due to the presence of long-range interactions—eventually, for a finite-length chain, such interactions wrap around the chain, invalidating the asymptotic Bethe ansatz approach.

In the QFT context, it is possible to estimate that wrapping effects are exponentially suppressed when ℓ is large, but nonetheless they are to be taken into account. One possibility, pioneered by Lüscher [69, 70], is to treat them as (perturbative) corrections to the spectrum predicted by the BY equations. Remarkably, Bajnok and Janik extended that approach to the AdS₅ × S⁵ NLSM providing further evidence of the correspondence [71].

One can do even better by fully exploiting the integrability properties of \mathbb{S} and the fact that, while directly dealing with (integrable) S matrices in finite volume is even hardly self-consistent, doing so at finite temperature is quite natural. The thermodynamic Bethe ansatz (TBA) is a tool to compute the free energy of finite-temperature integrable

QFTs [72]. One describes the thermal bath of particles in the grand canonical ensemble by taking into account all the particles appearing in the BY equations and their bound states, and requiring thermodynamic equilibrium. This means that the number of excitations is large, and their densities are fixed. Consequently, starting from the logarithm of eq. (1.22) one then obtains a set of non-linear coupled integral equations¹³.

Zamolodchikov realised that this can be used to find the ground-state energy of a finite-size theory from the free energy of a finite-temperature one [73], provided that these are related by exchanging the role of time and space by two Wick rotations—a “mirror” transformation. This procedure does not introduce any additional complication in the case of a relativistic theory, but is not straightforward in non-relativistic models. This can be seen by considering the effects of the mirror transformation

$$p \mapsto -iE = p_{\text{mirror}}, \quad E \mapsto -ip = E_{\text{mirror}}, \quad (1.28)$$

on the dispersion relation $E^2 = \omega_j(p)^2$, which leaves invariant the relativistic case (1.15) yields an entirely new relation when one uses eq. (1.16). In fact, the mirror transformation produces a novel *mirror theory*. Fortunately, even if many physical properties (such as the bound-state spectrum) differ in the string and mirror theories, their S matrices are related by analytic continuation, so that the information obtained by integrability carries over to the mirror theory [74, 75]. In the case of $\text{AdS}_5 \times \text{S}^5$, this allowed to write down a system of *mirror TBA* integral equations [76–79], or equivalently a “Y system” of functional equations [80–82], then simplified to a finite set of non-linear integral equations [83–85], from which in principle one can find the finite volume spectrum of the theory for any value of the ’t Hooft coupling. All these descriptions by construction yield a spectrum organised in multiplets of the superconformal algebra [86], for which finite-size effects are essential [S9]. The study of spectroscopy for $\text{AdS}_5/\text{CFT}_4$ by either analytical [87–92], [S8] or entirely numerical methods [93–95] has been initiated and provided substantial evidence in favour of the holographic duality.

A similar progress has been achieved also in the case of $\text{AdS}_4/\text{CFT}_3$, see refs. [32, 96]. As we will review, in the case of $\text{AdS}_3/\text{CFT}_2$, the understanding of the spectral problem is not quite as advanced, and the mirror TBA description is yet to be formulated—but hopefully not so far out of reach.

1.3 Aim and content of this thesis

This thesis aims at reviewing the recent progress in understanding the string worldsheet S matrices that arise in the context of the $\text{AdS}_3/\text{CFT}_2$ correspondence, focusing on the

¹³More precisely, one obtains such an integral equation for each species of particle configurations appearing in the thermodynamic limit. Identifying those usually requires some assumptions that go under the name of string hypothesis.

integrability approach.

We intend to offer a presentation as self-contained and accessible as possible. In view of the growing size of this field, we felt it was necessary to make a choice of topics: we will study in quite some detail the S matrix of massive excitations of superstrings in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ supported by pure Ramond-Ramond flux. This will be done both in terms of the superstring worldsheet theory and of a dual spin chain. Armed with a precise understanding of this simpler and prototypical case, we will devote the last chapter of this thesis to reviewing related topics, open issues and yet to be explored directions.

In chapter 2 we will discuss the worldsheet theory of free strings in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ in terms of a coset NLSM. We will focus on the massive excitations and derive their symmetries. This previously unpublished study follows closely ideas that were first developed for $\text{AdS}_5 \times \text{S}^5$ [56, 57], but highlights some novel unexpected features of the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ background. Some of the results presented here have also been found, among other things, in ref. [S1] by different techniques.

In chapter 3 we will review the main ideas behind integrability, and in particular the scattering factorisation, which ensures that any scattering process can be written in terms of a sequence of $2 \rightarrow 2$ scatterings. Under the assumptions of integrability, we will use the symmetries found in the previous chapter to find the two-body S matrix. This can be determined up to two scalar functions—the dressing factors—and satisfies several non-trivial consistency checks, most notably the Yang-Baxter equation. The results of this chapter were first found in [S3, S5], while the presentation follows [29, 58].

Chapter 4 is devoted to the study of crossing symmetry, and based on ref. [S2]. Firstly we will discuss crossing invariance for worldsheet excitations, which will be a generalisation of the familiar relativistic one [97]. Then we will propose crossing-invariant dressing factors for the S matrix found in the previous chapter and discuss some of their analytic properties. In particular we will show that our choice is compatible with the expected massive bound-state spectrum of the theory. This is a relevant check because crossing symmetry alone does not fix the form of the dressing factors completely.

We will switch gears in chapter 5, and introduce a spin-chain picture, constructed in such a way as to be dual to the worldsheet theory of the preceding chapters. Again the focus will be on its symmetries, and it will lead us to a two-magnon S matrix. We will relate it in a precise way to the worldsheet S matrix, and discuss the notion analogue to crossing in the spin-chain picture. This chapter follows more closely the original approach [S3, S5] in which the S matrix was found.

In chapter 6 we will make a first step from the decompactified theory towards a finite-size, albeit asymptotic, one. To this end, we introduce in quite some detail the asymptotic “coordinate” Bethe ansatz for both the spin-chain and worldsheet pictures. While the former case was worked out in refs. [S3, S4], the latter is an original result of this thesis. Since we will be dealing with non-diagonal S matrices, we will need to use the nesting

procedure in order to write the Bethe equations. As we will show, the Bethe ansatz in the spin-chain and worldsheet picture describe the same physical spectrum.

In chapter 7 we will put our construction to the test. Up to that point we assumed the worldsheet theory to be integrable at the quantum level, and derived an S matrix based on that assumption. We will check that S matrix, including the proposed dressing factors, against perturbative and semiclassical calculations in the worldsheet theory [50, 98–108] up to one-loop and including a non-trivial two-loop consistency check, finding complete agreement. Unfortunately it is much harder to perform such comparisons with the perturbative expansion of the dual CFT₂ [109]. In that regime, some puzzles remain.

In chapter 8, the final chapter, we will overview several directions in which the topics of this thesis can be and are being evolved. The most natural one is the inclusion of the massless fundamental excitations to the integrability picture, which was initiated in [110, 111] and more recently addressed in [S1]. Another is the extension of integrability to the other maximally supersymmetric AdS₃ background, *i.e.* AdS₃ × S³ × S³ × S¹ [112]. For that case, the all-loop massive S matrix and Bethe ansatz was proposed in [S4, S5] up to the dressing factors, but several open questions remain—notably what happens in the limit where one of the spheres blows up to give back AdS₃ × S³ × T⁴, up to a compactification. Finally, after the recent proof of their classical integrability [52], there has been progress in studying quantum integrability for backgrounds supported by a mixture of Ramond-Ramond and Neveu-Schwarz-Neveu-Schwarz fluxes [66, 67, 102], which may be a new playground to further our understanding of AdS/CFT integrability.

A note on notation

Most of the results collected in this thesis have already appeared elsewhere [S1–S5]. To streamline our presentation, our notations here may differ from some of the original ones. Furthermore, for the same reason we sometimes use slightly different convention and normalisations—most notably for the S matrix, which after all is a matrix and as such depends on the choice of basis. We also should warn the reader that some quantities that appear both in the worldsheet picture and in the spin-chain one (such as the S matrix, the generators of the symmetry algebra, *etc.*) will be indicated by the same letter, *despite not having the same value in the two cases*. We do so because these objects play the very same role and are *almost* identical in the two pictures, and because we feel that introducing different notations for each of them would be an unnecessary burden. When confusion may arise, we do clarify in the text what quantities we are referring to.

2 The non-linear σ model and its symmetries

In this chapter we will analyse in more detail some features of the NLSM on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$. In particular, we will write down its gauge-fixed action, restricting to massive excitations for simplicity. Out of that we will work out the symmetries of the theory, which will be important in the following chapters.

As anticipated in the introduction, we want to study our theory in the light-cone gauge. However, the technique to do so will be different and slightly more involved than the one sketched there. In fact, in AdS backgrounds it is impossible to impose conformal and light-cone gauge on top of each other [54, 113]. The way around this issue is to use first-order formalism, which has the further advantage naturally producing the Hamiltonian, whose eigenvalues are what we are after.

Firstly, we will demonstrate this approach in the case of a bosonic NLSM on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$. This will allow us to describe the strategy in some detail, without dealing with the complications due to fermions. In order to accommodate these, we could take two routes

1. Consider the Green-Schwarz (GS) action for the superstring [114, 115]. This is known up to quartic order in the fermions [116], which would give the complete action for the case of our interest.
2. Following [50], write the superstring action as a coset action

$$\frac{PSU(1, 1|2) \times PSU(1, 1|2)}{SU(1|1) \times SU(2)} \times U(1)^4. \quad (2.1)$$

The former method is completely general, but makes it harder to see how the isometries are realised. This is instead manifest in the coset formulation, which however requires a specific choice of which fermions we consider physical (the κ -gauge fixing). This choice is not suitable for studying massive and massless excitations at the same time. Since our focus will be on the massive excitations only, we will use the coset action, and work out its symmetries. As we will discuss in chapter 8, the GS approach has been used in ref. [S1] precisely to understand the role played by the massless excitations.

2.1 Bosonic strings in light-cone gauge

To exemplify the procedure we will follow later, let us first consider a bosonic NLSM action for closed strings, of the form

$$S = -\frac{h}{2} \int_{-\ell/2}^{\ell/2} d\tau d\sigma \gamma^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu G_{\mu\nu}(X), \quad (2.2)$$

where we consistently set to zero the Fradkin-Tseytlin $R_{(2)}$ term. Notice further that we replaced the worldsheet metric by the conformally invariant combination $\gamma^{\alpha\beta} \sqrt{|\gamma|} \rightarrow \gamma^{\alpha\beta}$.

Let us introduce the conjugate momenta

$$p_\mu = \frac{\delta S}{\delta \dot{X}^\mu} = -h \gamma^{0\beta} \partial_\beta X^\nu G_{\mu\nu}(X), \quad (2.3)$$

where $\dot{X}^\mu = \partial_0 X^\mu$. Then the action can be rewritten as [54, 113]

$$S = \int_{-\ell/2}^{\ell/2} d^2\sigma \left(p_\mu \dot{X}^\mu + \frac{\gamma^{01}}{\gamma^{00}} \mathcal{C}_1 + \frac{1}{2h \gamma^{00}} \mathcal{C}_2 \right), \quad (2.4)$$

where

$$\mathcal{C}_1 = p_\mu X'^\mu, \quad \mathcal{C}_2 = G^{\mu\nu} p_\mu p_\nu + h^2 G_{\mu\nu} X'^\mu X'^\nu, \quad (2.5)$$

with $X'^\mu = \partial_1 X^\mu$. The action (2.4) is no longer manifestly covariant on the worldsheet. However, due to the Virasoro constraints we have that it must be

$$\mathcal{C}_1 = 0, \quad \mathcal{C}_2 = 0, \quad (2.6)$$

so that the $\gamma^{\alpha\beta}$ plays the role of a Lagrange multiplier and will not appear in the Hamiltonian after (2.6) is imposed.

Let us specialise to the background $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$. Let t be the time-coordinate in AdS_3 and ϕ be an angle in S^3 . Translations and rotations in these directions are isometries, to which correspond two conserved charges

$$\mathbb{H}^{\text{t.s.}} = - \int_{-\ell/2}^{\ell/2} d\sigma p_t, \quad \mathbb{J} = \int_{-\ell/2}^{\ell/2} d\sigma p_\phi, \quad (2.7)$$

where p_t, p_ϕ are conjugate momenta. The target space energy $\mathbb{H}^{\text{t.s.}}$ is what, after quantisation, will give us the energy spectrum we are interested in, and \mathbb{J} is a distinguished angular momentum. Out of these two coordinates, we can construct light-cone ones [117]¹

$$x_- = \phi - t, \quad x_+ = \frac{1}{2}(\phi + t), \quad (2.8)$$

and denote by x^i the remaining coordinates. The light-cone gauge-fixing condition is then

$$x_+ = \tau, \quad p_+ = 1, \quad (2.9)$$

¹In fact, a more general coordinate choice is possible [55, 118], where $x_+ = a\phi + (1-a)t$.

where we assumed that there is no winding.² We will always restrict to this case, which is the only one where the large-tension expansion of the string is well-defined. Our coordinate choice implies that the light-cone momentum is $p_+ = \frac{1}{2}(p_\phi - p_t)$, so that the corresponding Noether charge is $\mathbb{P}_+ = \frac{1}{2}(\mathbb{J} + \mathbb{H}^{\text{t.s.}})$. In addition, the eigenvalue P_+ of \mathbb{P}_+ is

$$P_+ = \int_{-\ell/2}^{\ell/2} d\sigma p_+ = \ell \quad \implies \quad \ell = \frac{1}{2}(J + E^{\text{t.s.}}). \quad (2.10)$$

This shows explicitly how in light-cone gauge, invariance under worldsheet rescaling is lost, and in fact the worldsheet radius is fixed in term of physical charges. Moreover, relating t with τ will allow us to compute the target space energy $E^{\text{t.s.}}$ in terms of the worldsheet Hamiltonian.

One of the advantages of light-cone gauge fixing is that spurious degrees of freedom can be eliminated, which makes quantisation easier. In particular, we can get rid of the metric $\gamma^{\alpha\beta}$ by a suitable gauge fixing, whose precise form is irrelevant in the first-order formalism, and eliminate the longitudinal modes of the string x_\pm, p_\pm . To this end we solve $\mathcal{C}_1 = 0$ for x'_-

$$x'_- = -p_j x'^k. \quad (2.11)$$

Then, p_- can be eliminated by solving the non-linear constraint $\mathcal{C}_2 = 0$. We have, plugging in the light-cone gauge conditions

$$\begin{aligned} \mathcal{C}_2 = & \frac{1}{4}(G^{\phi\phi} - G^{tt})(p_-^2 + 4) + (G^{\phi\phi} + G^{tt})p_- + \frac{h^2}{4}(G_{\phi\phi} - G_{tt})x_-'^2 \\ & + G^{jk}p_j p_k + h^2 G_{jk}x'^j x'^k, \end{aligned} \quad (2.12)$$

where $G_{tt}, G_{\phi\phi}, G_{jk}$ are metric elements and x'_- is a function of the transverse fields as in eq. (2.11). Formally inverting the constraint equation gives

$$p_- = p_-(x^j, x'^j, p_j). \quad (2.13)$$

Plugging these expressions into the action (2.4) and dropping the total time derivative \dot{x}_- , we find

$$S = \int_{-\ell/2}^{\ell/2} d^2\sigma (p_j \dot{x}^j + p_-(x^j, x'^j, p_j)), \quad (2.14)$$

whence we can immediately identify the worldsheet Hamiltonian density

$$\mathcal{H} = -p_-(x^j, x'^j, p_j), \quad (2.15)$$

and the Poisson structure

$$\{x^k(\sigma), p_j(\tilde{\sigma})\} = \delta_j^k \delta(\sigma - \tilde{\sigma}). \quad (2.16)$$

²One can account for a winding of the form $x_+(r) - x_+(-r) = 2\pi W$, with $W \in \mathbb{Z}$ by allowing for a term linear in σ in x_+ .

For \mathcal{H} to be positive, p_- should be taken to be the *negative* root of (2.12). We also see that the worldsheet and target-space energies are related by

$$\mathbb{H} = \mathbb{H}^{\text{t.s.}} - \mathbb{J}, \quad (2.17)$$

where \mathbb{H} is the worldsheet Hamiltonian. When eliminating the longitudinal degrees of freedom we did not solve for x_- , but only for its derivative x'_- . For consistency, we have to impose that x_- is periodic

$$0 = \int_{-\ell/2}^{\ell/2} d\sigma x'_- = - \int_{-\ell/2}^{\ell/2} d\sigma p_j x'^j. \quad (2.18)$$

This last condition, the *level-matching constraint*, is difficult to impose before quantisation. Instead, we will impose it on the Hilbert space of the quantum theory. To this end, notice that the worldsheet momentum, *i.e.* the Noether charge corresponding to σ -translations, is given precisely by

$$\mathbb{P} = - \int_{-\ell/2}^{\ell/2} d\sigma p_j x'^j. \quad (2.19)$$

This should not be confused with the Noether charge \mathbb{P}_+ , which is constant. Therefore, the physical states in the quantum theory will be the ones annihilated by the (quantum) worldsheet momentum.

Perturbative expansion of the action

It is still not straightforward to quantise the Hamiltonian $\mathbb{H} = \int d\sigma \mathcal{H}$, which comes from inverting the non-linear constraint \mathcal{C}_2 and is highly interacting. In order to proceed, we need to systematically expand it. A way of doing so is to suitably redefine fields and coordinates so that the Hamiltonian density can be written as

$$\mathcal{H} = \mathcal{H}_2 + \frac{1}{h} \mathcal{H}_4 + \frac{1}{h^2} \mathcal{H}_6 + \dots, \quad (2.20)$$

where \mathcal{H}_2 is quadratic in the fields, \mathcal{H}_4 quartic, *etc.*, and take a large- h limit. One such procedure is the Berenstein-Maldacena-Nastase (BMN) limit [119], whereby

$$P_+ \rightarrow \infty, \quad h \rightarrow \infty, \quad \frac{P_+}{h} \text{ fixed}. \quad (2.21)$$

The action then admits a perturbative expansion in $1/P_+$.

Alternatively, one can first consider a *decompactification limit*

$$P_+ \rightarrow \infty, \quad h \text{ fixed}, \quad (2.22)$$

which is enough to identify asymptotic states and define an S matrix, since $\ell = P_+$. Then, to perform perturbative calculations we can take a large-tensions expansion in

$1/h$. Building on that, one can go on and construct a perturbative Hamiltonian and S matrix, see *e.g.* ref. [29] for more details. The advantage is this approach is to distinguish the decompactification of the worldsheet from the expansions of the Hamiltonian. In what follows, we will adopt this latter procedure.

Evaluation of the quadratic Hamiltonian

The line element of this geometry reads

$$ds^2 = -G_{tt}^2 dt^2 + G_{\phi\phi}^2 d\phi^2 + f_{ii}^2 dz_i^2 + g_{ii}^2 dy_i^2 + dX_i^2, \quad (2.23)$$

where we denoted the transverse coordinates by z_1, z_2 for AdS_3 , by y_1, y_2 for S^3 and by X_1, \dots, X_4 for the torus. The metric elements are

$$G_{tt} = \frac{4 + |z|^2}{4 - |z|^2}, \quad G_{\phi\phi} = \frac{4 - |y|^2}{4 + |y|^2}, \quad f_{ii} = \frac{4}{4 - |z|^2}, \quad g_{ii} = \frac{4}{4 + |y|^2}. \quad (2.24)$$

From (2.12), we see that it is convenient to perform the rescaling

$$\sigma \rightarrow h\sigma, \quad x^j \rightarrow h^{-1/2}x^j, \quad p_j \rightarrow h^{-1/2}p_j. \quad (2.25)$$

The first replacement suitably rescales the σ -derivatives, and the other two implement a field expansion. In (2.12) we can drop the subleading $(x'_-)^2$ term, and picking the appropriate solution have

$$p_- = -2 \frac{G_{\phi\phi}^{-2} + G_{tt}^{-2}}{G_{\phi\phi}^{-2} - G_{tt}^{-2}} + \frac{\sqrt{4G_{\phi\phi}^{-2}G_{tt}^{-2} - (G_{\phi\phi}^{-2} - G_{tt}^{-2})\mathcal{H}^\perp}}{G_{\phi\phi}^{-2} - G_{tt}^{-2}}, \quad (2.26)$$

with

$$\mathcal{H}^\perp = \frac{1}{h} (\delta^{jk} p_j p_k + \delta_{jk} x'^j x'^k) + O\left(\frac{1}{h^2}\right). \quad (2.27)$$

Expanding to leading order gives

$$\mathcal{H}_2 = \frac{1}{2} (|p_y|^2 + |y'|^2 + |y|^2) + \frac{1}{2} (|p_z|^2 + |z'|^2 + |z|^2) + \frac{1}{2} (|p_X|^2 + |X'|^2). \quad (2.28)$$

This is the (relativistic) Hamiltonian of four massive excitations (the four transverse coordinates in $\text{AdS}_3 \times S^3$) and four massless ones (the torus coordinates). By supersymmetry, we expect the superstring action at quadratic order to be given by the above \mathcal{H}_2 plus $4 + 4$ free fermions of the same masses.

At this order \mathbb{H} is free and can be quantised in a standard way in terms of raising and lowering operators $a_j^\dagger(p)$, $a^j(p)$ satisfying canonical commutation relations

$$[a^k(p), a_j^\dagger(\tilde{p})] = \delta^k_j \delta(p - \tilde{p}), \quad (2.29)$$

yielding³

$$\mathbb{H} = \int d\sigma \mathcal{H}_2 = \int dp \sum_{j=1}^8 \omega_j(p) a^j(p) a_j^\dagger(p), \quad \omega_j = \sqrt{m_j^2 + p^2}, \quad (2.30)$$

with $m_j = 0$ for the four number operators of the torus and $m_j = 1$ for $\text{AdS}_3 \times \text{S}^3$. The worldsheet momentum (2.19) is then

$$\mathbb{P} = \int dp \sum_{j=1}^8 p a^j(p) a_j^\dagger(p), \quad (2.31)$$

so that a multiparticle state $|p_1, \dots, p_n\rangle$ is physical if and only if $p_1 + \dots + p_n = 0$.

2.2 The $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ supercoset

In this section we will write down the coset action for the superstring, following [50]. Even if this construction is inspired by the one originally performed in $\text{AdS}_5 \times \text{S}^5$ [53], there are some remarkable differences due to the presence of the T^4 flat directions. In that case, the coset action coincided with the GS action *before* fixing the κ -gauge. Here, the correspondence holds only when κ -gauge is *completely fixed*. This will lead to some complications that fortunately are irrelevant as long as we focus on the massive modes alone. We will describe $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ as the coset (2.1) which can be essentially constructed out of *two copies* of the superalgebra $\mathfrak{psu}(1, 1|2)$.

The superalgebra $\mathfrak{psu}(1, 1|2)$

The superalgebra $\mathfrak{psu}(1, 1|2)$ consists of an even (“bosonic”) part given by a non-compact $\mathfrak{su}(1, 1)$ or $\mathfrak{sl}(2)$, and a compact $\mathfrak{su}(2)$ algebra. We can think of the former as being some of the isometries from AdS_3 and the latter as coming from S^3 . These are supplemented by eight (“fermionic”) supercharges.

Let us denote the generators of the non-compact bosonic algebra as \mathbb{L}_i , the ones from $\mathfrak{su}(2)$ as \mathbb{J}_i , and the supercharges as $\mathbb{Q}_{a\kappa l}$. In terms of raising and lowering operators the even part of the algebra, in a suitable real form, is given by

$$\begin{aligned} [\mathbb{L}_3, \mathbb{L}_\pm] &= \mp i \mathbb{L}_\pm, & [\mathbb{L}_+, \mathbb{L}_-] &= +2i \mathbb{L}_3, \\ [\mathbb{J}_3, \mathbb{J}_\pm] &= \mp i \mathbb{J}_\pm, & [\mathbb{J}_+, \mathbb{J}_-] &= -2i \mathbb{J}_3, \end{aligned} \quad (2.32)$$

and the supercharges are charged under the bosonic subalgebra

$$\begin{aligned} [\mathbb{L}_3, \mathbb{Q}_{\pm\kappa l}] &= \pm \frac{i}{2} \mathbb{Q}_{\pm\kappa l}, & [\mathbb{L}_\pm, \mathbb{Q}_{\pm\kappa l}] &= + \mathbb{Q}_{\mp\kappa l}, \\ [\mathbb{J}_3, \mathbb{Q}_{a\pm l}] &= \mp \frac{i}{2} \mathbb{Q}_{a\pm l}, & [\mathbb{J}_\pm, \mathbb{Q}_{a\mp l}] &= -i \mathbb{Q}_{a\pm l}, \end{aligned} \quad (2.33)$$

³Owing to the decompactification limit, the integration ranges from $-\infty$ to $+\infty$, and we omit to indicate the limits in the integral.

with $\kappa = \pm, \iota = \pm$ and $a = \pm$. Finally, the supercharges' anticommutators read

$$\begin{aligned} \{\mathbb{Q}_{\pm\pm\pm}, \mathbb{Q}_{\pm\pm\pm}\} &= \pm \mathbb{L}_{\mp}, & \{\mathbb{Q}_{\pm\pm-}, \mathbb{Q}_{\pm\pm-}\} &= \mp \mathbb{L}_{\mp}, \\ \{\mathbb{Q}_{\pm\pm\pm}, \mathbb{Q}_{-\pm\pm}\} &= \mp i \mathbb{J}_{\pm}, & \{\mathbb{Q}_{\pm\pm-}, \mathbb{Q}_{-\pm\pm}\} &= \pm i \mathbb{J}_{\pm}, \\ \{\mathbb{Q}_{\pm\pm\pm}, \mathbb{Q}_{-\mp\mp}\} &= i(+\mathbb{L}_3 \pm \mathbb{J}_3), & \{\mathbb{Q}_{\pm\pm\mp}, \mathbb{Q}_{-\mp\pm}\} &= i(-\mathbb{L}_3 \mp \mathbb{J}_3). \end{aligned} \quad (2.34)$$

In what follows, we will need to identify two copies of this algebra with part of the superisometries of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, and use an explicit representation for it. In order to do so, we will consider a realisation of the $\mathfrak{su}(1,1|2)$ in terms of supermatrices, and find $\mathfrak{psu}(1,1|2)$ as a quotient subalgebra.

Supermatrix realisation

Let us consider \mathbb{Z}_2 -graded vector space $\mathbb{C}^{2|2}$. The set of its linear endomorphisms form the superalgebra $\mathfrak{gl}(2|2)$, that can be represented in terms of 4×4 supermatrices

$$\mathcal{M} = \begin{pmatrix} A & \Theta \\ \Xi & B \end{pmatrix}, \quad (2.35)$$

where the 2×2 blocks A, B are even and Θ, Ξ are odd⁴. The subalgebra $\mathfrak{u}(1,1|2)$ can be singled out by imposing a suitable hermiticity condition

$$\mathcal{M}^\dagger + H^{-1} \mathcal{M} H = 0, \quad H = H^{-1} = \text{diag}(1, -1, 1, 1), \quad (2.36)$$

where H implements the non-euclidean signature. There are 8 odd and 8 even independent solutions to such condition. Among the latter, there are two central elements

$$\mathbb{I} = \text{diag}(1, 1, 1, 1), \quad \mathbb{I}^s = \text{diag}(1, 1, -1, -1). \quad (2.37)$$

The supertrace

$$\text{str } \mathcal{M} = \text{tr } A - \text{tr } B, \quad (2.38)$$

is an invariant of $\mathfrak{gl}(2|2)$ and $\mathfrak{u}(1,1|2)$ which can be used to impose the condition $\text{str } \mathcal{M} = 0$, that mods out \mathbb{I}^s and defines the matrix algebra $\mathfrak{su}(1,1|2)$. However, we cannot get rid of \mathbb{I} by consistently imposing $\text{tr } \mathcal{M} = 0$ because for a generic odd (and therefore traceless) element of $\mathfrak{u}(1,1|2)$ we have

$$\text{tr}(\{\mathcal{M}_{\text{odd}}, \mathcal{M}_{\text{odd}}\}) = -2 \text{tr}(\mathcal{M}_{\text{odd}} H^{-1} \mathcal{M}_{\text{odd}}^\dagger H) < 0. \quad (2.39)$$

Therefore $\mathfrak{psu}(1,1|2)$ does not admits a matrix realisation. With a small abuse of language we will refer to the quotient $\mathfrak{su}(1,1|2)/\mathbb{I}$ as the ‘‘matrix realisation’’ of $\mathfrak{psu}(1,1|2)$ and, when writing anticommutators, understand equalities modulo a multiple of the identity. In appendix A.1 we give the explicit form of the generators in terms of 4×4 supermatrices, as well as some additional properties of the superalgebra.

⁴Note that the odd elements of the supermatrix are nonetheless commuting (non-Grassmann) scalars.

The full algebra and \mathbb{Z}_4 -automorphism

The algebra generating the supercoset (2.1) is given by two copies of $\mathfrak{psu}(1, 1|2)$, which we will denote by “L” or “left” and “R” or “right” and four copies of $\mathfrak{u}(1)$. As it is always the case when considering a superalgebra that is direct sum of two subalgebras, $\mathfrak{psu}(1, 1|2)_L \oplus \mathfrak{psu}(1, 1|2)_R$ enjoys a \mathbb{Z}_4 -automorphism Ω , on top of the natural \mathbb{Z}_2 -grading due to the superalgebra structure. This can be defined, for instance, by a permutation of the two copies together with multiplication by the fermion sign of one of them.

There are several inequivalent ways to define a realisation of $\mathfrak{psu}(1, 1|2)_L \oplus \mathfrak{psu}(1, 1|2)_R$ in terms of 8×8 supermatrices \mathcal{M} . A possibility would be to consider two identical realisations of $\mathfrak{psu}(1, 1|2)$ and take their direct sum. However, this is not the most suitable choice for the coset construction that we will carry out in the next section.⁵ The elements of $\mathfrak{psu}(1, 1|2)_L \oplus \mathfrak{psu}(1, 1|2)_R$ can be split in two,

$$\mathcal{M}_L = \mathcal{M} \oplus \mathbb{O}, \quad \mathcal{M}_R = \mathbb{O} \oplus \widetilde{\mathcal{M}}, \quad (2.40)$$

where \mathbb{O} is the zero matrix, and \mathcal{M} and $\widetilde{\mathcal{M}}$ are in two matrix representations whose bosonic subalgebras have opposite notions of what the highest weight vectors are. In appendix A.1 we give the explicit form of these matrices and comment more on this choice. There are also several inequivalent ways of defining the \mathbb{Z}_4 automorphism Ω . We will rely on an exchange of the two $\mathfrak{psu}(1, 1|2)$ copies supplemented by a “fermionic” operation. In particular, Ω will take the form

$$\Omega(\mathcal{M}) = \mathcal{K}^{-1} \mathcal{M} \mathcal{K}, \quad \Omega^4(\mathcal{M}) = \mathcal{M}, \quad (2.41)$$

where \mathcal{K} permutes left and right and takes into account the fermionic signs, so that in terms of the (Hermitian) Pauli matrices we have

$$\mathcal{K} = \mathcal{F}_L \mathcal{P}, \quad \mathcal{F}_L = \mathbb{I}_2 \otimes (\sigma_3 \oplus \mathbb{I}_2), \quad \mathcal{P} = \sigma_1 \otimes (\mathbb{I}_2 \otimes \sigma_1). \quad (2.42)$$

In this way, Ω maps the left and right bosonic subalgebras into each other and accounts for their different matrix representations, see appendix A.1.

The advantage of such a construction is that it gives a natural way to decompose the symmetry algebra into the direct sum of four eigenspaces relative to the eigenvalues i^k with $k = 0, \dots, 3$. Using the matrix representation (2.35–2.40) it is easy to see that the eigenspaces relative to ± 1 consist of even elements. In particular, the eigenspace given by matrices such that $\Omega(\mathcal{M}) = \mathcal{M}$ consists of $\mathfrak{su}(1, 1) \oplus \mathfrak{su}(2)$. The corresponding group is precisely the quotient part of the coset. It only remains to extend the automorphism to $\mathfrak{u}(1)$ ⁴. A simple way of doing so is declaring that all of the $\mathfrak{u}(1)$ generators have eigenvalue -1 under Ω . A more “symmetric” choice is to introduce a \mathbb{Z}_2 -grading on $\mathfrak{u}(1)$ ⁸,

⁵In fact, it is easy to check that carrying out the construction with the more naïve representation would lead to an ill-defined Poisson structure in the resulting coset action.

which amounts to realising $U(1)^4$ as a coset $(U(1)^2/U(1))^4$. Since the latter is trivial, the two descriptions are equivalent.

Therefore, we have a vector-space decomposition of

$$\mathcal{A} = \mathfrak{psu}(1, 1|2) \oplus \mathfrak{psu}(1, 1|2) \oplus \mathfrak{u}(1)^4, \quad (2.43)$$

as

$$\mathcal{A} = \bigoplus_{k=0}^3 \mathcal{A}^{(k)}, \quad \Omega(\mathcal{A}^{(k)}) = i^k \mathcal{A}^{(k)}. \quad (2.44)$$

Using the fact that Ω is realised linearly and that

$$\Omega(\mathcal{M}_1 \mathcal{M}_2) = \Omega(\mathcal{M}_1) \Omega(\mathcal{M}_2), \quad (2.45)$$

we have that the decomposition turns \mathcal{A} into a \mathbb{Z}_4 -graded Lie superalgebra, *i.e.*

$$[\mathcal{A}^{(j)}, \mathcal{A}^{(k)}] \subset \mathcal{A}^{(j+k \bmod 4)}. \quad (2.46)$$

This structure has two important consequences:

- It provides a simple way to realise the supergroup coset starting from the superalgebra elements and to write down the string action and its symmetries, as we will do in the rest of this chapter.
- It ensures that the string equations of motions can be written as the flatness condition for a suitable (Lax) connection, a feature that guarantees classical integrability, see section 3.1.

Parametrisation of group elements

It is convenient to parametrise a generic element $g \in PSU(1, 1|2)^2 \times U(1)^4$ in terms of the “exponential” of the algebra elements. Several choices are possible. Here we will set

$$g(t, \phi, \chi, x, X) = \Lambda(t, \phi) g(\Psi_m) g(\Psi_l) g(x) g(X), \quad (2.47)$$

where

- t, ϕ are the time coordinate in AdS_3 and an angle in S^3 , respectively. We will later use them as light-cone coordinates.
- $\Psi_{m,l}$ are all the fermions, which we will split into massive fermions $\theta_{L,R}^i$ contained in Ψ_m , and massless fermions $\eta_{L,R}^i$ contained in Ψ_l .
- $x = (z_i, y_i)$ are the remaining four bosonic coordinates on $\text{AdS}_3 \times \text{S}^3$.
- $X = (X_1, \dots, X_4)$ are the four bosons from T^4 , so that $g(X)$ commutes with all the other parameters.

In particular, let us set

$$\Lambda(t, \phi) = e^{\frac{i}{2}(\mathbb{L}_3^L - \mathbb{L}_3^R) + \frac{\phi}{2}(\mathbb{J}_3^L - \mathbb{J}_3^R)} = e^{i x_+ \Sigma_+ - \frac{i}{2} x_- \Sigma_-}, \quad (2.48)$$

where we introduced the light-cone matrices in $\mathfrak{psu}(1, 1|2)^2$

$$i \Sigma_{\pm} = (\mathbb{L}_3^L - \mathbb{L}_3^R) \pm (\mathbb{J}_3^L - \mathbb{J}_3^R). \quad (2.49)$$

The remaining four fields $\{x_k\} = (z_1, z_2, y_1, y_2)$ appear through $g(x)$ which is written in terms of the remaining transverse angular momenta. For their form as well as for the forms of the fermion parametrisation we refer the reader to appendix A.2.

Lagrangian and Noether current

Let $g \in PSU(1, 1|2)^2 \times U(1)^4$. The one-form

$$A = -g^{-1} dg, \quad (2.50)$$

takes values in \mathcal{A} , and therefore can be decomposed according to (2.44) as

$$A = A^{(1)} + A^{(2)} + A^{(3)} + A^{(4)}. \quad (2.51)$$

Inspired by the case of $\text{AdS}_5 \times S^5$, let us define the Lagrangian

$$\mathcal{L} = -\frac{\hbar}{2} \left(\gamma^{\alpha\beta} \text{str}(A_{\alpha}^{(2)} A_{\beta}^{(2)}) + \varepsilon^{\alpha\beta} \text{str}(A_{\alpha}^{(1)} A_{\beta}^{(3)}) \right), \quad (2.52)$$

where definiteness we fixed the coefficient of the Wess-Zumino term $\text{str} A^{(1)} \wedge A^{(3)}$. This Lagrangian and the resulting action have several good properties. Firstly, as it is easy to check they are manifestly real. Furthermore, A and therefore \mathcal{L} is invariant under left multiplication of the group element g by any constant element g_0 —the theory is manifestly invariant under *global* (super)isometries.

One may worry that the action depends on g , rather than on the corresponding coset representative. However, it is easy to see that under a transformation of the $SU(1, 1) \times SU(2)$ from $\mathcal{A}^{(0)}$ all the current components appearing in the action $A^{(k)}$, $k = 1, \dots, 3$ transform covariantly, so that \mathcal{L} is left unchanged. Moreover, if we take g to be parametrised by (2.47) and set the coefficients of the odd algebra elements to zero, we have that the Lagrangian \mathcal{L} reduces to the one of the bosonic NLSM on $\text{AdS}_3 \times S^3 \times T^4$ given by eq. (2.2).

An important feature of (2.52) is that the resulting equations of motion can be written as a flatness condition for a Lax connection. This fact, familiar from $\text{AdS}_5 \times S^5$ [19] was pointed out in ref. [50] for the $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ background of which $\text{AdS}_3 \times S^3 \times T^4$ can be seen as a limit, at least in absence of winding. In ref. [50] it was also argued that the coset action coincides with the GS superstring action in a suitable κ -gauge, namely

one where all the massless fermions are eliminated by the gauge fixing. This was checked up to quadratic order in the fermions.

Finally, owing to the global \mathcal{A} -symmetry, there exists a conserved Noether current, which takes a simple form in the coset formulation [50]

$$\mathcal{J}^\alpha = g \left(\gamma^{\alpha\beta} A_\beta^{(2)} - \frac{1}{2} \varepsilon^{\alpha\beta} (A_\beta^{(1)} - A_\beta^{(3)}) \right) g^{-1}. \quad (2.53)$$

2.3 Massive modes in light-cone gauge

Following what we did in section 2.1, let us rewrite the Lagrangian (2.52) in the first-order formalism,

$$\mathcal{L} = - \left(\text{str}(\varpi A_0^{(2)}) + \frac{h}{2} \varepsilon^{\alpha\beta} \text{str}(A_\alpha^{(1)} A_\beta^{(3)}) + \frac{\gamma^{01}}{\gamma^{00}} \mathcal{C}_1 - \frac{1}{2h\gamma^{00}} \mathcal{C}_2 \right), \quad (2.54)$$

where we introduced the auxiliary field ϖ that without loss of generality we take to be equal to its component $\varpi^{(2)}$. The constraints are $\mathcal{C}_1 = \mathcal{C}_2 = 0$ where now

$$\mathcal{C}_1 = \text{str}(\varpi A_1^{(2)}), \quad \mathcal{C}_2 = \text{str}(\varpi^2 + h(A_1^{(2)})^2). \quad (2.55)$$

To preserve as much supersymmetry as possible, we want to fix light-cone gauge in terms of the coordinates x_\pm constructed in section 2.2. Such a gauge fixing is incompatible with the κ -symmetry fixing that was necessary to assume to have a coset description [50], and should we proceed in this way we would find that the massless fermions lack a good kinetic term (*i.e.* quadratic in a field expansion). In what follows, we will restrict to considering *only massive excitations*, *i.e.* truncate the coset to $\text{AdS}_3 \times \text{S}^3$. This will be enough to elucidate at least some of the symmetries of the theory, which is what we will later need to find the worldsheet S matrix for massive particles. Let us therefore set, from now on,

$$\Psi_l = 0, \quad X_1 = \dots = X_4 = 0. \quad (2.56)$$

Evaluation of the action

It is useful to split the current into an even and odd part under the \mathbb{Z}_4 decomposition, A_{even} and A_{odd} respectively. We also single out the part A_{even}^\perp of A_{even} which depends only on the transverse bosonic coordinates. In terms of these quantities (whose explicit expression we give in appendix A.2) and up to the imposing the Virasoro constraints, the Lagrangian can be rewritten as

$$\mathcal{L} = p_+ \dot{x}_- + \mathbf{p}_- \dot{x}_+ - \text{str}(\varpi A_{\text{even}}^\perp) - i \frac{h}{4} \varepsilon^{\alpha\beta} \text{str}(A_\alpha^{\text{odd}} \Omega(A_\beta^{\text{odd}})), \quad (2.57)$$

where we also made use of the \mathbb{Z}_4 automorphism. Note that p_+ is the momentum conjugate to x_- , while \mathbf{p}_- differs from the conjugate momentum to x_+ due to the contribution of the Wess-Zumino term. To see this, let us define the decomposition

$$\varpi = \frac{i}{2}\varpi_+\Sigma_+ + \frac{i}{4}\varpi_-\Sigma_- - \frac{1}{2}\varpi_k\Sigma_k, \quad (2.58)$$

valid up to trace contributions, where Σ_k corresponds to the bosonic generators of the transverse directions, see eq. (A.14). Then, using the explicit expressions from appendix A.2, we get indeed

$$p_+ = \varpi_+G_+ - \frac{1}{2}\varpi_-G_-, \quad \mathbf{p}_- = \frac{i}{2}\text{str}(\varpi\Sigma_+g_x(1 + 2\Psi_m^2)g_x). \quad (2.59)$$

Note that we introduced the short-hand notation $g_x = g(x)$ for the transverse bosonic coordinates and the metric in the light-cone directions, $G_{\pm} = \frac{1}{2}(G_{tt} \pm G_{\phi\phi})$.

Gauge fixing

Let us now fix light-cone gauge (with zero winding)

$$x_+ = \tau, \quad p_+ = 1. \quad (2.60)$$

From the latter equation we can immediately find the value of

$$\varpi_+ = \frac{1}{G_+}\left(1 + \frac{1}{2}\varpi_-G_-\right). \quad (2.61)$$

Substituting ϖ_+ in the $\mathcal{C}_1 = 0$ constraint we find as expected

$$-x'_- = p_k x'_k + \text{fermions}, \quad (2.62)$$

where the complete expression of the fermion contributions are given in eq. (A.29). Recall that it is this expression that appears in the level matching constrain, which as in the bosonic case amounts to vanishing of the worldsheet momentum

$$0 = - \int_{-\ell/2}^{\ell/2} d\sigma x'_- = \mathbb{P}. \quad (2.63)$$

The last longitudinal component of the auxiliary field ϖ_- can be found from the quadratic constraint $\mathcal{C}_2 = 0$, with

$$\mathcal{C}_2 = \varpi_+\varpi_- + \varpi_k^2 + h \text{str}\left((A_{\sigma}^{(2)})^2\right). \quad (2.64)$$

The expansion of and solution to this constraint is given in appendix A.2, where we also give the form of the auxiliary fields ϖ_k in terms of the conjugate momenta p_k as well as some other useful formulae. Using all this, in the next section we will write down explicit expressions to the leading order in a field expansion.

Noether charge

We conclude this section by writing down the conserved charges corresponding to the Noether current (2.53). By making use of the equations of motion for the auxiliary field ϖ

$$\varpi = h \gamma^{0\beta} A_\beta^{(2)}, \quad (2.65)$$

we can write down the charges in a form that is independent of the worldsheet metric,

$$\mathcal{Q} = \int_{-\ell/2}^{\ell/2} d\sigma g(x_\pm, x, \Psi_m) \left(\varpi - \frac{1}{2} h (A_1^{(1)} - A_1^{(3)}) \right) g^{-1}(x_\pm, x, \Psi_m), \quad (2.66)$$

which by using the \mathbb{Z}_4 -automorphism Ω can be recast in the form

$$\mathcal{Q} = \int_{-\ell/2}^{\ell/2} d\sigma \Lambda g_{\Psi_m} g_x \left(\varpi + \frac{i}{2} h g_x \Omega(\partial_0 \Psi_m) g_x^{-1} \right) g_x^{-1} g_{\Psi_m}^{-1} \Lambda^{-1}, \quad (2.67)$$

where we also expressed $g(x_\pm, x, \Psi_m)$ by means of (2.47). The Noether charge $\mathcal{Q} \in \mathfrak{psu}(1, 1|2)^2$ is written as a matrix, and its independent components may be projected by defining

$$\mathbb{Q}_{\mathcal{M}} = \text{str}(\mathcal{M} \mathcal{Q}), \quad \mathcal{M} \in \mathfrak{psu}(1, 1|2)^2, \quad (2.68)$$

which relates the superalgebra structure of $\mathfrak{psu}(1, 1|2)^2$ to the one induced in the phase space by the Poisson brackets⁶. By construction $\mathbb{Q}_{\mathcal{M}}$ will be conserved,

$$0 = \frac{d\mathbb{Q}_{\mathcal{M}}}{d\tau} = \frac{\partial \mathbb{Q}_{\mathcal{M}}}{\partial \tau} + \{\mathbb{H}, \mathbb{Q}_{\mathcal{M}}\}, \quad (2.69)$$

where in the last equality we have expressed the conservation law in terms of the Poisson structure. This highlights the fact that *some charges will not commute with the Hamiltonian*, namely the ones that depend explicitly on $\tau = x_+$. Only the remaining charges will constitute the manifest symmetry algebra of the theory, with \mathbb{H} as central element. This is similar to what we discussed in the introduction for strings in flat space, where the $\mathfrak{so}(1, 9)$ symmetry is broken down to $\mathfrak{so}(8)$.

The x_+ dependence enters (2.67) in a simple way, *i.e.* only through $\Lambda = \Lambda(x_+, x_-)$. This makes it easy to identify the elements of \mathcal{M} that give rise to charges commuting with \mathbb{H} . These are depicted in figure 2.1, and consist of the bosonic charges lying on the diagonal of \mathcal{M} and eight supercharges. Similarly, it is also easy to see that all of these supercharges carry a dependence on the unphysical (and highly non-local) field x_- , a fact that will be important later.

⁶This pairing can be understood in terms of the moment map [57, 120].

$$\mathcal{M} = \begin{pmatrix} \text{K} & \text{D} & \text{D} & \text{K} \\ \text{D} & \text{K} & \text{K} & \text{D} \\ \text{D} & \text{K} & \text{K} & \text{D} \\ \text{K} & \text{D} & \text{D} & \text{K} \end{pmatrix} \oplus \begin{pmatrix} \text{K} & \text{D} & \text{D} & \text{K} \\ \text{D} & \text{K} & \text{K} & \text{D} \\ \text{D} & \text{K} & \text{K} & \text{D} \\ \text{K} & \text{D} & \text{D} & \text{K} \end{pmatrix}$$

Figure 2.1: The elements of the $\mathfrak{psu}(1,1|2)^2$ matrix \mathcal{M} of (2.68), distinguished by the dependence on x_{\pm} in the resulting charge $\mathbb{Q}_{\mathcal{M}}$. Elements on a white background yield an x_+ -dependent charge (that does not commute with \mathbb{H}), while the one highlighted in yellow **K**, **D** yield conserved charges. We further distinguish between kinematical (K), *i.e.* x_- -independent charges, and dynamical ones (D).

Perturbative evaluation at leading order

We now want to find explicit expressions for the action and its symmetries perturbatively in a field expansion. To this end, as discussed in section 2.1 we will take the decompactification limit and perform a field expansion. We are interested in the quadratic Hamiltonian, that comes at leading order in $1/h$, and should be a suitable supersymmetric completion of (2.28), together with its symmetry algebra.

Quadratic expressions in the fields

Using the expansion of the auxiliary fields provided in appendix A.2 we find that the Lagrangian can be written, at leading order, as

$$\mathcal{L}_2 = p_k \dot{x}_k - \frac{i}{2} \text{str}(\Sigma_+ \Psi_m \dot{\Psi}_m) - \mathcal{H}_2, \quad (2.70)$$

where the quadratic Hamiltonian is

$$\mathcal{H}_2 = \frac{1}{2} |p_k|^2 + \frac{1}{2} |x'_k|^2 + \frac{1}{2} |x_k|^2 - \frac{1}{2} \text{str}(\Sigma_+ \Psi_m \Omega(\Psi'_m)) + \frac{1}{2} \text{str}(\Psi_m^2). \quad (2.71)$$

As expected, this is a supersymmetric relativistic extension of what computed in the purely bosonic NLSM. In order to write down more explicitly these results, let us use the fermion parametrisation (A.15) so that the Lagrangian becomes

$$\mathcal{L}_2 = p_k \dot{x}_k + i \bar{\theta}_{j_L} \dot{\theta}^{j_L} + i \bar{\theta}_{j_R} \dot{\theta}^{j_R} - \mathcal{H}_2, \quad (2.72)$$

where the fermion fields are conjugate to each other, $\bar{\theta} = \theta^\dagger$, and indices are raised and lowered with δ^{jk}, δ_{jk} . The quadratic Hamiltonian is

$$\mathcal{H}_2 = \frac{1}{2} |p_k|^2 + \frac{1}{2} |x'_k|^2 + \frac{1}{2} |x_k|^2 + \theta_L^j \theta'_{j_R} - \bar{\theta}'_{j_R} \bar{\theta}_{j_L} + \bar{\theta}_{j_L} \theta_L^j + \bar{\theta}_{j_R} \theta_R^j. \quad (2.73)$$

In particular, this implies that the non-vanishing Poisson brackets are

$$\{x_j(\sigma), p_k(\tilde{\sigma})\} = \delta_{jk} \delta(\sigma - \tilde{\sigma}), \quad (2.74)$$

and

$$\{\theta_L^j(\sigma), \bar{\theta}_{k_L}(\tilde{\sigma})\} = \delta_k^j \delta(\sigma - \tilde{\sigma}), \quad \{\theta_R^j(\sigma), \bar{\theta}_{k_R}(\tilde{\sigma})\} = \delta_k^j \delta(\sigma - \tilde{\sigma}). \quad (2.75)$$

Let us now work out the charges that commute with the Hamiltonian. We start from the bosonic charges. From figure 2.1 we see that they can be found from contracting \mathcal{Q} with a diagonal matrix \mathcal{M} . Taking into account that $\mathcal{M} \in \mathfrak{psu}(1,1|2)^2$, we are left with four independent choices of \mathcal{M} , yielding as many central charges. Two of these are well familiar:

$$\frac{1}{2i} \text{str}(\Sigma_+ \mathcal{Q}) = \mathbb{H}, \quad \frac{1}{2i} \text{str}(\Sigma_- \mathcal{Q}) = -2\mathbb{P}_+ = -2P_+. \quad (2.76)$$

The remaining two come from

$$\mathcal{M} = \mathbb{S}_L^3 + \mathbb{S}_R^3 + \mathbb{J}_L^3 + \mathbb{J}_R^3, \quad \mathcal{N} = \mathbb{S}_L^3 + \mathbb{S}_R^3 - \mathbb{J}_L^3 - \mathbb{J}_R^3, \quad (2.77)$$

and are given in appendix A.3. We define the supercharges that commute with the Hamiltonian by contracting \mathcal{Q} with odd supermatrices with a single non-vanishing entry (akin to raising and lowering operators). As expected we find eight supercharges, that we label

$$\mathbb{Q}^{1L,R}, \quad \mathbb{Q}^{2L,R}, \quad \bar{\mathbb{Q}}^{1L,R}, \quad \bar{\mathbb{Q}}^{2L,R}, \quad (2.78)$$

and which are Hermitian conjugate in pairs, $(\mathbb{Q}^{jL,R})^\dagger = \bar{\mathbb{Q}}^{jL,R}$. The general form of the charges in terms of fields is

$$\begin{aligned} \mathbb{Q} &\approx \int d\sigma e^{+\frac{i}{2}x-} (p\theta + x\bar{\theta}' + x\theta), \\ \bar{\mathbb{Q}} &\approx \int d\sigma e^{-\frac{i}{2}x-} (p\theta + x\bar{\theta}' + x\theta), \end{aligned} \quad (2.79)$$

where we considered only the leading order contribution inside the brackets but kept track of the $e^{\pm\frac{i}{2}x-}$ factor coming from $\Lambda(x_\pm)$. The precise form of all charges is given in appendix A.3. It is however more convenient to take a Fourier transform and introduce a basis of raising and lowering operators, thereby making the action of the σ -derivatives more apparent, and diagonalizing some of the charges. In particular, we will take all of the central elements to be proportional to the number operator.

Momentum-space representation

Let us introduce the bosonic creation and annihilation operators $a_{L,R}^{\dagger\pm\pm}$ and $a_{L,R}^{\pm\pm}$, as defined in appendix A.3. An annihilation operator takes the schematic form

$$a_{\pm\pm}(q) \approx \frac{1}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_q}} (\omega_q x(\sigma) + ip(\sigma)) e^{-iq\sigma} \quad (2.80)$$

where

$$\omega_p = \sqrt{m^2 + p^2}. \quad (2.81)$$

The respective creation operator is the complex conjugate of (2.80). The definitions in appendix A.3 are given in such a way that the operators have canonical commutation relations

$$[a_L^{\kappa\iota}(p), a_L^\dagger{}_{\bar{\kappa}\bar{\iota}}(\tilde{p})] = \delta_{\bar{\kappa}}^\kappa \delta_{\bar{\iota}}^\iota \delta(p - \tilde{p}), \quad [a_R^{\kappa\iota}(p), a_R^\dagger{}_{\bar{\kappa}\bar{\iota}}(\tilde{p})] = \delta_{\bar{\kappa}}^\kappa \delta_{\bar{\iota}}^\iota \delta(p - \tilde{p}). \quad (2.82)$$

Similarly, we introduce the fermionic operators $a_{L,R}^\dagger{}_{\pm\mp}$ and $a_{L,R}^{\pm\mp}$, also defined in appendix A.3, that have the general form *e.g.*

$$a_{\pm\mp}(q) \approx \frac{1}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_q}} (f_q \bar{\theta}(\sigma) + g_q \theta(\sigma)) e^{-iq\sigma}, \quad (2.83)$$

where we used the fermion wave-function parameters

$$f_p = \sqrt{\frac{\omega_p + m}{2}}, \quad g_p f_p = \frac{p}{2}, \quad f_p^2 - g_p^2 = m, \quad f_p^2 + g_p^2 = \omega_p. \quad (2.84)$$

The fermionic operators are defined to satisfy

$$\{a_L^{\kappa\iota}(p), a_L^\dagger{}_{\bar{\kappa}\bar{\iota}}(\tilde{p})\} = \delta_{\bar{\kappa}}^\kappa \delta_{\bar{\iota}}^\iota \delta(p - \tilde{p}), \quad \{a_R^{\kappa\iota}(p), a_R^\dagger{}_{\bar{\kappa}\bar{\iota}}(\tilde{p})\} = \delta_{\bar{\kappa}}^\kappa \delta_{\bar{\iota}}^\iota \delta(p - \tilde{p}). \quad (2.85)$$

Since the Hamiltonian (2.73) contains only particles of unit mass, from now on we set $m = 1$. At leading order, the Hamiltonian and the other central charges take the form

$$\begin{aligned} \mathbb{H} &= \int dp \sum_{\kappa,\iota=\pm} \omega_p (a_L^\dagger{}_{\kappa\iota} a_L^{\kappa\iota} + a_R^\dagger{}_{\kappa\iota} a_R^{\kappa\iota}), \\ \mathbb{M} &= \int dp \sum_{\kappa,\iota=\pm} (a_L^\dagger{}_{\kappa\iota} a_L^{\kappa\iota} - a_R^\dagger{}_{\kappa\iota} a_R^{\kappa\iota}), \\ \mathbb{N} &= \int dp \sum_{\kappa=\pm} \kappa (a_L^\dagger{}_{\kappa\kappa} a_L^{\kappa\kappa} - a_R^\dagger{}_{\kappa\kappa} a_R^{\kappa\kappa}). \end{aligned} \quad (2.86)$$

In particular, we see that L-excitations have charge +1 under the angular momentum \mathbb{M} , while R-excitations have charge -1. We will therefore refer to these excitations as “left” and “right”. However, *these do not correspond to left- and right-moving excitations on the worldsheet*. Since the angular momentum \mathbb{M} has a component on AdS_3 , these excitations can instead be thought of as left- and right-movers in the dual CFT_2 .

The supercharges are given by

$$\begin{aligned} \mathbb{Q}^{1L} &= \int dp \sum_{\iota=\pm} (f_p a_L^\dagger{}_{-l} a_L^{+\iota} - g_p a_R^\dagger{}_{+l} a_R^{-\iota}), \\ \mathbb{Q}^{1R} &= \int dp \sum_{\iota=\pm} (f_p a_R^\dagger{}_{-l} a_R^{+\iota} - g_p a_L^\dagger{}_{+l} a_L^{-\iota}), \end{aligned} \quad (2.87)$$

and

$$\begin{aligned}\mathbb{Q}^{2L} &= \int dp \sum_{\kappa=\pm} \kappa \left(f_p a_{L\kappa}^\dagger a_L^{\kappa+} - g_p a_{R\kappa}^\dagger a_R^{\kappa-} \right), \\ \mathbb{Q}^{2R} &= \int dp \sum_{\kappa=\pm} \kappa \left(f_p a_{R\kappa}^\dagger a_R^{\kappa+} - g_p a_{L\kappa}^\dagger a_L^{\kappa-} \right),\end{aligned}\tag{2.88}$$

together with their Hermitian conjugates $\overline{\mathbb{Q}}^{jL,R} = (\mathbb{Q}^{jL,R})^\dagger$. Notice how the form of the supercharges is completely symmetric under exchange of the labels $L \leftrightarrow R$, a fact that is true also when we write them in terms of the field, see eq. (A.46).

Finally, let us recall that the worldsheet momentum is given by

$$\mathbb{P} = \int dp \sum_{\kappa,\ell=\pm} p \left(a_{L\kappa\ell}^\dagger a_L^{\kappa\ell} + a_{R\kappa\ell}^\dagger a_R^{\kappa\ell} \right).\tag{2.89}$$

2.4 Symmetry algebra and its representation

We can now use the perturbative information we just obtained to study the symmetry algebra and its representations.

Commutation relations

From the explicit results of the previous section it is easy to read off the anticommutation relations. The non-vanishing ones are

$$\begin{aligned}\{\mathbb{Q}^{1L}, \overline{\mathbb{Q}}^{1L}\} &= \{\mathbb{Q}^{2L}, \overline{\mathbb{Q}}^{2L}\} = \frac{1}{2}\mathbb{H} + \frac{1}{2}\mathbb{M}, \\ \{\mathbb{Q}^{1R}, \overline{\mathbb{Q}}^{1R}\} &= \{\mathbb{Q}^{2R}, \overline{\mathbb{Q}}^{2R}\} = \frac{1}{2}\mathbb{H} - \frac{1}{2}\mathbb{M},\end{aligned}\tag{2.90}$$

and

$$\begin{aligned}\{\mathbb{Q}^{1L}, \mathbb{Q}^{1R}\} &= \{\mathbb{Q}^{2L}, \mathbb{Q}^{2R}\} = -\frac{1}{2}\mathbb{P}, \\ \{\overline{\mathbb{Q}}^{1L}, \overline{\mathbb{Q}}^{1R}\} &= \{\overline{\mathbb{Q}}^{2L}, \overline{\mathbb{Q}}^{2R}\} = -\frac{1}{2}\mathbb{P}.\end{aligned}\tag{2.91}$$

From (2.90) we recognise two copies of $\mathfrak{su}(1|1)^2$, and from (2.91) we see that each of them gets centrally extended by two $\mathfrak{u}(1)$ charges, which at leading order coincide and are proportional to the worldsheet momentum⁷. We conclude that the symmetry algebra is

$$\left[(\mathfrak{su}(1|1)_L \oplus \mathfrak{su}(1|1)_R) \oplus (\mathfrak{u}(1) \oplus \mathfrak{u}(1)) \right]^2 / \mathfrak{u}(1)^4,\tag{2.92}$$

where *e.g.* $\mathbb{Q}^{1L}, \overline{\mathbb{Q}}^{1L}$ and their central charge form one of the four $\mathfrak{su}(1|1)$, and $\dots \oplus \mathfrak{u}(1)^2$ denotes the central extension by two copies of \mathbb{P} . The quotient ensures that the central

⁷From this leading-order analysis we cannot establish whether the two right-hand-sides of (2.91) also coincide at higher orders, but only that such charges should be Hermitian conjugate to each other. We will later see they do not coincide, by more carefully investigating the structure of the supercharges.

charges of commutators differing by $1 \leftrightarrow 2$ coincide. Since this algebra can be found from centrally extending $\mathfrak{psu}(1|1)^4$, we indicate it as $\mathfrak{psu}(1|1)_{c.e.}^4$.

It is quite remarkable that the algebra has been extended by two elements proportional to the worldsheet momentum \mathbb{P} , which was not part of the original superisometries. Such an extended algebra should not be a symmetry of the physical states, which is guaranteed by the fact that the level matching constrain (2.63) imposes

$$\mathbb{P}|\text{physical state}\rangle = 0. \quad (2.93)$$

Therefore the algebra (2.92) is valid *off-shell*, whereas *the on-shell algebra* is just

$$(\mathfrak{su}(1|1)_L \oplus \mathfrak{su}(1|1)_R)^2 / \mathfrak{u}(1)^2. \quad (2.94)$$

Representations on one-excitation states

Let us consider excitations of a defined momentum p , which can be created by acting on a vacuum as

$$|\Phi_{\pm\pm}^{L,R}(p)\rangle = a_{\pm\pm L,R}^\dagger(p)|0\rangle, \quad |\Phi_{\pm\mp}^{L,R}(p)\rangle = a_{\pm\mp L,R}^\dagger(p)|0\rangle, \quad (2.95)$$

where the former and latter expressions correspond to bosonic and fermionic excitations, respectively. We should distinguish between representation of the full, off-shell algebra and representations of the on-shell one. The former is realised for general p , while the latter when we impose $p = 0$. Let us start from this simpler case.

On-shell representations

On-shell states are annihilated by the total momentum operator \mathbb{P} , so that one-particle on-shell states are just given by particles with momentum $p = 0$. As a consequence, the on-shell action of the supercharges (2.87–2.88) and their conjugates is easily found, by using that when $p = 0$, we have $f_{p=0} = \omega_{p=0} = 1$ and $g_{p=0} = 0$. We see that L- and R-excitations form two distinct representations that are charged under \mathbb{Q}^L 's and \mathbb{Q}^R 's only, respectively. In figure 2.2 we draw the action of the supercharges. We see that the left excitations transform into a bifundamental of a $\mathfrak{psu}(1|1)_L^2$ consisting of the four L-supercharges. The right excitations are in a bifundamental representation of $\mathfrak{psu}(1|1)_R^2$.

In figure 2.2 we have depicted the left and right representation slightly differently. This is because we cannot take the same raising operators to be the same in the L- and R-algebra if we want this representation to smoothly extend to the off-shell one. If we took the raising operators to be, *e.g.*

$$\overline{\mathbb{Q}}^{1L}, \quad \overline{\mathbb{Q}}^{2L}, \quad \overline{\mathbb{Q}}^{1R}, \quad \overline{\mathbb{Q}}^{2R}, \quad (2.96)$$




Figure 2.2: The action of the supercharges on a zero-momentum (on-shell) worldsheet excitation (2.95). In this case an L-excitation is charged only under Q^{jL} and \bar{Q}^{jL} , and similarly for R-excitations. As explained in the text, the raising operators are \bar{Q}^{jL} and Q^{jR} .

it would be impossible for the anticommutator $\{\bar{Q}^{jL}, \bar{Q}^{kL}\}$ to be non-vanishing on a highest weight state, which should happen if we deform p to be different from zero. Therefore, we will take the raising operators to be

$$\bar{Q}^{1L}, \quad \bar{Q}^{2L}, \quad Q^{1R}, \quad Q^{2R}. \quad (2.97)$$

If we want to think of this algebra as embedded into $\mathfrak{psu}(1,1|2)_L \oplus \mathfrak{psu}(1,1|2)_R$, it then follows that the two copies of $\mathfrak{psu}(1,1|2)$ are in *different gradings* with respect to each other. As we discuss in appendix A.1, this nicely ties to our choice of different supermatrix representatives for the left and right copies of $\mathfrak{psu}(1,1|2)$.

Off-shell representations

Let us now consider an excitation of arbitrary momentum p , *e.g.* a left excitation. Now this is charged under all of the supercharges—the left ones act proportionally to f_p , and the right ones to g_p . However, no supercharge can transform it into a right excitation, since nowhere in (2.87–2.88) appears an operator of the form $a_R^\dagger a_L$. The left representation is then an irreducible representation of the whole $\mathfrak{psu}(1|1)_{c.e.}^4$.

Clearly, the same reasoning can be applied to any right-moving excitation. Therefore, fundamental massive particles of the superstring transform into two irreducible (L and R) representations of the off-shell symmetry algebra.

Tensor-product structure

From the form of the on-shell representation depicted in figure 2.2 it is easy to see that we can describe the symmetries by means of a tensor product structure, whose factors are related to the indices 1 and 2 of the supercharges. We then can obtain the bifundamental representation of *e.g.* $\mathfrak{psu}(1|1)_L^2$ from the tensor product of two fundamental representations of a $\mathfrak{psu}(1|1)_L$.




Figure 2.3: The action of the supercharges $q^{L,R}$ and $\bar{q}^{L,R}$ of $\mathfrak{psu}(1|1)_L \oplus \mathfrak{psu}(1|1)_R$ centrally extended on an arbitrary momentum (off-shell) excitation (2.102).

Moreover, a tensor product structure exists in the off-shell algebra too. To see this let us introduce bosonic and fermionic operators $a_{L,R}^\dagger, a_{L,R}$ and $d_{L,R}^\dagger, d_{L,R}$, respectively. They obey canonical (anti)commutation relations

$$[a_L(p), a_L^\dagger(\tilde{p})] = \delta(p - \tilde{p}), \quad \{d_L(p), d_L^\dagger(\tilde{p})\} = \delta(p - \tilde{p}), \quad (2.98)$$

and similarly for $L \leftrightarrow R$. By means of them we can define the supercharges

$$q^L = \int dp (f_p d_L^\dagger a_L - g_p a_R^\dagger d_R), \quad q^R = \int dp (f_p d_R^\dagger a_R - g_p a_L^\dagger d_L), \quad (2.99)$$

and their Hermitian conjugates \bar{q}^L and \bar{q}^R . It is easy to verify that these satisfy a centrally extended $\mathfrak{psu}(1|1)^2$ algebra given by

$$\begin{aligned} \{q^L, \bar{q}^L\} &= \frac{1}{2}\mathfrak{h} + \frac{1}{2}\mathfrak{m}, & \{q^R, \bar{q}^R\} &= \frac{1}{2}\mathfrak{h} - \frac{1}{2}\mathfrak{m}, \\ \{q^L, q^R\} &= -\frac{1}{2}\mathfrak{p}, & \{\bar{q}^L, \bar{q}^R\} &= -\frac{1}{2}\mathfrak{p}, \end{aligned} \quad (2.100)$$

where the central charges are

$$\begin{aligned} \mathfrak{h} &= \int dp \omega_p (a_L^\dagger a_L + d_L^\dagger d_L + a_R^\dagger a_R + d_R^\dagger d_R), \\ \mathfrak{m} &= \int dp (a_L^\dagger a_L + d_L^\dagger d_L - a_R^\dagger a_R - d_R^\dagger d_R), \\ \mathfrak{p} &= \int dp p (a_L^\dagger a_L + d_L^\dagger d_L + a_R^\dagger a_R + d_R^\dagger d_R). \end{aligned} \quad (2.101)$$

Let us introduce excitations

$$|\phi^{L,R}(p)\rangle = a_{L,R}^\dagger(p) |0\rangle, \quad |\psi^{L,R}(p)\rangle = d_{L,R}^\dagger(p) |0\rangle. \quad (2.102)$$

On shell we have $f_{p=0} = 1$ and $g_{p=0} = 0$, so that the excitations indeed transform in two fundamental representations of $\mathfrak{su}(1|1)_L$ and $\mathfrak{su}(1|1)_R$, respectively. Off-shell, these two representations get deformed as depicted in figure 2.3.

The symmetry algebra of our worldsheet theory is in fact just given by the tensor product of two copies of (2.99) together with the relative central charges. To see this,

note that the annihilation operators are

$$a_L^{++} = a_L \otimes a_L, \quad a_L^{--} = d_L \otimes d_L, \quad a_L^{+-} = a_L \otimes d_L, \quad a_L^{-+} = d_L \otimes a_L, \quad (2.103)$$

and similarly for the creation operators and for $L \leftrightarrow R$. When we want to emphasise that the excitations $\phi^{L,R}$ and $\psi^{L,R}$ are to be taken as elements of a tensor product we will instead denote them as

$$|\Phi_+^{L,R}\rangle = |\phi^{L,R}\rangle, \quad |\Phi_-^{L,R}\rangle = |\psi^{L,R}\rangle, \quad (2.104)$$

so that $\Phi_{\kappa L}^L = \Phi_{\kappa}^L \otimes \Phi_L^L$, and $\Phi_{\kappa L}^R = \Phi_{\kappa}^R \otimes \Phi_L^R$. For the charges, we have

$$\mathbb{Q}^{1L} = q^L \otimes \mathbb{I}, \quad \mathbb{Q}^{2L} = \mathbb{I} \otimes q^L, \quad \mathbb{Q}^{1R} = q^R \otimes \mathbb{I}, \quad \mathbb{Q}^{2R} = \mathbb{I} \otimes q^R, \quad (2.105)$$

and similarly for the $\overline{\mathbb{Q}}$'s. The tensor product here should respect the \mathbb{Z}_2 grading, and yield a minus sign when two fermionic operators are swapped. By taking this into account we see that the charges defined in this way precisely agree with (2.87–2.88).

This scenario is quite similar to what happens for the $\text{AdS}_5 \times S^5$ superstring. In that case, the off-shell algebra is given by two copies of $\mathfrak{su}(2|2)$ centrally extended, and the excitations transform in a representation that is the tensor product of two fundamental representations of that algebra. In our case, $\mathfrak{su}(1|1)_L \oplus \mathfrak{su}(1|1)_R$ plays the role of $\mathfrak{su}(2|2)$. Excitations now transform in *two distinct representations* that are once again of the tensor product form. This will have important consequences for the form of the S matrix that we will compute in the next section. Before moving to that, let us further investigate the central charges appearing in the off-shell algebra.

The central charges

Studying the symmetry algebra at quadratic order in the fields we have seen that it has the form of a subalgebra of the original isometries, supplemented by two central charges that we will denote by \mathbb{C} and $\overline{\mathbb{C}}$ and that vanish on-shell. At quadratic order it so happens that these two central charges are equal and proportional to the worldsheet momentum \mathbb{P} . In particular, in terms of the fields we had *e.g.*

$$\{\mathbb{Q}^{jL}, \mathbb{Q}^{kR}\} = -\frac{1}{2} \delta^{jk} \int d\sigma x'_-, \quad (2.106)$$

at leading order.

While—as the coset construction highlighted—the rest of the algebra is fixed by the embedding into $\mathfrak{psu}(1,1|2)^2$, these additional central charges are not, and it is important to understand how they are modified if we consider higher order terms in our field expansion. In particular, we would like to understand

- whether there is any additional central charge that vanishes on-shell and appearing at higher order in the expansion,

- and what is the form of $\mathbb{C}, \bar{\mathbb{C}}$ when we account for higher order terms.

The former point is negatively answered by observing that the one we are considering is already the *maximal* non-trivial central extension of $\mathfrak{psu}(1,1)^4$. As for the latter point, the scenario we have here is very similar to the one found in $\text{AdS}_5 \times S^5$, and in fact find an answer by repeating an argument employed there, which spares us the—hardly feasible—computation of the charges at very high orders in the field expansion.

The key observation is that in the coset construction the unphysical coordinate x_- is neatly packaged in $\Lambda(x_{\pm})$ so that the supercharges can be cast in the general form

$$\mathbb{Q} = \int d\sigma e^{+\frac{i}{2}x_-} \left(\theta \cdot (\mathcal{G}^{(1)}(p, x) + \frac{1}{h}\mathcal{G}^{(3)}(p, x) + \dots) + \mathcal{O}(\theta^3) \right), \quad (2.107)$$

see also appendix A.3. Here we have expanded the charge density first order by order in the fermions, and then we have expanded the coefficient of each such term in $1/h$. Therefore, $\mathcal{G}^{(1)}(p, x)$ is linear in the bosons, $\mathcal{G}^{(3)}(p, x)$ is cubic, and so on. Since the central charges should vanish on-shell, they should be a function of \mathbb{P} which vanishes at zero. In particular, if we find the functional dependence of \mathbb{C} and $\bar{\mathbb{C}}$ on the bosonic fields alone, we will be able to unambiguously fix their full form [57]. The bosonic part of the Poisson bracket of two supercharges is then *e.g.*

$$\{\mathbb{Q}^{jL}, \mathbb{Q}^{kR}\} \approx \delta^{jk} \int d\sigma e^{ix_-} \left(\mathcal{G}_{jL}^{(1)} \mathcal{G}_{kR}^{(1)} + \frac{1}{h} (\mathcal{G}_{jL}^{(1)} \mathcal{G}_{kR}^{(3)} + \mathcal{G}_{jL}^{(3)} \mathcal{G}_{kR}^{(1)}) + \dots \right). \quad (2.108)$$

In fact, the leading order term is precisely what we have already calculated, that is x'_- restricted to the bosonic fields only. Therefore, at the leading order of this hybrid expansion we find

$$\{\mathbb{Q}^{jL}, \mathbb{Q}^{kR}\} = -\frac{\hbar}{2} \delta^{jk} \int d\sigma e^{ix_-} x'_- = i\frac{\hbar}{2} (e^{ix_- (+\infty)} - e^{ix_- (-\infty)}) \delta_{jk}. \quad (2.109)$$

Repeating this calculation to include higher orders of the bosonic expansion, one can check that these do not spoil the form of eq. (2.109). It is actually convenient to rewrite this in a way that makes the dependence on the worldsheet momentum manifest,⁸

$$\{\mathbb{Q}^{jL}, \mathbb{Q}^{kR}\} = \delta^{jk} \mathbb{C} = i \delta^{jk} \zeta \frac{\hbar}{2} (e^{i\mathbb{P}} - 1), \quad (2.110)$$

where we isolated the boundary condition $\zeta = e^{ix_- (-\infty)}$. In fact, on the one-particle representation we can consistently set *e.g.* $x_-(-\infty) = 0$ so that $\zeta = 1$. A similar calculation shows that, as required by hermiticity

$$\{\bar{\mathbb{Q}}^{jL}, \bar{\mathbb{Q}}^{kR}\} = \delta_{jk} \bar{\mathbb{C}} = -i \delta_{jk} \bar{\zeta} \frac{\hbar}{2} (e^{-i\mathbb{P}} - 1). \quad (2.111)$$

Of course the non-linearity of $\mathbb{C}, \bar{\mathbb{C}}$ in the off-shell algebra means that the same must be true in each copy of the tensor product for \mathfrak{c} and $\bar{\mathfrak{c}}$, while the last line of (2.100) holds only at leading order in $1/h$.

⁸Eq. (2.91) can be recovered by performing the rescaling $\mathbb{P} \rightarrow \mathbb{P}/h$ and expanding in $1/h$.

2.5 Chapter summary

The main result of this chapter is that we derived the off-shell symmetries of the massive excitations of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ from a perturbative analysis of the NLSM action. The symmetry algebra is given by two copies of a central extension of $\mathfrak{su}(1|1)_L \oplus \mathfrak{su}(1|1)_R$, which has anticommutation relations

$$\begin{aligned} \{\mathfrak{q}^L, \bar{\mathfrak{q}}^L\} &= \frac{1}{2}\mathfrak{h} + \frac{1}{2}\mathfrak{m}, & \{\mathfrak{q}^R, \bar{\mathfrak{q}}^R\} &= \frac{1}{2}\mathfrak{h} - \frac{1}{2}\mathfrak{m}, \\ \{\mathfrak{q}^L, \mathfrak{q}^R\} &= \mathfrak{c}, & \{\bar{\mathfrak{q}}^L, \bar{\mathfrak{q}}^R\} &= \bar{\mathfrak{c}}. \end{aligned} \quad (2.112)$$

The central charges $\mathfrak{c}, \bar{\mathfrak{c}}$ vanish on-shell, and otherwise are non-linear functions of the worldsheet momentum \mathfrak{p} , which for an appropriate choice of boundary conditions can be written as

$$\mathfrak{c} = +i\frac{\hbar}{2}(e^{+i\mathfrak{p}} - 1), \quad \bar{\mathfrak{c}} = -i\frac{\hbar}{2}(e^{-i\mathfrak{p}} - 1). \quad (2.113)$$

The role of the central extensions for $\mathfrak{su}(1|1)^2$ was originally discussed in ref. [121], in the context of $\text{AdS}_5/\text{CFT}_4$ duality, and in the case of $\text{AdS}_3/\text{CFT}_2$ in refs. [50, 122] and more recently in ref. [102].⁹ We found two one-particle representation of the algebra (2.112), which are deformations of the fundamental representation of $\mathfrak{psu}(1|1)$: the “left” representation (L) has a bosonic highest weight state, whereas the “right” representation (R) has a fermionic one. Massive particles of the worldsheet theory are in two tensor product representations, $L \otimes L$ and $R \otimes R$. This accounts for all of the $8 = 4_L + 4_R$ massive excitations.

Some remarks are in order. Firstly, owing to the coset construction, even if our approach is perturbative we were able to capture the general form of the algebra including the non-linear central extension. We should still bear in mind that our calculation was entirely classical, and in principle some of these result might still be spoiled by quantum anomalies.

To exploit the coset formulation we “froze” all the massless excitations. Could taking them into account modify the symmetries we identified? We know that the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ backgrounds has 16 supersymmetries, and that the choice of light-cone gauge will break at least half of those. Therefore, we cannot expect any new supercharge to appear from including the massless modes. In principle, we may be overlooking some bosonic symmetries that should supplement the algebra we found. Still, we know that the off-shell theory has *at least* the symmetries given by two copies of (2.112). Of course this does not say anything on what are the symmetries of the *massless* excitations. We do know that since \mathbb{H} (and therefore the mass) appears in the symmetry algebra, the massless excitations cannot transform in the same representation as the massive ones. We will come back to this in chapter 8.

⁹The same symmetry algebra was also found in the analysis of the Pohlmeyer reduced sigma model of the $\text{AdS}_3 \times \text{S}^3$ in ref. [123].

Let us finally stress that even if physical excitations transform under the on-shell symmetry algebra, our interest lies mainly in the off-shell symmetries, which are the ones that constrain the S matrix.

3 The all-loop integrable S matrix

In this chapter we will conjecture the scattering matrix for fundamental massive excitations of the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ superstring. To do so, we will rely on two results:

- the off-shell symmetry algebra found in the previous chapter,
- and the fact that the underlying classical field theory enjoys a large amount of symmetries that make it “integrable” [50–52].¹

We will supplement these fact by two assumptions, namely

- that the off-shell symmetries which we found persist in the full quantum theory,
- and that the integrable structure also extends to the quantum theory.

In this way, we will be able to derive an essentially unique S matrix \mathbb{S} , up to some prefactors—the so-called dressing factors. Our first task will be to briefly explore the idea of integrability in the classical and quantum theory, and derive restrictions on the resulting S matrix. Then we will formulate the off-shell symmetries at the level of the S matrix. This will be sufficient to find an \mathbb{S} which will have several desirable physical properties, guaranteeing the self-consistency of our procedure, as we will see. Ultimately, the validity of our assumptions will have to be tested by comparing the S matrix against perturbative calculations. We will come back to this in chapter 7.

3.1 Classical and quantum integrability

Integrability is a broad concept that first emerged in the context of Hamiltonian systems and later was extended to classical and quantum field theories. There is no universal mathematical definition of what an integrable theory is. Colloquially, an integrable theory is one that enjoys so much symmetry so that its dynamics is completely constrained and can be solved “exactly”. Here we will try to make this idea more precise. Note that our discussion of this general framework will be quite essential. We refer the reader to *e.g.* refs. [125, 126] for further details on classical integrability and the inverse scattering method and to refs. [127–129] for scattering factorisation and quantum integrable theories.

¹Classical integrability is also seen by studying the Gubser-Klebanov-Polyakov “spinning string” [124].




Figure 3.1: In classical mechanics, the Liouville-Arnol'd theorem establishes that, when enough conserved quantities in involution exist, the phase space is foliated in tori where the dynamics is supported. Here we depict such a foliation, with the arrow representing the motion of a particle.

Classical integrable theories

In the case of an Hamiltonian system with M degrees of freedom $(p_1, q_1, \dots, p_M, q_M)$, integrability can be defined through the Liouville-Arnol'd theorem. This states that if there exists M Poisson-commuting independent quantities $\mathbb{F}_1, \dots, \mathbb{F}_M$, then there exists a foliation of the phase space into invariant tori² on which the motion is supported, as depicted in figure 3.1. Furthermore there exists a canonical transformation to action-angle variables $(K_1, \varphi_1, \dots, K_M, \varphi_M)$ that takes the equations of motion to the form

$$\dot{K}_j = 0, \quad \dot{\varphi}_j = 1, \quad j = 1, \dots, M. \quad (3.1)$$

Typically, we know only one of the \mathbb{F}_j 's, which is the Hamiltonian. The other charges can sometimes be found from the geometry of the problem. When that is possible—which happens quite exceptionally—even if the Hamiltonian appears to be highly non-linear, the theorem guarantees that we are in fact dealing with a simple system “in disguise”—such as the Kepler problem or the Euler top.

The idea of having enough commuting conserved quantities is at the hearth of integrability, but it is not the most convenient approach to extend it to classical field theory. In that case there are infinitely many degrees of freedom, so that we would need to exhibit infinitely many quantities in involution. This may be possible, but how many should we actually produce? Two infinite sets of charges may have the same cardinality while one is strictly included in the other. A way around this complication is given by the *Lax formalism*³. Consider a two-dimensional classical field theory, and assume that the resulting equations of motion can be cast in the form

$$\partial_\tau U(\tau, \sigma, x) - \partial_\sigma V(\tau, \sigma, x) - [V(\tau, \sigma, x), U(\tau, \sigma, x)] = 0, \quad (3.2)$$

²If the orbits are compact. The generalisation to non-compact orbits is straightforward.

³Such a formalism can also be very useful when dealing with finite dimensional integrable systems, but this is beyond the purpose of our discussion.

where U, V are matrices depending on the fields and on the complex parameter x . Let us define the *monodromy matrix* $T(\tau, x)$ by the path-ordered exponential

$$T(\tau, x) = \overleftarrow{\exp} \left(\int_{-\ell/2}^{\ell/2} d\sigma U(\tau, \sigma, x) \right). \quad (3.3)$$

An explicit computation shows that this obeys the evolution equation

$$\partial_\tau T(\tau, x) = [V(\tau, -\ell/2, x), T(\tau, x)]. \quad (3.4)$$

As a consequence, $\partial_\tau \text{tr}[T(\tau, x)^j] = 0$ and all of the eigenvalues of the monodromy matrix are conserved by the time evolution. In fact, an object that nicely encodes all of these conserved quantities is the complex curve defined by the eigenvalue equation

$$\Gamma(x, \mu(x)) = 0, \quad \Gamma(x, \mu) = \det(T(\tau, x) - \mu \mathbf{1}), \quad (3.5)$$

where $\mu(x)$ is such as to solve the leftmost equation. The resulting $\Gamma(x)$ is called the *spectral curve*.

On top of this set of conserved quantities, the Lax formulation guarantees more. Without loss of generality, we may think of the equations of motion (3.2) as arising from a linear system⁴

$$\begin{aligned} \partial_\sigma \Psi(\tau, \sigma, x) &= U(\tau, \sigma, x) \Psi(\tau, \sigma, x), \\ \partial_\tau \Psi(\tau, \sigma, x) &= V(\tau, \sigma, x) \Psi(\tau, \sigma, x), \end{aligned} \quad (3.6)$$

by requiring the compatibility condition $(\partial_\sigma \partial_\tau - \partial_\tau \partial_\sigma) \Psi = 0$. This gives us the means of solving the non-linear equations (3.2) in terms of linear ones—a procedure sometimes called *the inverse scattering method* [130–133]. The matrices U, V are functions of the fields, and in particular $U(0, \sigma, x)$ is completely determined once we fix the initial conditions for the fields at $\tau = 0$. From this, finding $\Psi(0, \sigma, x)$ amounts to solving a linear equation. So does finding the time evolution for $\Psi(\tau, \sigma, x)$. Once this is determined, the only missing step is to reconstruct $U(\tau, \sigma, x)$ (and therefore the value of the fields at any given τ) from $\Psi(\tau, \sigma, x)$. This can be done by solving an integral equation, the Gel'fand-Levitan-Marchenko equation [134, 135], which again is linear.

In conclusion, the mere fact that the equations of motion for a two dimensional classical field theory can be cast in the form (3.2) implies that the system has infinitely many conserved charges and that its equations of motions are relatively simple to solve. Moreover, for such systems it is generally possible to relate the matrix structure with the Poisson structure of the original Hamiltonian description in such a way as to guarantee that the conserved quantities Poisson-commute.

⁴In fact, the monodromy matrix takes its name from representing the monodromy of a solution of this linear system, $\Psi(\tau, r, x) = T(\tau, x) \Psi(\tau, -r, x)$.

Integrability for two-dimensional QFTs

Let us assume that we can quantise an integrable theory as the one described before, and that we can do so without spoiling its infinitely many symmetries. Let

$$\mathbb{F}_1, \dots, \mathbb{F}_n, \dots, \quad [\mathbb{F}_j, \mathbb{F}_k] = 0 \quad (3.7)$$

be the set of commuting conserved charges. These charges can be simultaneously diagonalised. For definiteness, let us suppose that this happens in a basis where Hilbert space states are identified by their momentum and a flavour label α . Then we will have

$$\mathbb{F}_n |p\rangle_\alpha^{(\text{in}, \text{out})} = F_n(p; \alpha) |p\rangle_\alpha^{(\text{in}, \text{out})}. \quad (3.8)$$

For instance, in the particular case of the sine-Gordon model where there is only one scalar excitation of mass m , one would have

$$F_{2n+1}(p) = p^{2n+1}, \quad F_{2n}(p) = p^{2n} \sqrt{p^2 + m^2}, \quad (3.9)$$

which give *higher charges* generalizing momentum and energy. In a more general theory, the higher charges will feature some invariant tensor constructed out of the labels α and infinitely many functions of the particle momentum.

Let us consider the action of such a charge on an M -particle state. Considering *e.g.* an in-state, we have

$$\mathbb{F}_n |p_1, \dots, p_M\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in})} = (F_n(p_1; \alpha_1) + \dots + F_n(p_M; \alpha_M)) |p_1, \dots, p_M\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in})}. \quad (3.10)$$

After we evolve the state $|p_1, \dots, p_M\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in})}$ we obtain some corresponding out-state $|\tilde{p}_1, \dots, \tilde{p}_{\tilde{M}}\rangle_{\tilde{\alpha}_1, \dots, \tilde{\alpha}_{\tilde{M}}}^{(\text{out})}$. Since the charges are conserved, it must be that

$$\sum_{j=1}^M F_n(p_j; \alpha_j) = \sum_{k=1}^{\tilde{M}} F_n(\tilde{p}_k; \tilde{\alpha}_k). \quad (3.11)$$

The only way for these sums to be equal *for all of the* $F_n(p)$ is if the set of “in” momenta appearing $\{p_j\}$ corresponds to the set of “out” momenta $\{\tilde{p}_k\}$. In particular, it must be $M = \tilde{M}$.

This scenario, together with the peculiar topology of two-dimensional QFTs, has deep implications for a scattering event. Let us start by considering a two-particle scattering event. The in-state is $|p_1, p_2\rangle_{\alpha_1, \alpha_2}^{(\text{in})}$. Since this state is defined at time $\tau = -\infty$, momenta are ordered so that $p_1 > p_2$. As time evolves, the particles move on a line until at some point they come together and scatter. After the scattering, the products move away from each other and can again be considered as two real particles. Because of the conservation law (3.11), the resulting state is proportional to $|p_2, p_1\rangle_{\tilde{\alpha}_2, \tilde{\alpha}_1}$, where the momenta are exactly the same and the labels $\tilde{\alpha}_2, \tilde{\alpha}_1$ may have changed.




Figure 3.2: Different ways of resolving the $3 \rightarrow 3$ particle scattering (central panel). Taking time to flow upwards, we have in the left panel the sequence of scattering (12)-(13)-(23), while in the right panel we have (23)-(13)-(12). The Yang-Baxter equation imposes the equality of the two resolutions.

This can be readily extended to a M -particle in-state $|p_1, \dots, p_M\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in})}$ with $p_1 > p_2 > \dots > p_M$. At some point two of the particles undergo a $2 \rightarrow 2$ scattering like the one described above. After that, the we are left with M real (as opposed to virtual) particles propagating on a line. After a sequence of $\frac{1}{2}M(M-1)$ scattering events, the particles are spatially ordered as in the out-state, having momenta p_M, p_{M-1}, \dots, p_1 from left to right. Therefore, an M -particle scattering event *factorises* into a sequence of two-particle events, and was first put forward by the Zamolodchikov brothers in the seminal paper [128]. This is the special property of the multiparticle S matrix that we alluded to in the introduction. The object that we need to determine is only the two-particle S matrix $\mathbb{S}(p_1, p_2)$.

For internal consistency, we have to require some conditions on the S matrix. The most obvious one is that there is no $2 \rightarrow M$ scattering unless $M = 2$, in sharp contrast with what happens *e.g.* in a typical particle collider experiment. Another obstacle is that there are several apparently inequivalent ways of resolving an $M \rightarrow M$ scattering in terms of a sequence of $2 \rightarrow 2$ ones. In figure 3.2 we depict a generic $3 \rightarrow 3$ particle scattering, and the two ways of resolving it. We must require that these two sequences of scattering events yield the same result. Each of them gives a cubic expression in the S matrix, and their equality results in a non-linear matrix equation—the *Yang-Baxter equation*. The physical reason for the equivalence of the pictures in fig. 3.2 can be traced back to the existence of the higher conserved charges \mathbb{F}_n . These generate unitary transformations that shift the momentum p_j of a particle by an amount of order $(p_j)^{n-1}$, and effectively transform the leftmost panel of fig. 3.2 into the rightmost one.

To better investigate the constraints that $\mathbb{S}(p_1, p_2)$ should satisfy it is convenient to introduce a more formal algebraic framework.

The Zamolodchikov-Faddeev algebra

The Zamolodchikov-Faddeev (ZF) algebra [128, 136] is a tool to encode the integrability properties of a two-dimensional QFT. It is defined in terms of abstract raising and lowering operators $A_\alpha^\dagger(p)$ and $A^\alpha(p)$ which create or destroy a particle of definite momentum and flavour. Their action on the vacuum is

$$\begin{aligned} A^\alpha(p) |0\rangle &= 0, \\ A_\alpha^\dagger(p) |0\rangle &= |p\rangle_\alpha^{(\text{in})} = |p\rangle_\alpha^{(\text{out})}, \end{aligned} \quad (3.12)$$

where we used the fact that in an integrable theory like the one we want to describe a one-particle state undergoes a trivial time evolution. Following the discussion of the previous section, it is natural to relate the two-particle in- and out-states as

$$\begin{aligned} |p_1, p_2\rangle_{\alpha_1, \alpha_2}^{(\text{in})} &= A_{\alpha_1}^\dagger(p_1) A_{\alpha_2}^\dagger(p_2) |0\rangle, \\ |p_1, p_2\rangle_{\alpha_3, \alpha_4}^{(\text{out})} &= (-1)^{\varepsilon(\alpha_3)\varepsilon(\alpha_4)} A_{\alpha_3}^\dagger(p_2) A_{\alpha_4}^\dagger(p_1) |0\rangle, \end{aligned} \quad \text{with } p_1 > p_2. \quad (3.13)$$

Notice that we used that the set of momenta is conserved and we ordered the action of the creation and annihilation operators according to the ordering of the particles in the final states. We have also written explicitly the sign arising from the exchange of two fermions, where $\varepsilon(\alpha) = 1$ for fermionic particles and zero otherwise.

The two states in (3.13) are precisely the ones related by the S matrix from (1.18) in the $2 \rightarrow 2$ case

$$|p_1, p_2\rangle_{\alpha_1, \alpha_2}^{(\text{in})} = S_{\alpha_1, \alpha_2}^{\alpha_3, \alpha_4}(p_1, p_2) |p_1, p_2\rangle_{\alpha_3, \alpha_4}^{(\text{out})}, \quad (3.14)$$

where we indicated by $S_{j_1, j_2}^{j_3, j_4}$ the matrix element of \mathbb{S} . This equation, together with (3.13), yields the commutation relations for the ZF operators

$$A_{\alpha_1}^\dagger(p_1) A_{\alpha_2}^\dagger(p_2) = (-1)^{\varepsilon(\alpha_3)\varepsilon(j_4)} S_{\alpha_1, \alpha_2}^{\alpha_3, \alpha_4}(p_1, p_2) A_{\alpha_3}^\dagger(p_2) A_{\alpha_4}^\dagger(p_1), \quad (3.15)$$

which makes it evident that A^\dagger, A satisfy a *different algebra* than the one of the canonical raising and lowering operators a^\dagger, a .

To avoid carrying around too many indices it is useful to introduce a matrix basis. Let E_j and E^j be rows and column vectors with a single non-vanishing unit entry in the j th place. These give bases of the two dual vector spaces \mathcal{V}^* and \mathcal{V} . A basis of the space of matrices on \mathcal{V} is given by $E_j^k = E_j \otimes E^k$, so that by indices are naturally contracted to give

$$E^l E_k^j = \delta_k^l E^j, \quad E_l E_j^k = \delta_k^l E_j, \quad E_k^l E_i^j = \delta_i^l E_k^j. \quad (3.16)$$

We can then introduce vector and row operators of the ZF algebra

$$\mathbb{A}^\dagger = \mathbb{A}^\dagger(p) = A_\alpha^\dagger(p) E^\alpha, \quad \mathbb{A} = \mathbb{A}(p) = A^\alpha(p) E_\alpha. \quad (3.17)$$

It is also useful to reabsorb a permutation and the fermionic signs into the \mathbb{R} matrix

$$\mathbb{R}(p, q) = R_{\alpha_1, \alpha_2}^{\alpha_3, \alpha_4}(p, q) E_{\alpha_3}^{\alpha_1} \otimes E_{\alpha_4}^{\alpha_2}, \quad R_{\alpha_1, \alpha_2}^{\alpha_4, \alpha_3}(p, q) = (-1)^{\varepsilon(\alpha_3)\varepsilon(\alpha_4)} S_{\alpha_1, \alpha_2}^{\alpha_3, \alpha_4}(p, q), \quad (3.18)$$

so that \mathbb{R} differs from \mathbb{S} by a graded permutation Π^g , $\mathbb{R} = \Pi^g \mathbb{S}$. In these terms the commutation relations (3.15) take the form

$$\mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger = \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(1)}^\dagger \mathbb{R}_{(12)}, \quad (3.19)$$

where we added subscript indices specifying on which factors of the tensor product $\mathcal{V} \otimes \mathcal{V}$ the operators act.

In the same way as we have obtained (3.15) we can now derive similar relations involving the annihilation operators, thus completing the ZF algebra

$$\begin{aligned} \mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger &= \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(1)}^\dagger \mathbb{R}_{(12)}, & \mathbb{A}_{(1)} \mathbb{A}_{(2)} &= \mathbb{R}_{(12)} \mathbb{A}_{(2)} \mathbb{A}_{(1)}, \\ \mathbb{A}_{(1)} \mathbb{A}_{(2)}^\dagger &= \mathbb{A}_{(2)} \mathbb{R}_{(21)} \mathbb{A}_{(1)}^\dagger + \delta(p_1 - p_2) \mathbb{I}. \end{aligned} \quad (3.20)$$

Consistency conditions

The algebra (3.20) in principle constitutes a tool to express an in-particle state in terms of the out-particle basis and *vice versa*, *i.e.* a way of computing S-matrix elements. For this to be true, however, we have to impose some further consistency conditions. We immediately find that

$$\mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger = \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(1)}^\dagger \mathbb{R}_{(12)} = \mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger \mathbb{R}_{(21)} \mathbb{R}_{(12)}, \quad (3.21)$$

so that it must be

$$\mathbb{R}_{(21)} \mathbb{R}_{(12)} = \mathbb{R}_{(12)} \mathbb{R}_{(21)} = \mathbb{I}. \quad (3.22)$$

This condition, called *braiding unitarity*, supplements the usual *physical unitarity* condition, which states that \mathbb{S} and therefore \mathbb{R} should be unitary as a matrix:

$$\mathbb{R}_{(12)} \mathbb{R}_{(12)}^\dagger = \mathbb{R}_{(12)}^\dagger \mathbb{R}_{(12)} = \mathbb{I}. \quad (3.23)$$

So far we have used the ZF algebra only on two-particle states. We now want to extend it to arbitrary multiparticle states in such a way as to implement the factorisation of the scattering. To do so, we extend the definition (3.13) by

$$\begin{aligned} |p_1, \dots, p_M\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{in})} &= A_{\alpha_1}^\dagger(p_1) \cdots A_{\alpha_M}^\dagger(p_M) |0\rangle, \\ |p_1, \dots, p_M\rangle_{\alpha_1, \dots, \alpha_M}^{(\text{out})} &= (-1)^{\sum_{k < l} \varepsilon(\alpha_k) \varepsilon(\alpha_l)} A_{\alpha_1}^\dagger(p_M) \cdots A_{\alpha_M}^\dagger(p_1) |0\rangle, \end{aligned} \quad (3.24)$$

again with $p_1 > p_2 > \cdots > p_M$. If we want to use (3.20) to express in-states in terms of out-states, we are faced with an ambiguity. Consider, for instance, the combination $\mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(3)}^\dagger$ acting on $\mathcal{V} \otimes \mathcal{V} \otimes \mathcal{V}$. We can rewrite it in two different ways:

$$\begin{aligned} \mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(3)}^\dagger &= \mathbb{A}_{(3)}^\dagger \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(1)}^\dagger \mathbb{R}_{(12)} \mathbb{R}_{(13)} \mathbb{R}_{(23)}, \\ \mathbb{A}_{(1)}^\dagger \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(3)}^\dagger &= \mathbb{A}_{(3)}^\dagger \mathbb{A}_{(2)}^\dagger \mathbb{A}_{(1)}^\dagger \mathbb{R}_{(23)} \mathbb{R}_{(13)} \mathbb{R}_{(12)}. \end{aligned} \quad (3.25)$$

These coincide only provided that

$$\mathbb{R}_{(12)}\mathbb{R}_{(13)}\mathbb{R}_{(23)} = \mathbb{R}_{(23)}\mathbb{R}_{(13)}\mathbb{R}_{(12)}. \quad (3.26)$$

This is the *Yang-Baxter equation* that we described in the previous subsection. In fact, comparing with figure 3.2 one can see that the product of the matrices appearing in the equation precisely corresponds to the sequences of scattering events depicted there. It is then straightforward to see that, by repeatedly using the Yang-Baxter equation, it is possible to rearrange a string of ZF operators of arbitrary length, which ensures factorisation of any multiparticle scattering.

Symmetries

In our subsequent study, it will be important to make use of the transformation properties of S matrix under the off-shell symmetry algebra.

The simplest case is given by the central charges, which make up the whole bosonic part of the off-shell symmetries. Let us consider a charge that is proportional to the number operator

$$\mathbb{X}_f = \int dp f(p) \mathbb{A}^\dagger(p) \mathbb{A}(p), \quad (3.27)$$

where $f(p)$ is an arbitrary function of the momentum. Using braiding unitarity it is easy to prove [29] that \mathbb{X}_f must satisfy the relations

$$\mathbb{X}_f \mathbb{A}^\dagger(p) = \mathbb{A}^\dagger(p) (f(p) + \mathbb{X}_f), \quad \mathbb{X}_f \mathbb{A}(p) = \mathbb{A}(p) (-f(p) + \mathbb{X}_f). \quad (3.28)$$

Therefore, these functions must form an abelian subalgebra of the ZF algebra. In particular, we expect the worldsheet momentum, the Hamiltonian and the two central charges found in the previous section to be part of it.

Let us now consider a more general non-abelian (super)algebra \mathcal{G} , and let us assume that the one-particle vector space \mathcal{V} carries one or more irreducible representations of \mathcal{G} . Eventually we will identify \mathcal{G} with the full off-shell algebra. It is then natural to take \mathcal{G} to commute with the worldsheet momentum and the particle number⁵ as well as with all the higher conserved charges—compatibly with our assumption of integrability. Then we can write the action a symmetry generator $\mathbb{Q} \in \mathcal{G}$ on the zero-, one-, two-particle Hilbert spaces, *etc.* as

$$\begin{aligned} \mathbb{Q} \cdot |0\rangle &= 0, \\ \mathbb{Q} \cdot A_\alpha^\dagger(p) |0\rangle &= Q_\alpha^\beta(p) A_\beta^\dagger(p) |0\rangle, \\ \mathbb{Q} \cdot A_{\alpha_1}^\dagger(p) A_{\alpha_2}^\dagger(q) |\Omega\rangle &= Q_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p, q) A_{\beta_1}^\dagger(p) A_{\beta_2}^\dagger(q) |0\rangle, \end{aligned} \quad (3.29)$$

⁵Since we are dealing with a supersymmetric theory, we also require that $(-1)^{\mathcal{F}}$, where \mathcal{F} is the fermion number, to be conserved. Note that \mathcal{F} itself is generally not conserved, resulting in processes such as fermion-fermion \rightarrow boson-boson.

and so on.

Since \mathbb{Q}_α is a symmetry of the theory, we can simultaneously transform the in- and out-states without affecting the S-matrix elements. In particular using (3.13) in

$$\mathbb{Q} \cdot |p_1, p_2\rangle_{\alpha_1, \alpha_2}^{(\text{in})} = \mathbb{S} \cdot \mathbb{Q} \cdot |p_1, p_2\rangle_{\alpha_3, \alpha_4}^{(\text{out})}, \quad (3.30)$$

we get the invariance condition

$$\begin{aligned} S_{\beta_3, \beta_4}^{\alpha_3, \alpha_4}(p_1, p_2) Q_{\alpha_1, \alpha_2}^{\beta_3, \beta_4}(p_1, p_2) &= (-1)^{\varepsilon(\alpha_3)\varepsilon(\alpha_4) + \varepsilon(\beta_3)\varepsilon(\beta_4)} \\ &Q_{\beta_3, \beta_4}^{\alpha_3, \alpha_4}(p_2, p_1) S_{\alpha_1, \alpha_2}^{\beta_3, \beta_4}(p_1, p_2), \end{aligned} \quad (3.31)$$

which can be more compactly expressed in the matrix notation in terms of \mathbb{R} :

$$\mathbb{R}_{(12)}(p_1, p_2) \mathbb{Q}_{(12)}(p_1, p_2) = \mathbb{Q}_{(21)}(p_2, p_1) \mathbb{R}_{(12)}(p_1, p_2). \quad (3.32)$$

Let us further investigate the form of the structure constants. Since our symmetry algebra \mathcal{G} has a non-trivial centre, any of its irreducible representations is labelled by the value of the momentum and the other central charges $\{c\}$. Therefore we in principle we have to allow for⁶

$$\begin{aligned} Q_\alpha^\beta(p) &\rightarrow Q_\alpha^\beta(p; \{c\}), \\ Q_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p_1, p_2) &\rightarrow Q_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p_1, p_2; \{c_1, c_2\}). \end{aligned} \quad (3.33)$$

In a matrix notation, we can decompose the action of \mathbb{Q}_α on the factors of the tensor product as

$$\mathbb{Q}_{(12)}(p_1, p_2; \{c_1, c_2\}) = \mathbb{Q}(p_1, \{c_1\}) \otimes \mathbb{I} + \mathbb{I}^g (\mathbb{I} \otimes \mathbb{Q}(p_2, \{c_2\})) \mathbb{I}^g, \quad (3.34)$$

where \mathbb{I}^g is the graded identity matrix.

In order to further specify the form of this representation we will need to input some information on the symmetry algebra. We will do so in the next section.

3.2 Representations of $\mathfrak{su}(1|1)^2$ centrally extended

Rather than focusing on the whole off-shell symmetry algebra $\mathfrak{psu}(1|1)_{c.e.}^4$, let us focus on a single $\mathfrak{su}(1|1)_{c.e.}^2$. The commutation relations are given by (2.112), and we rewrite them here for convenience

$$\begin{aligned} \{\mathfrak{q}^L, \bar{\mathfrak{q}}^L\} &= \mathfrak{h}^L, & \{\mathfrak{q}^R, \bar{\mathfrak{q}}^R\} &= \mathfrak{h}^R, \\ \{\mathfrak{q}^L, \mathfrak{q}^R\} &= \mathfrak{c}, & \{\bar{\mathfrak{q}}^L, \bar{\mathfrak{q}}^R\} &= \bar{\mathfrak{c}}, \end{aligned} \quad (3.35)$$

where we introduce a left- and right-“Hamiltonian” $\mathfrak{h}^{L,R}$ so that

$$\mathfrak{h} = \mathfrak{h}^L + \mathfrak{h}^R, \quad \mathfrak{m} = \mathfrak{h}^L - \mathfrak{h}^R. \quad (3.36)$$

⁶This notation is a bit heavy, and we will keep the dependence on the central charges implicit where no confusion may arise.

We already know that how to represent this algebra on the space of left- and right-excitations $\phi^{L,R}, \psi^{L,R}$ when $\mathfrak{c} = \bar{\mathfrak{c}} = -\frac{1}{2}\mathbb{P}$. In terms of oscillators, that representation is given by (2.100) and (2.101). Here we want to deform that representation to allow for

$$\mathfrak{c} = +i\zeta \frac{h}{2}(e^{+i\mathbb{P}} - 1), \quad \bar{\mathfrak{c}} = -i\bar{\zeta} \frac{h}{2}(e^{-i\mathbb{P}} - 1). \quad (3.37)$$

This will amount to suitably deforming the parameters f_p, g_p and ω_p appearing in the representation. The deformed parameters will depend on the momentum p , the coupling constant h and the phase ζ that we found in (2.110).

The phase ζ has the meaning of a boundary condition on the unphysical field x_- in the coset model, $\zeta = e^{ix_-(-\infty)}$. When we want the representation of $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ to describe excitations of the superstring, we will set $\zeta = 1$. Then

$$\mathfrak{c} = +i \frac{h}{2}(e^{+i\mathbb{P}} - 1), \quad \bar{\mathfrak{c}} = -i \frac{h}{2}(e^{-i\mathbb{P}} - 1). \quad (3.38)$$

Shortening condition

In the previous chapter, we have found two irreducible representations of dimension two. These are both short (or atypical) representations of $\mathfrak{su}(1|1)_{\text{c.e.}}^2$. In fact, if we consider *e.g.* the left representation, we have that $|\phi^L\rangle$ is the highest weight state, annihilated by the two raising operators

$$\bar{\mathfrak{q}}^L |\phi^L\rangle = 0, \quad \mathfrak{q}^R |\phi^L\rangle = 0. \quad (3.39)$$

However, the highest weight is also annihilated by a combination of the lowering operators

$$(\mathfrak{h}^R \mathfrak{q}^L - \mathfrak{c} \bar{\mathfrak{q}}^R) |\phi^L\rangle = 0. \quad (3.40)$$

The vanishing of this combination of charges is the *shortening condition*. It also implies the vanishing of the anticommutator

$$0 = \{\bar{\mathfrak{q}}^L, \mathfrak{h}^R \mathfrak{q}^L - \mathfrak{c} \bar{\mathfrak{q}}^R\} = \mathfrak{h}^L \mathfrak{h}^R - \bar{\mathfrak{c}} \mathfrak{c}, \quad (3.41)$$

which we will also refer to as shortening condition.

Had we considered the right-representation, we would have found a fermionic highest weight state

$$\bar{\mathfrak{q}}^L |\psi^R\rangle = 0, \quad \mathfrak{q}^R |\psi^R\rangle = 0. \quad (3.42)$$

The same combination appearing in (3.40) annihilates $|\psi^R\rangle$. In fact, we can take (3.41) as shortening condition of both the left- and right-representations.

One-particle representation

Since the excitations span a space \mathcal{V} of dimension only four, it is quite handy to introduce a 4×4 matrix representation for the (super)charges. Let us pick a basis

$$\mathcal{B} = (\phi^L, \psi^L, \phi^R, \psi^R). \quad (3.43)$$

We can then make an ansatz for the supercharges in the one-particle representation,

$$\begin{aligned} \mathfrak{q}^L &= \left(\begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ a^L & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & \bar{b}^R \\ 0 & 0 & 0 & 0 \end{array} \right), & \mathfrak{q}^R &= \left(\begin{array}{cc|cc} 0 & b^L & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & a^R & 0 \end{array} \right), \\ \bar{\mathfrak{q}}^L &= \left(\begin{array}{cc|cc} 0 & \bar{a}^L & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & \bar{b}^R & 0 \end{array} \right), & \bar{\mathfrak{q}}^R &= \left(\begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ \bar{b}^L & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & \bar{a}^R \\ 0 & 0 & 0 & 0 \end{array} \right), \end{aligned} \quad (3.44)$$

where we made it manifest that $(\mathfrak{q}^{L,R})^\dagger = \bar{\mathfrak{q}}^{L,R}$. The parameters $a^{L,R}, b^{L,R}$ and their conjugates characterise the L- and R-representations, which sit in the diagonal matrix blocks. We can think of these matrices as explicit realisations of the tensors $Q_j^k(p; \{c\})$ introduced in the previous section. The set of parameters $\{c\}$ consists of h and ζ , on which $a^{L,R}, b^{L,R}$ and their conjugates depend.

From the anticommutators, we find immediately the central charges

$$\begin{aligned} \mathfrak{h}^L &= \left(\begin{array}{c|c} |a^L|^2 \mathbb{I} & 0 \\ \hline 0 & |b^R|^2 \mathbb{I} \end{array} \right), & \mathfrak{h}^R &= \left(\begin{array}{c|c} |b^L|^2 \mathbb{I} & 0 \\ \hline 0 & |a^R|^2 \mathbb{I} \end{array} \right), \\ \mathfrak{c} &= \left(\begin{array}{c|c} a^L b^L \mathbb{I} & 0 \\ \hline 0 & a^R b^R \mathbb{I} \end{array} \right), & \bar{\mathfrak{c}} &= \left(\begin{array}{c|c} \bar{a}^L \bar{b}^L \mathbb{I} & 0 \\ \hline 0 & \bar{a}^R \bar{b}^R \mathbb{I} \end{array} \right). \end{aligned} \quad (3.45)$$

It can be explicitly checked that the shortening condition holds. Furthermore, can use the fact that the angular momentum

$$\mathfrak{m} = \mathfrak{h}^L - \mathfrak{h}^R = \text{diag}(+1, +1, -1, -1), \quad (3.46)$$

is quantised and should not receive corrections, together with the explicit form (3.37) of $\mathfrak{c}, \bar{\mathfrak{c}}$ in terms of the momentum to solve for the representation parameters. To do so, let us introduce the *Zhukovski paramterisation* of the momentum in terms of variables x_p^\pm satisfying

$$\frac{x_p^+}{x_p^-} = e^{ip}, \quad \left(x_p^+ + \frac{1}{x_p^+} \right) - \left(x_p^- + \frac{1}{x_p^-} \right) = \frac{2i}{h}. \quad (3.47)$$

We find

$$\begin{aligned} a_p^L &= a_p^R = \sqrt{\zeta} e^{\frac{1}{4}ip} \eta_p, & \bar{a}_p^L &= \bar{a}_p^R = \frac{1}{\sqrt{\zeta}} e^{-\frac{3}{4}ip} \eta_p, \\ b_p^L &= b_p^R = -\sqrt{\zeta} \frac{e^{-\frac{3}{4}ip}}{x_p^-} \eta_p, & \bar{b}_p^L &= \bar{b}_p^R = -\frac{1}{\sqrt{\zeta}} \frac{e^{\frac{1}{4}ip}}{x_p^+} \eta_p, \end{aligned} \quad (3.48)$$

where we explicitly indicated the momentum dependence on the representation parameters and introduced the function

$$\eta_p = e^{ip/4} \sqrt{i \frac{h}{2} (x_p^- - x_p^+)}. \quad (3.49)$$

The definition (3.49) is given in such a way that η_p will have nice analyticity properties that we will use in the next chapter. Note that if we want to identify the vectors in \mathcal{V} with states of the superstring, we should restrict to representations having $\zeta = 1$ to reproduce (3.38).

There is some arbitrariness the choice of the representation coefficients, *e.g.* corresponding to a change of normalisation of the basis vectors in \mathcal{B} . We fix some of this freedom by requiring that we are truly deforming the relativistic representation found perturbatively in the previous chapter. In this way, if we rescale $p \rightarrow p/h$ and expand the coefficients in $1/h$, we find as expected

$$a_p^{L,R} = f_p + O(1/h) = \bar{a}_p^{L,R}, \quad b_p^{L,R} = g_p + O(1/h) = \bar{b}_p^{L,R}. \quad (3.50)$$

Dispersion relation

By explicitly evaluating $\mathfrak{h} = \mathfrak{h}^L + \mathfrak{h}^R$ we find

$$\mathfrak{h} = \frac{h}{2i} \left(x_p^+ - \frac{1}{x_p^+} - x_p^- + \frac{1}{x_p^-} \right) \mathbb{I} = \sqrt{1 + 4h^2 \sin^2 \left(\frac{p}{2} \right)} \mathbb{I}. \quad (3.51)$$

We anticipated the presence of such a non-relativistic dispersion relation in the introduction. In fact, it is an immediate consequence of the shortening condition (3.41) and the non-linear form of \mathfrak{c} and $\bar{\mathfrak{c}}$. Let us rewrite (3.41) as

$$0 = (\mathfrak{h} + \mathfrak{m}) (\mathfrak{h} - \mathfrak{m}) - 4\bar{\mathfrak{c}} \mathfrak{c}, \quad (3.52)$$

Using the fact that the eigenvalues of \mathfrak{m} are ± 1 so that $\mathfrak{m}^2 = 1$, we have

$$\mathfrak{h}^2 = 1 + 4\bar{\mathfrak{c}} \mathfrak{c} = 1 + 4h^2 \sin^2 \left(\frac{\mathbb{P}}{2} \right), \quad (3.53)$$

i.e. precisely the dispersion relation (3.51).

Left-right symmetry

The identity between left- and right-representation coefficients in (3.48) suggests the presence of a discrete symmetry relating them. This is not surprising when we think of the

original coset model, where L and R were introduced as arbitrary labels for the two copies of $\mathfrak{psu}(1, 1|2)$. Furthermore, as discussed in the previous chapter (see also appendix A.3), the supercharges in the symmetry algebra are naturally split in two sets differing only by the relabelling $L \leftrightarrow R$, while the energy and the central charges take have the same form on both representations. As a consequence, up to exchanging raising and lowering operators, we can map the left representation into the right one and *vice versa*.

This results in a discrete \mathbb{Z}_2 -symmetry, which we will call left-right (LR) symmetry. When extended to multiparticle states, this will tell us that *e.g.* any configuration involving only left excitations has an equivalent realisation in terms of right excitations only, *etc.*, which will yield restrictions on the S-matrix elements.

Two-particle representation

The two-particle representation is the tensor product of two one-particle representations like the one we just constructed. Its structure constants will be of the form $Q_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p_1, p_2; \{c_1, c_2\})$ where p_1, p_2 and c_1, c_2 are the momenta and the central charges of the one-particle representations. Even if the two-particle representation will have central charges of the form (3.38), *i.e.* with $\zeta = 1$, this may not necessarily be the case for the two factors. To see that, let us evaluate the structure constant $Q_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p_1, p_2; \{c_1, c_2\})$ for the central charge \mathfrak{c} of (3.38). Since this acts diagonally, we have

$$\mathfrak{c} \cdot A_{\alpha_1}^\dagger(p_1) A_{\alpha_2}^\dagger(p_2) |0\rangle = i \frac{\hbar}{2} \left(e^{i(p_1+p_2)} - 1 \right) \delta_{\alpha_1}^{\beta_1} \delta_{\alpha_2}^{\beta_2} A_{\alpha_1}^\dagger(p_1) A_{\alpha_2}^\dagger(p_2) |0\rangle, \quad (3.54)$$

where we used (3.28) which implies that the worldsheet momentum of this state is $p_1 + p_2$ as we expect. From this, we can read off

$$C_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p_1, p_2; \{c_1, c_2\}) = i \frac{\hbar}{2} \left(e^{i(p_1+p_2)} - 1 \right) \delta_{\alpha_1}^{\beta_1} \delta_{\alpha_2}^{\beta_2}, \quad (3.55)$$

On the other hand, we can commute \mathfrak{c} with one creation operator at the time, finding instead

$$C_{\alpha_1, \alpha_2}^{\beta_1, \beta_2}(p_1, p_2; \{c_1, c_2\}) = i \frac{\hbar}{2} \left(\zeta_1 (e^{ip_1} - 1) + \zeta_2 (e^{ip_2} - 1) \right) \delta_{\alpha_1}^{\beta_1} \delta_{\alpha_2}^{\beta_2}, \quad (3.56)$$

Clearly, (3.55) and (3.56) should match, which cannot happen if $\zeta_1 = \zeta_2 = 1$. We conclude that the two-particle representation is the tensor product of two one-particle representations *with non-trivial central charge values* $\zeta_{1,2}$. It is easy to check that the two expressions we found match if and only if

$$(I): \quad \zeta_1 = 1, \quad \zeta_2 = e^{ip_1} \quad \text{or} \quad (II): \quad \zeta_1 = e^{ip_2}, \quad \zeta_2 = 1. \quad (3.57)$$

As discussed, for consistency with left-right symmetry it natural to take the same choice of $\zeta_{1,2}$ for both the L and R irreducible representations. Therefore, we can label the

whole representation space \mathcal{V} by the choice of the central charge, $\mathcal{V} = \mathcal{V}(p_i, \zeta_i)$. Then the action of the S matrix on the two-particle (reducible) representation is

$$\mathbb{S}: \mathcal{V}(p_1, \zeta_1) \otimes \mathcal{V}(p_2, \zeta_2) \rightarrow \mathcal{V}(p_2, \zeta_2) \otimes \mathcal{V}(p_1, \zeta_1). \quad (3.58)$$

If we assign the value of $\zeta_{1,2}$ on the initial states according to either choice in (3.57), we find that \mathbb{S} exchanges (I) with (II) or *vice versa*. In what follows we will consider the choice (I) for the initial states, which as we will see will reproduce the perturbative results.

We can now rewrite more explicitly (3.34) for a supercharge as

$$\mathbb{Q}_{(12)}(p_1, p_2) = \mathbb{Q}(p_1, \zeta_1 = 1) \otimes \mathbb{I} + \Sigma \otimes \mathbb{Q}(p_2, \zeta_2 = e^{ip_1}), \quad (3.59)$$

where we used that \mathbb{Q} is odd in order to rewrite the action of the graded identity in terms of the fermion sign matrix, which in the basis \mathcal{B} is

$$\Sigma = \text{diag}(+1, -1, +1, -1). \quad (3.60)$$

The 16×16 matrix representation is then found by taking the one-particle charge \mathbb{Q} to be equal to each of the matrices (3.44).

The choice of a non-trivial ζ_1 gives the general form of the two-particle representation

$$\mathbb{Q}_{(12)} = \mathbb{Q}(p_1) \otimes \mathbb{I} + e^{\pm \frac{i}{2} p_1} \Sigma \otimes \mathbb{Q}(p_2), \quad (3.61)$$

where the sign in the exponent depends on which supercharge we consider—positive for \mathbb{Q} and negative for $\overline{\mathbb{Q}}$. This can be compared to the more usual relation

$$\mathbb{Q}_{(12)} = \mathbb{Q}(p_1) \otimes \mathbb{I} + \Sigma \otimes \mathbb{Q}(p_2). \quad (3.62)$$

In algebraic terms, the latter is a trivial (graded) *coproduct*, *i.e.* the most natural way of extending the action of an operator to a tensor product space. We can then say that the symmetries of the superstring have a non-trivial coproduct. The mathematical object to deal with such structures is an *Hopf algebra*⁷. The relevance of Hopf algebras in AdS/CFT is well known [137], see also refs. [138, 139] for a review. In fact, it is possible to use such an algebraic approach to find \mathbb{R} in the case at hand. Here we will follow an equivalent and more direct route, and refer the reader to ref. [S3] for the Hopf-algebraic derivation of the S matrix.

In principle we could go on and construct the representation of the symmetries on higher multiparticle states. This can be done either explicitly [29] or by subsequent applications of the Hopf-algebra coproduct. However, since our focus is on the two-particle S matrix, we will not need to work out the resulting expressions.

⁷More precisely, an Hopf algebra is also equipped with an “antipode” involution, which has the physical interpretation of a particle-to-antiparticle transformation.

3.3 Finding the S matrix

In this section we want to find the S matrix of massive fundamental excitations of the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ superstring. This can be done working with \mathbb{R} or \mathbb{S} , but it will be useful for us to introduce the graded matrix $\check{\mathbb{S}}(p, q)$ satisfying

$$\check{\mathbb{S}}(p, q) = \mathbb{I}^g \mathbb{S}(p, q) = \Pi \mathbb{R}(p, q), \quad (3.63)$$

where Π is the permutation and \mathbb{I}^g is the graded identity. This will make some of our expressions easier to manipulate, because all permutations of tensor product factors will be automatically accounted for, and match with refs. [S3, S5].

Our strategy will be to write down the most general operator

$$\check{\mathbb{S}} : \mathcal{V}(p, 1) \otimes \mathcal{V}(q, e^{ip}) \rightarrow \mathcal{V}(q, e^{ip}) \otimes \mathcal{V}(p, 1), \quad (3.64)$$

as 16×16 matrix, and require that it satisfies suitable physical properties, which we now list.

Off-shell symmetries

The S matrix should commute with the whole off-shell symmetry algebra. Since the latter is generated by the supercharges, it is enough to impose that the S matrix commutes with $\mathbb{Q}^{\text{L,R}}$ and $\bar{\mathbb{q}}^{\text{L,R}}$ in the 16×16 matrix representation given by (3.59).

Explicitly, in terms of $\check{\mathbb{S}}$, we have

$$\check{\mathbb{S}}_{(12)}(p, q) \mathbb{Q}_{(12)}(p, q) = \mathbb{Q}_{(12)}(q, p) \check{\mathbb{S}}_{(12)}(p, q), \quad (3.65)$$

where $\mathbb{Q}_{(12)}(p, q)$ is defined by (3.59). This should be imposed for all four supercharges. Notice that, unlike (3.32), this equations does not feature $\mathbb{Q}_{(21)}$.

Since (3.65) is a linear equation, any solution can be multiplied by a prefactor. In our case, the charges (3.44) have a block-diagonal structure due to the presence of two (L and R) irreducible representations in \mathcal{V} . Therefore, we expect (3.65) to determine $\check{\mathbb{S}}$ at best up to four scalar factors for the LL, LR, RL and RR blocks. We will refer to the part of \mathbb{S} determined independently from these factors (*i.e.*, suitable ratios of the matrix elements) as its “matrix part”.

Left-right symmetry

An additional constraint is the discrete left-right symmetry, which amounts to imposing that scattering processes differing only by relabelling $\text{L} \leftrightarrow \text{R}$ should be indistinguishable. In particular, this removes part of the ambiguity due to scalar factors, relating the LL block to the RR one, and the LR one to the RL one.

Braiding and physical unitarity

In section 3.1 we have established that the \mathbb{R} matrix must satisfy braiding unitarity as well as be unitary as a matrix. In terms of \check{S} this gives

$$\check{S}_{(12)}(q, p) \check{S}_{(12)}(p, q) = \mathbb{I}, \quad (\check{S}_{(12)}(p, q))^\dagger \check{S}_{(12)}(p, q) = \mathbb{I}. \quad (3.66)$$

One of these two equations quadratic constrains can be eliminated in terms of the linear one

$$\check{S}_{(12)}(q, p) = (\check{S}_{(12)}(p, q))^\dagger. \quad (3.67)$$

While this linear equation will put a restriction on the reality properties of the scalar factors, it is easy to see that the matrix part should automatically satisfy it. In fact, taking the conjugate of (3.65), we find

$$\overline{\mathbb{Q}}_{(12)}(p, q) (\check{S}_{(12)}(p, q))^\dagger = (\check{S}_{(12)}(p, q))^\dagger \overline{\mathbb{Q}}_{(12)}(q, p). \quad (3.68)$$

Since for any charge $\mathbb{Q}_{(12)}$ its conjugate $\overline{\mathbb{Q}}_{(12)}$ is also part of the algebra, we have that $\check{S}_{(12)}(p, q)^\dagger$ satisfies the same invariance condition that define $\check{S}_{(12)}(q, p)$. Therefore the part of each of the two S matrices that is completely determined by symmetries must coincide.

The reflectionless $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ S matrix

It turns out that imposing all the symmetries together with the unitarity requirements gives two physically distinct solutions for $\check{S}_{pq} = \check{S}_{(12)}(p, q)$. These coincide on the LL and RR sectors, but are different on the mixed sectors. One solution gives

$$(T): \quad \langle \mathcal{X}^L \mathcal{Y}^R | \check{S}_{pq} | \mathcal{X}^L \mathcal{Y}^R \rangle = 0 \quad \text{and} \quad \langle \mathcal{X}^R \mathcal{Y}^L | \check{S}_{pq} | \mathcal{X}^L \mathcal{Y}^R \rangle \neq 0, \quad (3.69)$$

and similarly for $L \leftrightarrow R$, whereas the other gives

$$(R): \quad \langle \mathcal{X}^L \mathcal{Y}^R | \check{S}_{pq} | \mathcal{X}^L \mathcal{Y}^R \rangle \neq 0 \quad \text{and} \quad \langle \mathcal{X}^R \mathcal{Y}^L | \check{S}_{pq} | \mathcal{X}^L \mathcal{Y}^R \rangle = 0, \quad (3.70)$$

where \mathcal{X}, \mathcal{Y} are two generic excitations. Keeping into account that the matrix \check{S} permutes the final states, the case we indicated by (T) corresponds to *pure transmission* of the target-space chirality, while (R) corresponds to *pure reflection*. From the symmetry properties, there is no reason to choose one over the other. However, by a perturbative calculation it is easy to check which are the non-vanishing matrix elements. In fact, a tree-level calculation suffices [100], and shows that the case (R) cannot reproduce the worldsheet superstring S matrix. From now on, we will restrict our considerations to the case (T) of a *pure-transmission* (or *reflectionless*) S matrix.

$$\left(\begin{array}{cccccccc|cccccccc}
A_{pq}^{LL} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & C_{pq}^{LL} & 0 & 0 & D_{pq}^{LL} & 0 & 0 & 0 & A_{pq}^{RL} & 0 & 0 & 0 & 0 & F_{pq}^{RL} & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & D_{pq}^{RL} & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & B_{pq}^{LL} & 0 & 0 & E_{pq}^{LL} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & F_{pq}^{LL} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & C_{pq}^{RL} & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & B_{pq}^{RL} & 0 & 0 & 0 & 0 & E_{pq}^{RL} & 0 & 0 \\
\hline
0 & 0 & A_{pq}^{LR} & 0 & 0 & 0 & 0 & F_{pq}^{LR} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & D_{pq}^{LR} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{pq}^{RR} & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & C_{pq}^{RR} & 0 & 0 & D_{pq}^{RR} & 0 & 0 \\
0 & 0 & 0 & C_{pq}^{LR} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & B_{pq}^{LR} & 0 & 0 & 0 & 0 & E_{pq}^{LR} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & B_{pq}^{RR} & 0 & 0 & E_{pq}^{RR} & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & F_{pq}^{RR}
\end{array} \right)$$

Figure 3.3: The matrix representation of $\check{\mathbb{S}}_{(12)}(p, q)$ in the basis constructed out of the tensor product of \mathcal{B} . Notice how each block is given by a 4×4 S matrix suitably embedded in a tensor product structure.

Constructing the tensor-product basis out of \mathcal{B} , we can explicitly represent $\check{\mathbb{S}}_{pq}$ as a 16×16 matrix which naturally splits in four blocks depending on the target-space chiralities of the particle scattered

$$\check{\mathbb{S}}_{pq} = \left(\begin{array}{c|c} \check{\mathbb{S}}_{pq}^{LL} & \check{\mathbb{S}}_{pq}^{RL} \\ \check{\mathbb{S}}_{pq}^{LR} & \check{\mathbb{S}}_{pq}^{RR} \end{array} \right). \quad (3.71)$$

The full form of the S matrix is given in figure 3.3. Let us further investigate its structure. Due to LR symmetry, it is natural to distinguish the scattering of particles of the same or opposite chirality.

Same-chirality scattering

Let us consider particles of the same, let us say LL, chirality. Then the non-vanishing scattering processes are

$$\begin{aligned}
\check{\mathbb{S}} |\phi_p^L \phi_q^L\rangle &= A_{pq}^{LL} |\phi_q^L \phi_p^L\rangle, & \check{\mathbb{S}} |\phi_p^L \psi_q^L\rangle &= B_{pq}^{LL} |\psi_q^L \phi_p^L\rangle + C_{pq}^{LL} |\phi_q^L \psi_p^L\rangle, \\
\check{\mathbb{S}} |\psi_p^L \psi_q^L\rangle &= F_{pq}^{LL} |\psi_q^L \psi_p^L\rangle, & \check{\mathbb{S}} |\psi_p^L \phi_q^L\rangle &= D_{pq}^{LL} |\phi_q^L \psi_p^L\rangle + E_{pq}^{LL} |\psi_q^L \phi_p^L\rangle.
\end{aligned} \quad (3.72)$$

These are determined up to a single, overall scalar factor $\mathcal{S}_{pq}^{\text{LL}}$, and they read

$$\begin{aligned}
A_{pq}^{\text{LL}} &= \mathcal{S}_{pq}^{\text{LL}} e^{\frac{i}{2}(p-q)} \frac{x_p^- - x_q^+}{x_p^+ - x_q^-}, & B_{pq}^{\text{LL}} &= \mathcal{S}_{pq}^{\text{LL}} e^{-\frac{i}{2}q} \frac{x_p^+ - x_q^+}{x_p^+ - x_q^-}, \\
C_{pq}^{\text{LL}} &= \mathcal{S}_{pq}^{\text{LL}} e^{\frac{i}{4}(p-3q)} \frac{-\frac{2i}{h} \eta_p \eta_q}{x_p^+ - x_q^-}, & D_{pq}^{\text{LL}} &= \mathcal{S}_{pq}^{\text{LL}} e^{\frac{i}{2}p} \frac{x_p^- - x_q^-}{x_p^+ - x_q^-}, \\
E_{pq}^{\text{LL}} &= \mathcal{S}_{pq}^{\text{LL}} e^{-\frac{i}{4}(p+q)} \frac{-\frac{2i}{h} \eta_p \eta_q}{x_p^+ - x_q^-}, & F_{pq}^{\text{LL}} &= -\mathcal{S}_{pq}^{\text{LL}}.
\end{aligned} \tag{3.73}$$

In the RR sector we find exactly the same formulae, in terms of a scalar factor $\mathcal{S}_{pq}^{\text{RR}}$. Requiring LR-symmetry implies, for instance,

$$\langle \phi_q^{\text{L}} \phi_p^{\text{L}} | \check{\mathbb{S}} | \phi_p^{\text{L}} \phi_q^{\text{L}} \rangle = \langle \phi_q^{\text{R}} \phi_p^{\text{R}} | \check{\mathbb{S}} | \phi_p^{\text{R}} \phi_q^{\text{R}} \rangle, \tag{3.74}$$

so that it must be

$$\mathcal{S}_{pq}^{\text{LL}} = \mathcal{S}_{pq}^{\text{RR}} = \mathcal{S}_{pq}. \tag{3.75}$$

Opposite-chirality scattering

If we now consider processes of LR chirality we find

$$\begin{aligned}
\check{\mathbb{S}} | \phi_p^{\text{L}} \phi_q^{\text{R}} \rangle &= A_{pq}^{\text{LR}} | \phi_q^{\text{R}} \phi_p^{\text{L}} \rangle + B_{pq}^{\text{LR}} | \psi_q^{\text{R}} \psi_p^{\text{L}} \rangle, & \check{\mathbb{S}} | \phi_p^{\text{L}} \psi_q^{\text{R}} \rangle &= C_{pq}^{\text{LR}} | \psi_q^{\text{R}} \phi_p^{\text{L}} \rangle, \\
\check{\mathbb{S}} | \psi_p^{\text{L}} \psi_q^{\text{R}} \rangle &= E_{pq}^{\text{LR}} | \psi_q^{\text{R}} \psi_p^{\text{L}} \rangle + F_{pq}^{\text{LR}} | \phi_q^{\text{R}} \phi_p^{\text{L}} \rangle, & \check{\mathbb{S}} | \psi_p^{\text{L}} \phi_q^{\text{R}} \rangle &= D_{pq}^{\text{LR}} | \phi_q^{\text{R}} \psi_p^{\text{L}} \rangle,
\end{aligned} \tag{3.76}$$

while for RL

$$\begin{aligned}
\check{\mathbb{S}} | \phi_p^{\text{R}} \phi_q^{\text{L}} \rangle &= A_{pq}^{\text{RL}} | \phi_q^{\text{L}} \phi_p^{\text{R}} \rangle + B_{pq}^{\text{RL}} | \psi_q^{\text{L}} \psi_p^{\text{R}} \rangle, & \check{\mathbb{S}} | \phi_p^{\text{R}} \psi_q^{\text{L}} \rangle &= C_{pq}^{\text{RL}} | \psi_q^{\text{L}} \phi_p^{\text{R}} \rangle, \\
\check{\mathbb{S}} | \psi_p^{\text{R}} \psi_q^{\text{L}} \rangle &= E_{pq}^{\text{RL}} | \psi_q^{\text{L}} \psi_p^{\text{R}} \rangle + F_{pq}^{\text{RL}} | \phi_q^{\text{L}} \phi_p^{\text{R}} \rangle, & \check{\mathbb{S}} | \psi_p^{\text{R}} \phi_q^{\text{L}} \rangle &= D_{pq}^{\text{RL}} | \phi_q^{\text{L}} \psi_p^{\text{R}} \rangle.
\end{aligned} \tag{3.77}$$

The elements read

$$\begin{aligned}
A_{pq}^{\text{LR}} &= \mathcal{S}_{pq}^{\text{LR}} e^{\frac{i}{2}(p+q)} \frac{1 - x_p^+ x_q^-}{1 - x_p^+ x_q^+}, & B_{pq}^{\text{LR}} &= \mathcal{S}_{pq}^{\text{LR}} e^{\frac{i}{4}(3p-q)} \frac{\frac{2i}{h} \eta_p \eta_q}{1 - x_p^+ x_q^+}, \\
C_{pq}^{\text{LR}} &= \mathcal{S}_{pq}^{\text{LR}} e^{\frac{i}{2}(2p+q)} \frac{1 - x_p^- x_q^-}{1 - x_p^+ x_q^+}, & D_{pq}^{\text{LR}} &= \mathcal{S}_{pq}^{\text{LR}} e^{\frac{i}{2}p}, \\
E_{pq}^{\text{LR}} &= -\mathcal{S}_{pq}^{\text{LR}} e^{ip} \frac{1 - x_p^- x_q^+}{1 - x_p^+ x_q^+}, & F_{pq}^{\text{LR}} &= \mathcal{S}_{pq}^{\text{LR}} e^{\frac{i}{4}(p-q)} \frac{-\frac{2i}{h} \eta_p \eta_q}{1 - x_p^+ x_q^+},
\end{aligned} \tag{3.78}$$

and

$$\begin{aligned}
A_{pq}^{\text{RL}} &= \mathcal{S}_{pq}^{\text{RL}} e^{-\frac{i}{2}(p+q)} \frac{1 - x_p^+ x_q^-}{1 - x_p^- x_q^-}, & B_{pq}^{\text{RL}} &= \mathcal{S}_{pq}^{\text{RL}} e^{-\frac{i}{4}(p+3q)} \frac{\frac{2i}{h} \eta_p \eta_q}{1 - x_p^- x_q^-}, \\
C_{pq}^{\text{RL}} &= \mathcal{S}_{pq}^{\text{RL}} e^{-\frac{i}{2}q}, & D_{pq}^{\text{RL}} &= \mathcal{S}_{pq}^{\text{RL}} e^{-\frac{i}{2}(p+2q)} \frac{1 - x_p^+ x_q^+}{1 - x_p^- x_q^-}, \\
E_{pq}^{\text{RL}} &= -\mathcal{S}_{pq}^{\text{RL}} e^{-iq} \frac{1 - x_p^- x_q^+}{1 - x_p^- x_q^-}, & F_{pq}^{\text{RL}} &= \mathcal{S}_{pq}^{\text{RL}} e^{-\frac{i}{4}(3p+5q)} \frac{-\frac{2i}{h} \eta_p \eta_q}{1 - x_p^- x_q^-}.
\end{aligned} \tag{3.79}$$

Now LR-symmetry implies *e.g.*

$$\langle \psi_q^R \phi_p^L | \check{\mathbb{S}} | \phi_p^L \psi_q^R \rangle = \langle \psi_q^L \phi_p^R | \check{\mathbb{S}} | \phi_p^R \psi_q^L \rangle . \quad (3.80)$$

This can be solved in terms of a single scalar factor $\widetilde{\mathcal{F}}_{pq}$

$$\begin{aligned} \mathcal{S}_{pq}^{\text{LR}} &= \widetilde{\mathcal{F}}_{pq} e^{-\frac{i}{2}(p+q)} \left(\frac{1 - x_p^+ x_q^+}{1 - x_p^- x_q^-} \right)^{+1/2} , \\ \mathcal{S}_{pq}^{\text{RL}} &= \widetilde{\mathcal{F}}_{pq} e^{+\frac{i}{2}(p+q)} \left(\frac{1 - x_p^+ x_q^+}{1 - x_p^- x_q^-} \right)^{-1/2} . \end{aligned} \quad (3.81)$$

By this choice, we find not only $\mathbf{C}_{pq}^{\text{RL}} = \mathbf{C}_{pq}^{\text{LR}}$, but also $\mathbf{A}_{pq}^{\text{RL}} = \mathbf{A}_{pq}^{\text{LR}}$, $\mathbf{B}_{pq}^{\text{RL}} = \mathbf{B}_{pq}^{\text{LR}}$, and so on.

Scalar factors

The requirements of braiding unitarity and physical unitarity pose constraints on the scalar factors \mathcal{S}_{pq} and $\widetilde{\mathcal{F}}_{pq}$. In particular, they must satisfy

$$\begin{aligned} \mathcal{S}(p, q) \mathcal{S}(q, p) &= 1, & |\mathcal{S}(p, q)|^2 &= 1, \\ \widetilde{\mathcal{F}}(p, q) \widetilde{\mathcal{F}}(q, p) &= 1, & |\widetilde{\mathcal{F}}(p, q)|^2 &= 1. \end{aligned} \quad (3.82)$$

i.e. they are given by two antisymmetric phases.

There is an additional condition that we can require on the S matrix, that is *crossing symmetry*. This is an extension of the well-known relativistic covariance under particle-antiparticle transformation to our non-relativistic S matrix. Crossing invariance will put stringent requirements on the analytic structure of the dressing factors, which we will analyse at length in the next chapter.

The full S matrix

The full S matrix that we are eventually interested in is not the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ -invariant one. Instead, it is invariant under two copies of such an algebra, with the excitations transforming in a tensor product representation (2.103–2.105). One way to find it is to construct a representation of the full $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$, similar to (3.59), in terms of 64×64 matrices. The S matrix one would find then by repeating the procedure of this section is precisely the tensor product of two copies of the 16×16 matrix we just found,⁸

$$\check{\mathbb{S}}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} = \check{\mathbb{S}}_{\mathfrak{su}(1|1)_{\text{c.e.}}^2} \otimes \check{\mathbb{S}}_{\mathfrak{su}(1|1)_{\text{c.e.}}^2} . \quad (3.83)$$

⁸The tensor product yields a 256×256 matrix with many vanishing elements, rather than 64×64 one. This is because in (2.103) we do not consider any state of the form *e.g.* $\phi^L \otimes \phi^R$, that do exist in the tensor product space.

The tensor product should take into account the signs arising from permuting the fermions. In components this gives

$$\begin{aligned} \left(\check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4}\right)_{\alpha_1, \alpha_2}^{\alpha_3, \alpha_4} &= \left(\check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4}\right)_{\kappa_1 \iota_1, \kappa_2 \iota_2}^{\kappa_3 \iota_3, \kappa_4 \iota_4} = \\ &(-1)^{\varepsilon(\iota_1)\varepsilon(\kappa_2)+\varepsilon(\iota_3)\varepsilon(\kappa_4)} \left(\check{S}_{\mathfrak{su}(1|1)_{\text{c.e.}}^2}\right)_{\kappa_1, \kappa_2}^{\kappa_3, \kappa_4} \left(\check{S}_{\mathfrak{su}(1|1)_{\text{c.e.}}^2}\right)_{\iota_1, \iota_2}^{\iota_3, \iota_4}, \end{aligned} \quad (3.84)$$

where we used the notation introduced in (2.104), with $\alpha = \kappa\iota$. From this, all of the scattering elements can be readily calculated, and again we have that the scattering can be decomposed based on the LR chirality, with the matrix elements being the “square” of what found in the previous section. To illustrate this, let us work out a few processes in the LL sector

$$\begin{aligned} \check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{++q}^L\rangle &= A_{pq}^{LL} A_{pq}^{LL} |\Phi_{++q}^L \Phi_{++p}^L\rangle, \\ \check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{--q}^L\rangle &= B_{pq}^{LL} B_{pq}^{LL} |\Phi_{--q}^L \Phi_{++p}^L\rangle + C_{pq}^{LL} C_{pq}^{LL} |\Phi_{++q}^L \Phi_{--p}^L\rangle \\ &\quad + B_{pq}^{LL} C_{pq}^{LL} (|\Phi_{-+q}^L \Phi_{+-p}^L\rangle - |\Phi_{+-q}^L \Phi_{-+p}^L\rangle), \end{aligned} \quad (3.85)$$

where we used the short-hand notation $|\Phi_{\pm\pm p}^L\rangle = |\Phi_{\pm\pm}^L(p)\rangle$ and so on. Similarly, in the LR sector we have for instance

$$\begin{aligned} \check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{++q}^R\rangle &= A_{pq}^{LR} A_{pq}^{LR} |\Phi_{++q}^R \Phi_{++p}^L\rangle - B_{pq}^{LR} B_{pq}^{LR} |\Phi_{--q}^R \Phi_{--p}^L\rangle \\ &\quad + A_{pq}^{LR} B_{pq}^{LR} (|\Phi_{-+q}^R \Phi_{+-p}^L\rangle + |\Phi_{+-q}^R \Phi_{-+p}^L\rangle), \\ \check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{--q}^R\rangle &= C_{pq}^{LR} C_{pq}^{LR} |\Phi_{--q}^R \Phi_{++p}^L\rangle. \end{aligned} \quad (3.86)$$

In particular, in $\check{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4}$ there are still only *two* undetermined scalar factors, *i.e.* \mathcal{S}^2 and $\check{\mathcal{S}}^2$.

In what follows, unless confusion may arise, we will drop the subscript invariant algebra from \check{S} .

Yang-Baxter equation and integrability

Once we have successfully derived the $2 \rightarrow 2$ S matrix by maxing use of the off-shell symmetry, the question remains as to whether such an S matrix can be used to define an arbitrary $M \rightarrow M$ scattering. The condition for this is expressed by Yang-Baxter equation (3.26), which in terms of $\check{S}_{pq} = \check{S}(p, q)$ reads

$$\check{S}_{qr} \otimes \mathbb{I} \cdot \mathbb{I} \otimes \check{S}_{pr} \cdot \check{S}_{pq} \otimes \mathbb{I} = \mathbb{I} \otimes \check{S}_{pq} \cdot \check{S}_{pr} \otimes \mathbb{I} \cdot \mathbb{I} \otimes \check{S}_{qr}, \quad (3.87)$$

where we made the three-particle tensor product explicit.

It is easy to check that the Yang-Baxter equation is non-trivially⁹ satisfied by the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ S matrix that we found, as well as by the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ one—a fact following from the tensor-product structure. Therefore, they can be both used to define an integrable two-dimensional QFT.

⁹ In the case of $\check{S}_{\mathfrak{su}(1|1)_{\text{c.e.}}^2}$, once the YB equation is spelled out in components, one finds 112 equations involving the non-zero matrix elements, which vanish by using their explicit form.

Comparison with the $\mathfrak{su}(2|2)$ -symmetric S matrix

The symmetry algebra $\mathfrak{su}(1|1)_L \oplus \mathfrak{su}(1|1)_R$ centrally extended can be embedded into $\mathfrak{su}(2|2)_{\text{c.e.}}$. This last algebra is also related to an integrable S matrix. The S matrix of fundamental string excitations in $\text{AdS}_5 \times S^5$ is in fact invariant under two copies of $\mathfrak{su}(2|2)_{\text{c.e.}}$, with the excitations transforming in a tensor product representation akin to the one we encountered here, see *e.g.* [29].

Since $\mathfrak{su}(1|1)_{\text{c.e.}}^2 \subset \mathfrak{su}(2|2)_{\text{c.e.}}$ we could expect that the $\mathfrak{su}(2|2)_{\text{c.e.}}$ -invariant S matrix should arise of a special limit of our $\check{\mathbb{S}}$. Differently from $\mathfrak{su}(1|1)^2$, in $\mathfrak{su}(2|2)_{\text{c.e.}}$ there exists an additional $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ symmetry. The excitations $\phi^{L,R}$ then would transform as a doublet under the former $\mathfrak{su}(2)$, while $\psi^{L,R}$ would be a doublet under the latter. This would require the scattering to take the form

$$\begin{aligned} \check{\mathbb{S}}_{pq} (|\phi_p^L \phi_q^R\rangle + |\phi_p^R \phi_q^L\rangle) &= \# (|\phi_p^L \phi_q^R\rangle + |\phi_p^R \phi_q^L\rangle), \\ \check{\mathbb{S}}_{pq} (|\phi_p^L \phi_q^R\rangle - |\phi_p^R \phi_q^L\rangle) &= \# (|\phi_p^L \phi_q^R\rangle - |\phi_p^R \phi_q^L\rangle) + \# (|\psi_p^L \psi_q^R\rangle - |\psi_p^R \psi_q^L\rangle). \end{aligned} \quad (3.88)$$

We immediately see that we cannot obtain this general form from the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ S matrix that we have constructed. The reason is that we have imposed on the S matrix the discrete LR symmetry, which is incompatible with the additional $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ invariance. It is this symmetry that distinguishes the massive sector of $\text{AdS}_3 \times S^3 \times T^4$ from a truncation of the $\text{AdS}_5 \times S^5$ superstring at all loops. Notice also that in the latter case the excitations form a single irreducible representation of the symmetry algebra, and hence a single scalar factor is left undetermined.

3.4 Chapter summary

The main result of this chapter is the derivation of the two-particle S matrix $\check{\mathbb{S}}$ (which is related to \mathbb{S} by some fermion signs) for the scattering of fundamental massive excitations of the $\text{AdS}_3 \times S^3 \times T^4$ superstring, and of the non-relativistic dispersion relation (3.53). The matrix $\check{\mathbb{S}}$ satisfies the Yang-Baxter equation as well as braiding unitarity and physical unitarity, so that we can use it to unambiguously define any multiparticle scattering. $\check{\mathbb{S}}$ is defined up to two antisymmetric unit-norm functions $\mathcal{S}_{pq}, \widetilde{\mathcal{F}}_{pq}$ —the *dressing factors*.

It may appear strange that we could determine $\check{\mathbb{S}}$ for the massive excitations while completely ignoring the massless ones. Even if we expect those to transform in a different irreducible representation of the symmetry algebra, it is easy to see that perturbatively there exist quartic massive-massless interactions in the Lagrangian—the two sectors *are* coupled. Therefore, the effective vertices for massive particles will contain massless excitations running in the loops, even if they cannot appear in the final states. We can draw a parallel of sorts with $SU(N)$ Yang-Mills theory, with the de Wit-Fadeev-Popov ghosts playing the role of the massless modes in the massive sector. Even if we ignore




Figure 3.4: The bootstrap condition for the bound-state S matrix. We consider two particles ϕ with complex momenta $p_1 = \frac{1}{2}P + iv$ and $p_2 = \frac{1}{2}P - iv$ that form a bound state Φ of momentum P (thicker violet line). The scattering of a third particle $\phi(q)$ with $\Phi(P)$ can be resolved in terms of the scattering with its constituents. Note that the point in the diagram at which the two particles fuse to give a bound state should not be interpreted as a scattering event.

the ghosts, we can still correctly predict that the S-matrix elements will be $SU(N)$ invariant. However, to compute their value we need to take the ghosts into account—not everything is fixed by the $SU(N)$ symmetry. Here $SU(N)$ is replaced by a much larger symmetry, involving the off-shell algebra as well as the higher conserved charges, and the only underdetermined elements are the dressing factors. They will contain the dynamical information about the theory, featuring poles and cuts in such a way as to also account for processes involving virtual massless particles. We will return to this topic in the next chapter.

We should also stress that we only computed the S matrix of *fundamental* massive excitations. If the theory admits bound states, the S matrix that scatter those should also be determined to have a complete handle on the spectrum, in particular to compute the wrapping effects [74, 76]. Typically, this can be done again by means of integrability. The scattering of a particle with a bound state can be defined as illustrated in figure 3.4, *i.e.* in terms of the scattering with its constituents—a procedure called *bootstrap*. This is particularly simple when one takes advantage of the Hopf algebra structure, see *e.g.* refs. [138–140]. In the case of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, such a calculation has not been performed so far.

4 Crossing symmetry and dressing factors

In this chapter we will see that crossing symmetry puts stringent requirements on the analytic structure of the dressing factors.

In a relativistic theory, crossing symmetry is a consequence of the fact that fields are constructed out of particle *and* antiparticle creation and annihilation operators. One can then show that any two elements of the S matrix related by a particle-to-antiparticle transformation are equivalent up to performing an analytic continuation, which exchanges the branches of the dispersion relation. This invariance is illustrated pictorially in figure 4.1. There we have introduced the relativistic rapidity ϑ satisfying

$$E = m \cosh \vartheta, \quad p = m \sinh \vartheta, \quad (4.1)$$

which parametrises the positive branch of the dispersion relation $E^2 = m^2 + p^2$. In a theory where all particles coincide with their antiparticles (*e.g.* photons), each of one can be mapped into each of the other by shifting

$$\vartheta \rightarrow \vartheta + i\pi. \quad (4.2)$$

This flips both the sign of E and p . If particles and antiparticles do not coincide, the transformation (4.2) should be supplemented by a linear map sending the particle-representation into the antiparticle one—*e.g.* in the case of quarks and antiquarks, this would be a map between the fundamental and anti-fundamental representation of $\mathfrak{su}(3)$.

These concept can be extended to the non-relativistic S matrix that we computed in the previous chapter. In order to do so, it is first convenient to introduce a *rapidity* variable z akin to ϑ . Then, we will study the charge-conjugation properties of the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ or equivalently $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ algebras that we constructed in the previous chapter. This will lead us to formulate the *crossing equations* constraining the dressing factors, for which we will find a solution in section 4.1.

4.1 Crossing symmetry

Let us start by defining the rapidity z and its domain.




Figure 4.1: In a relativistic theory, crossing invariance can be understood by looking at a scattering process in two different ways. In the left panel, time flows upwards, and the scattering of particles 1 and 2 happens with rapidity $\vartheta = \vartheta_2 - \vartheta_1$. In the right panel, time flows from left to right, and the same scattering involves the antiparticle of 2 (moving backwards in time) and 1. The scattering happens with rapidity $\vartheta^c = \vartheta + i\pi$, which can be understood as a Lorentzian angle.

Uniformising the dispersion relation

The all-loop dispersion relation for massive excitations on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ reads

$$E^2(p) = 1 + 4h^2 \sin^2 \frac{p}{2}. \quad (4.3)$$

This is the same functional form appearing in $\text{AdS}_5 \times \text{S}^5$. To uniformise the dispersion relation we can introduce a rapidity variable following ref. [97]. Let z satisfy

$$p(z) = 2 \operatorname{am} z, \quad \sin \frac{p(z)}{2} = \operatorname{sn}(z, \kappa), \quad E(z) = \operatorname{dn}(z, \kappa), \quad (4.4)$$

in terms of Jacobi's elliptic functions, where the elliptic modulus is $\kappa = -4h^2$. This defines a torus with a real period $2\omega_1$ and an imaginary period $2\omega_2$ that depend on h through

$$\omega_1 = 2\operatorname{K}(\kappa), \quad \omega_2 = 2i\operatorname{K}(1 - \kappa) - 2\operatorname{K}(\kappa), \quad (4.5)$$

where K is the complete elliptic integral of the first kind. In this parameterisation the real z -axis corresponds to real momentum and positive energy. In order to describe bound states, it is useful to consider complex momenta, and therefore complex rapidities z in the torus. One can check that all the definitions we have given are also ω_1 -periodic, so that we can always restrict to $|\operatorname{Re}(z)| \leq \omega_1/2$, corresponding to $-\pi < p \leq \pi$.

The Zhukovski variables $x^\pm(z)$ are meromorphic functions on the torus

$$x^\pm(z) = \frac{1}{2h} \left(\frac{\operatorname{cn}(z, \kappa)}{\operatorname{sn}(z, \kappa)} \pm i \right) (1 + \operatorname{dn}(z, \kappa)), \quad (4.6)$$

and satisfy the relations (3.47). Furthermore we can resolve the square root in η_p by

$$\eta(z) = \frac{\operatorname{dn} \frac{z}{2} (\operatorname{cn} \frac{z}{2} + i \operatorname{sn} \frac{z}{2} \operatorname{dn} \frac{z}{2})}{1 + 4h^2 \operatorname{sn}^4 \frac{z}{2}}. \quad (4.7)$$




Figure 4.2: The rapidity torus with several significant curves. The solid blue line is the real z -axis (physical region), the dashed blue line is the $z = \omega_2$ axis (“crossed” region). In the leftmost figure the torus is divided in four regions by $|x^\pm| = 1$ and in the central figure it is divided by $\text{Im}(x^\pm) = 0$. The rightmost picture depicts both sets of curves, which intersect in eight points with real part $\pm\omega_1/4$, denoted by stars.

The S-matrix elements that we computed in the previous chapter can now be expressed purely in terms of z . Remarkably, they are all *rational functions* on two copies¹ of the elliptic torus, up to the dressing factors.

In figure 4.2 we depict the rapidity torus, highlighting several significant curves. The red ones correspond to $|x^\pm(z)| = 1$. In analogy with $\text{AdS}_5 \times \text{S}^5$, we will call the region they delimit which contains the real z -axis *the physical region*—this is where the complex momenta of physical bound state are expected to be found [74]. It can be identified by requiring $|x^\pm(z)| > 1$. In what follows, we will consider all S-matrix elements corresponding to physical processes to be evaluated at rapidities lying in the physical region.

A shift $z \rightarrow z + \omega_2$ flips the sign of energy and momentum (4.4). The region containing the line $\text{Im}(z) = \omega_2$ corresponds to real crossed variables, having real momentum and real negative energy. In terms of the Zukhovski variables we have that

$$x^+(z \pm \omega_2) = \frac{1}{x^+(z)}, \quad x^-(z \pm \omega_2) = \frac{1}{x^-(z)}, \quad \eta(z \pm \omega_2) = \frac{\pm i}{x^+(z)} \eta(z), \quad (4.8)$$

so that the shift maps the physical region $|x^\pm| > 1$ to $|x^\pm| < 1$. The scattering of

¹Recall that $\mathbb{S}(p_1, p_2)$ or more precisely $\mathbb{S}(z_1, z_2)$ depends on z_1 and z_2 separately, in contrast with the relativistic case where it would be $\mathbb{S}(\vartheta_1, \vartheta_2) = \mathbb{S}(\vartheta_1 - \vartheta_2)$.

antiparticles, which take the negative branch of the dispersion relation, will give S-matrix elements evaluated for z in the crossed region, *i.e.* for $|x^\pm(z)| < 1$.

Charge conjugation

Let us go back to the representations of $\mathfrak{psu}(1|1)_{c.e.}^4$, or equivalently of $\mathfrak{su}(1|1)_{c.e.}^2$. Even better, since we know how to obtain multiparticle representations in terms of a non-trivial coproduct, we can even restrict to the one-particle representation given by the 4×4 supermatrices (3.44) and by the relative central charges. We can label this pair of representations as

$$\mathcal{V}_{\text{LR}}(e, c, \bar{c}) = \mathcal{V}_{\text{L}}(e, c, \bar{c}) \oplus \mathcal{V}_{\text{R}}(e, c, \bar{c}), \quad (4.9)$$

where e labels the energy and c, \bar{c} the central charges eigenvalues.

By transforming all of the charges and supercharges \mathfrak{x} by means of a supertransposition

$$\mathfrak{x} \rightarrow -\mathfrak{x}^{\text{st}}, \quad (4.10)$$

we find a different pair of representations of $\mathfrak{su}(1|1)_{c.e.}^2$. Notice that the minus sign is needed to ensure that the anticommutation relations are preserved. This clearly flips the sign of the central charges. In particular, we now have that $E < 0$ and that left and right excitation have swapped their charge under \mathfrak{m} .

Combining $\mathfrak{x} \rightarrow -\mathfrak{x}^{\text{st}}$ with the crossing transformation $z \rightarrow z + \omega_2$, we can *almost* obtain a pair of representations with the same central charges as the one that we started. If that were the case, that would mean that there exists a (pseudo)unitary transformation relating $-\mathfrak{x}^{\text{st}}(z + \omega_2)$ with $\mathfrak{x}(z)$ —that is, a charge conjugation matrix. There are two obstructions for this to be the case. Firstly, for the eigenvalues of \mathfrak{m} to be swapped, charge conjugation must exchange the left and right irreducible representations. Secondly, the crossing transformation does flip the sign of energy, but acts on $C(z)$ as

$$c(z) = i\zeta \frac{\hbar}{2} \left(e^{ip(z)} - 1 \right) \rightarrow c(z + \omega_2) = -e^{-ip(z)} c(z), \quad (4.11)$$

and similarly for $\bar{c}(z)$. These additional phase factors can be accounted for if we additionally transform the supercharges $\mathfrak{q} = \mathfrak{q}^{\text{L,R}}$ by a phase

$$\begin{aligned} (-\mathfrak{q}^{\text{st}}(z + \omega_2)) &\rightarrow e^{+\frac{i}{2}P} (-\mathfrak{q}^{\text{st}}(z + \omega_2)), \\ (-\bar{\mathfrak{q}}^{\text{st}}(z + \omega_2)) &\rightarrow e^{-\frac{i}{2}P} (-\bar{\mathfrak{q}}^{\text{st}}(z + \omega_2)). \end{aligned} \quad (4.12)$$

This transformation is an $U(1)$ automorphism of $\mathfrak{su}(1|1)_{c.e.}^2$. Using it, we can finally conclude that, for the supercharges it must be

$$\begin{aligned} -e^{+\frac{i}{2}P} \mathfrak{q}^{\text{st}}(z + \omega_2) &= \mathcal{C} \mathfrak{q}(z) \mathcal{C}^{-1}, \\ -e^{-\frac{i}{2}P} \bar{\mathfrak{q}}^{\text{st}}(z + \omega_2) &= \mathcal{C} \bar{\mathfrak{q}}(z) \mathcal{C}^{-1}, \end{aligned} \quad (4.13)$$

where \mathcal{C} is the charge-conjugation matrix, which exchanges the left and right representations. It can be written in the basis \mathcal{B} as

$$\mathcal{C} = \left(\begin{array}{cc|cc} 0 & 0 & \xi^{\text{RL}} & 0 \\ 0 & 0 & 0 & -i\xi^{\text{RL}} \\ \hline \xi^{\text{LR}} & 0 & 0 & 0 \\ 0 & -i\xi^{\text{LR}} & 0 & 0 \end{array} \right). \quad (4.14)$$

The two normalisation constants ξ^{LR} and ξ^{RL} , one per representation, can be set without loss of generality to $e.g.$

$$\xi^{\text{LR}} = \xi^{\text{RL}} = 1. \quad (4.15)$$

In this way $\mathcal{C}^2 = \Sigma$ and $\mathcal{C}^{-1} = \mathcal{C}^\dagger$.

By using the fact supertransposition acts on the supercharges as $q^{\text{st}} = q^\dagger \Sigma$, we can rewrite eq. (4.13) as

$$\begin{aligned} q^\dagger(z \pm \omega_2) &= \mp e^{-\frac{i}{2}p} \Sigma \mathcal{C}^\dagger q(z) \mathcal{C}, \\ \bar{q}^\dagger(z \pm \omega_2) &= \mp e^{+\frac{i}{2}p} \Sigma \mathcal{C}^\dagger \bar{q}(z) \mathcal{C}, \end{aligned} \quad (4.16)$$

where we also used the fact that $\Sigma \mathcal{C} = \mathcal{C} \Sigma$ and $\Sigma q = -q \Sigma$. Note how these equations are related by Hermitian conjugation.

Crossing equations

The fact that the charges enjoy the invariance (4.13) together with the fundamental invariance property of the S matrix

$$\mathbb{R}_{(12)} \mathbb{Q}_{(12)} = \mathbb{Q}_{(21)} \mathbb{R}_{(12)}, \quad (4.17)$$

allows us to derive an additional property of \mathbb{R}_{12} .² To this end, let us focus on the case where \mathbb{Q} is one of the supercharges $q^{\text{L,R}}$ or of their conjugates—the invariance under the central charges follows trivially. Furthermore, let us recall that $\mathbb{Q}_{(12)}$ is given by the coproduct (3.59), and similarly $\mathbb{Q}_{(21)}$ is given by

$$\mathbb{Q}_{(21)}(p, q) = \mathbb{Q}(p, \zeta_1 = e^{iq}) \otimes \Sigma + \mathbb{I} \otimes \mathbb{Q}(q, \zeta_2 = 1). \quad (4.18)$$

It is then easy to write down the invariance of \mathbb{R} under $q = q^{\text{L,R}}$

$$\begin{aligned} \mathbb{R}_{(12)}(z_1, z_2) &\left(q(z_1) \otimes \mathbb{I} + e^{\frac{i}{2}p_1} \Sigma \otimes q(z_2) \right) \\ &= \left(e^{\frac{i}{2}p_2} q(z_1) \otimes \Sigma + \mathbb{I} \otimes q(z_2) \right) \mathbb{R}_{(12)}(z_1, z_2), \end{aligned} \quad (4.19)$$

²We choose to work with \mathbb{R} here rather than \mathbb{S} or $\check{\mathbb{S}}$ to obtain a more compact and familiar-looking final expression the crossing equations.

where $p_j = p(z_j)$. If we take the transpose of this equation with respect to *e.g.* the first space, we find

$$\begin{aligned} & (-\Sigma \mathfrak{q}^\dagger(z_1) \otimes \mathbb{I} + \Sigma \otimes \mathfrak{q}(z_2)) \mathbb{R}_{(12)}^{t_1}(z_1, z_2) \\ &= \mathbb{R}_{(12)}^{t_1}(z_1, z_2) \left(-e^{\frac{i}{2}p_2} \Sigma \mathfrak{q}^\dagger(z_1) \otimes \Sigma + e^{\frac{i}{2}p_1} \mathbb{I} \otimes \mathfrak{q}(z_2) \right). \end{aligned} \quad (4.20)$$

where we also multiplied by $-\Sigma$ from the left. This equation holds for any z_1, z_2 . If we now perform a shift $z_1 \rightarrow z_1 + \omega_2$ and use (4.16), we can recast the equation in the form

$$\begin{aligned} & \left(\mathfrak{q}(z_1) \otimes \mathbb{I} + e^{\frac{i}{2}p_1} \Sigma \otimes \mathfrak{q}(z_2) \right) \mathcal{C}_{(1)} \mathbb{R}_{(12)}^{t_1}(z_1 + \omega_2, z_2) \mathcal{C}_{(1)}^\dagger \\ &= \mathcal{C}_{(1)} \mathbb{R}_{(12)}^{t_1}(z_1 + \omega_2, z_2) \mathcal{C}_{(1)}^\dagger \left(e^{\frac{i}{2}p_2} \mathfrak{q}(z_1) \otimes \Sigma + \mathbb{I} \otimes \mathfrak{q}(z_2) \right), \end{aligned} \quad (4.21)$$

where $\mathcal{C}_{(1)} = \mathcal{C} \otimes \mathbb{I}$. Comparing this expression with (4.19), we see that the combination $\mathcal{C}_{(1)} \mathbb{R}_{(12)}^{t_1}(z_1 + \omega_2, z_2) \mathcal{C}_{(1)}^\dagger$ has the same invariance property with respect to \mathfrak{q}^\dagger as the inverse of $\mathbb{R}_{(12)}(z_1, z_2)$. In fact, it is not hard to see that the same calculation applies to all of the supercharges.³ We can therefore require that it is

$$\mathcal{C}_{(1)} \mathbb{R}_{(12)}^{t_1}(z_1 + \omega_2, z_2) \mathcal{C}_{(1)}^\dagger = \left(\mathbb{R}_{(12)}(z_1, z_2) \right)^{-1}. \quad (4.22)$$

This is the *crossing equation*. If we had repeated a similar calculation taking the transpose in the second space, we would have found

$$\mathcal{C}_{(2)} \mathbb{R}_{(12)}^{t_2}(z_1, z_2 - \omega_2) \mathcal{C}_{(2)}^\dagger = \left(\mathbb{R}_{(12)}(z_1, z_2) \right)^{-1}. \quad (4.23)$$

In fact, the two crossing equations are related by braiding unitarity, as we will see explicitly in the next subsection.

What we have just found with regard to crossing symmetry is somewhat similar to what happened for the unitarity condition (3.68). The invariance property we have found is seemingly new, but it can be found from manipulating eq. (4.19). Since the matrix part of \mathbb{R} has been found imposing (4.19), that will *automatically* satisfy crossing invariance. However, this requirement will mean that the scalar factors \mathcal{S}_{pq} and $\widetilde{\mathcal{S}}_{pq}$ may not be arbitrary, but satisfy the analyticity requirements coming from (4.22) or equivalently (4.23).

Here we have derived the crossing symmetry requirements for the $\mathfrak{su}(1|1)_{c.e.}^2$ S matrix. The ones for the bigger $\mathfrak{psu}(1|1)_{c.e.}^4$ S matrix follow immediately by requiring that it is given by the tensor product (3.83) of two *crossing invariant* $\mathfrak{su}(1|1)_{c.e.}^2$ S matrices. Equivalently, it is easy to repeat the construction of the charge-conjugation matrix for the charges in $\mathfrak{psu}(1|1)_{c.e.}^4$ and obtain the crossing equations in that way.

³Note that in the case of $\mathfrak{q}^{L,R}$ the sign in the phase shifts $e^{\pm \frac{i}{2}p}$ is everywhere opposite to the one we used for $\mathfrak{q}^{L,R}$.

Constrains on the scalar factors

Let us write the requirements imposed by crossing symmetry on the scalar factors of the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ S matrix. For later convenience, let us normalise the scalar factors by introducing two functions $\sigma(p_1, p_2)$ and $\tilde{\sigma}(p_1, p_2)$ satisfying

$$\begin{aligned}\mathcal{S}^2(p_1, p_2) &= \sigma^{-2}(p_1, p_2) \frac{x_1^+ - x_2^-}{x_1^- - x_2^+} \frac{1 - \frac{1}{x_1^- x_2^+}}{1 - \frac{1}{x_1^+ x_2^-}}, \\ \tilde{\mathcal{S}}^2(p_1, p_2) &= \tilde{\sigma}^{-2}(p_1, p_2) \frac{1 - \frac{1}{x_1^- x_2^+}}{1 - \frac{1}{x_1^+ x_2^-}},\end{aligned}\tag{4.24}$$

In this way the phase for the same-chirality diagonal processes features the Beisert-Dippel-Staudacher [141] matrix element. The scalar factors have the form

$$\sigma(p_1, p_2) = e^{i\theta(p_1, p_2)}, \quad \tilde{\sigma}(p_1, p_2) = e^{i\tilde{\theta}(p_1, p_2)},\tag{4.25}$$

where $\theta(p_1, p_2)$ and $\tilde{\theta}(p_1, p_2)$ are antisymmetric real analytic functions for real p_1, p_2 —a requirement that follows from unitarity.

The crossing equations that we have found then imply

$$\begin{aligned}\sigma(p_1^c, p_2)^2 \tilde{\sigma}(p_1, p_2)^2 &= g(p_1, p_2), & \sigma(p_1, p_2)^2 \tilde{\sigma}(p_1^c, p_2)^2 &= \tilde{g}(p_1, p_2), \\ \sigma(p_1, p_2^c)^2 \tilde{\sigma}(p_1, p_2)^2 &= \frac{1}{\tilde{g}(p_2^c, p_1)}, & \sigma(p_1, p_2)^2 \tilde{\sigma}(p_1, p_2^c)^2 &= \frac{1}{g(p_2^c, p_1)},\end{aligned}\tag{4.26}$$

where

$$\begin{aligned}g(p_1, p_2) &= \left(\frac{x_2^-}{x_2^+}\right)^2 \frac{\left(1 - \frac{1}{x_1^+ x_2^+}\right) \left(1 - \frac{1}{x_1^- x_2^-}\right)}{\left(1 - \frac{1}{x_1^+ x_2^-}\right)^2} \frac{x_1^- - x_2^+}{x_1^+ - x_2^-}, \\ \tilde{g}(p_1, p_2) &= \left(\frac{x_2^-}{x_2^+}\right)^2 \frac{(x_1^- - x_2^+)^2}{(x_1^+ - x_2^+) (x_1^- - x_2^-)} \frac{1 - \frac{1}{x_1^- x_2^+}}{1 - \frac{1}{x_1^+ x_2^-}}.\end{aligned}\tag{4.27}$$

and the superscript c indicates crossing. More precisely, we have

$$p_1^c = p(z_1 + \omega_2), \quad p_2^c = p(z_2 - \omega_2).\tag{4.28}$$

It is then easy to check that the four equations (4.26) are related by the antisymmetry of the scalar factors, *i.e.* owing to unitarity. Therefore, it will be enough to restrict ourselves to *e.g.*

$$\sigma(p_1^c, p_2)^2 \tilde{\sigma}(p_1, p_2)^2 = g(p_1, p_2), \quad \sigma(p_1, p_2)^2 \tilde{\sigma}(p_1^c, p_2)^2 = \tilde{g}(p_1, p_2),\tag{4.29}$$

which are the ones due to (4.22).




Figure 4.3: The paths used for analytic continuation from z to $z + \omega_2$ (in purple) are vertical segments that lie close to the boundary of $|\text{Re}(\gamma)| < \omega_1/4$. They cross the red lines $|x^\pm| = 1$ when $\text{Im}(x^\pm) < 0$.

We have remarked how the matrix part of \mathbb{R} is a meromorphic function on the rapidity torus. However, by iterating the crossing transformation twice we find that the dressing factors are not $2\omega_2$ -periodic:

$$\begin{aligned} \frac{\sigma(z_1 + 2\omega_2, z_2)^2}{\sigma(z_1, z_2)^2} &= \frac{g(z_1 + \omega_2, z_2)}{\tilde{g}(z_1, z_2)} = \left(\frac{x_1^+ - x_2^+}{x_1^+ - x_2^-} \frac{x_1^- - x_2^-}{x_1^- - x_2^+} \right)^2, \\ \frac{\tilde{\sigma}(z_1 + 2\omega_2, z_2)^2}{\tilde{\sigma}(z_1, z_2)^2} &= \frac{\tilde{g}(z_1 + \omega_2, z_2)}{g(z_1, z_2)} = \left(\frac{1 - \frac{1}{x_1^+ x_2^-}}{1 - \frac{1}{x_1^- x_2^+}} \frac{1 - \frac{1}{x_1^- x_2^-}}{1 - \frac{1}{x_1^+ x_2^+}} \right)^2. \end{aligned} \quad (4.30)$$

Therefore, they must have cuts on the rapidity torus, so that the whole crossing-invariant S matrix is defined on some more complicated surface. In fact, even if we considered shifts by $4\omega_2, 6\omega_2, \text{etc.}$ we would still find no periodicity, meaning that the dressing phases should live on an infinite cover of the z -torus.

Even in the better-understood case of $\text{AdS}_5 \times \text{S}^5$ it is unknown how to define a generalised rapidity that resolves the additional cuts and describes that cover. Therefore, in what follows we will continue using z (or x^\pm), keeping in mind that when describing paths on the torus additional care should be used in the case where a cut is crossed.

4.2 Solving the crossing equations

A solution for the crossing equations is given by two antisymmetric phases satisfying eq. (4.29). It is convenient to rewrite this in terms of crossing equations for the sum and the difference of the two phases $\theta(p_1, p_2)$ and $\tilde{\theta}(p_1, p_2)$. Let us denote the product and the ratio of the dressing factors by

$$\sigma^+(p_1, p_2) = \sigma(p_1, p_2) \tilde{\sigma}(p_1, p_2), \quad \sigma^-(p_1, p_2) = \frac{\sigma(p_1, p_2)}{\tilde{\sigma}(p_1, p_2)}, \quad (4.31)$$

and corresponding phases by $\theta^+(p_1, p_2)$ and $\theta^-(p_1, p_2)$. By analogy with the case of $\text{AdS}_5 \times \text{S}^5$ [25], it is also useful to rewrite each phase as

$$\theta(p_1, p_2) = \chi(x_1^+, x_2^+) + \chi(x_1^-, x_2^-) - \chi(x_1^+, x_2^-) - \chi(x_1^-, x_2^+), \quad (4.32)$$

where χ is an antisymmetric function. Similar expressions can be introduced for $\tilde{\theta}(p_1, p_2)$, $\theta^+(p_1, p_2)$ and $\theta^-(p_1, p_2)$.

Dressing phases in $\text{AdS}_5 \times \text{S}^5$

Due to the many similarities of our S matrix with the $\text{AdS}_5 \times \text{S}^5$, it is worth briefly describing the solution of the crossing equation in that case. In the $\text{AdS}_5 \times \text{S}^5$ S matrix, a single dressing phase appears. An all-loop solution to its crossing equation [97] was found by Beisert, Eden and Staudacher (BES) [28]

$$\begin{aligned} \sigma^{\text{BES}}(z_1, z_2) \sigma^{\text{BES}}(z_1 + \omega_2, z_2) &= h(x_1^\pm, x_2^\pm), \\ h(x_1^\pm, x_2^\pm) &= \frac{x_2^-}{x_2^+} \frac{x_1^- - x_2^+}{x_1^+ - x_2^-} \frac{1 - \frac{1}{x_1^+ x_2^+}}{1 - \frac{1}{x_1^+ x_2^-}}. \end{aligned} \quad (4.33)$$

A particularly useful representation of this phase was given by Dorey, Hofman and Maldacena (DHM) [142]

$$\chi^{\text{BES}}(x, y) = i \oint \frac{dw}{2\pi i} \oint \frac{d\tilde{w}}{2\pi i} \frac{1}{x-w} \frac{1}{y-\tilde{w}} \log \frac{\Gamma[1 + \frac{i}{2}h(w + 1/w - \tilde{w} - 1/\tilde{w})]}{\Gamma[1 - \frac{i}{2}h(w + 1/w - \tilde{w} - 1/\tilde{w})]}. \quad (4.34)$$

This is valid in the physical region $|x| < 1$, $|y| < 1$. On the boundary of this region, the integral representations has cuts so that a path should be chosen to analytically continue the expression to the crossed region.

As it is discussed in detail in [75], for the $\text{AdS}_5 \times \text{S}^5$ crossing equations to be solved we can chose the path depicted in figure 4.3. These are curves $\gamma(z)$ that go from z to $z + \omega_2$ with constant $\text{Re}(\gamma)$, and which lie close to the boundaries of the region $|\text{Re}(\gamma)| < \omega_1/4$ and crossing the lines $|x^\pm| = 1$ in the region $\text{Im}(x^\pm) < 0$. Then, the precise statement of the crossing equation (4.33) is that the BES dressing factor evaluated for z in the physical

region as defined by the DHM double integral, times itself analytically continued through the cuts of (4.34) along $\gamma(z)$, equals the rational function on the right hand side.⁴

Our crossing equations (4.29) will be interpreted in a similar sense. In fact, we will from now consider the crossing transformation as a continuation along the paths $\gamma(z)$ described in figure 4.3. This will be understood every time we write p^c , z^c and so on.

At leading order $O(h^{-1})$ in a large- h expansion,⁵ the BES phase coincides with the Arutyunov-Frolov-Staudacher (AFS) phase [25]. This was found based on rather general considerations on the large- h behaviour of the energy of string states, and for this reason it is expected to be common to several AdS/CFT duals. It can be written as

$$\sigma^{\text{AFS}}(x_1, x_2) = \left(\frac{1 - \frac{1}{x_1^+ x_2^+}}{1 - \frac{1}{x_1^+ x_2^-}} \right) \left(\frac{1 - \frac{1}{x_1^+ x_2^-} 1 - \frac{1}{x_1^- x_2^+}}{1 - \frac{1}{x_1^+ x_2^+} 1 - \frac{1}{x_1^- x_2^-}} \right)^{\frac{i}{2} h(x_1+1/x_1-x_2-1/x_2)}, \quad (4.35)$$

The next-to-leading-order term in strong-coupling expansion is the Hernández-López (HL) phase [144],

$$\chi^{\text{HL}}(x, y) = \frac{\pi}{2} \oint \frac{dw}{2\pi i} \oint \frac{d\tilde{w}}{2\pi i} \frac{1}{x-w} \frac{1}{y-\tilde{w}} \text{sign}(\tilde{w} + 1/\tilde{w} - w - 1/w). \quad (4.36)$$

The HL phase appears at order $O(1)$, and indeed its expression is h -independent. For later convenience, let us perform one of the two integrals in (4.36) and obtain the representation

$$\chi^{\text{HL}}(x, y) = \left(\oint - \oint \right) \frac{dw}{4\pi} \frac{1}{x-w} (\log(y-w) - \log(y-1/w)), \quad (4.37)$$

where the two integrals are performed in the upper and lower unit semi-circle respectively, counterclockwise in both cases. The HL phase solves the “odd” part of the AdS₅ crossing equation [65]

$$\sigma^{\text{HL}}(z_1, z_2) \sigma^{\text{HL}}(z_1 + \omega_2, z_2) = \sqrt{\frac{h_{12}}{h_{\bar{1}2}}} = \sqrt{h_{12} (h_{12})^*}, \quad (4.38)$$

where

$$h_{12} (h_{12})^* = \frac{\ell^{\text{HL}}(x_1^+, x_2^-) \ell^{\text{HL}}(x_1^-, x_2^+)}{\ell^{\text{HL}}(x_1^+, x_2^+) \ell^{\text{HL}}(x_1^-, x_2^-)}, \quad \ell^{\text{HL}}(x, y) = \frac{x-y}{1-xy}, \quad (4.39)$$

where complex conjugation amounts to sending $x_k^\pm \rightarrow x_k^\mp$.⁶

⁴Actually, we are restricting to a subset of all the allowed paths used in the case of AdS₅ × S⁵, see section 4 in [75]. This more special choice will be the one suitable for our crossing equations.

⁵To find the asymptotic expansion of the BES phase at strong coupling one can expand the integrand using that $i \log \frac{\Gamma(1+ix)}{\Gamma(1-ix)} = -x \log \frac{x^2}{e^2} - \frac{\pi}{2} \text{sign}(\text{Re } x) - 2 \sum_{n=0}^{\infty} \frac{\zeta(-2n-1)}{2n+1} \frac{(-1)^n}{x^{2n+1}}$ for $\text{Re } x \neq 0$. This expression corrects some typos in the expansion given in [143].

⁶One can check that in order for (4.37) to solve (4.38) it is necessary to choose the path of analytic continuation as in figure 4.3. To do this one can mimic the arguments presented in appendix A.4.

Solution for the sum of the phases

Taking the product of the two crossing equations (4.29), we find an equation for σ^+

$$\sigma^+(z_1, z_2)^2 \sigma^+(z_1 + \omega_2, z_2)^2 = g_{12} \tilde{g}_{12}. \quad (4.40)$$

We observe that the r.h.s of this equation can be written in terms of the function h_{12} appearing on the r.h.s of the AdS₅ crossing equation (4.33)

$$g_{12} \tilde{g}_{12} = \frac{(h_{12})^3}{(h_{12})^*}, \quad (4.41)$$

where we used the constrain (3.47). The above relation allows us to solve the crossing equation (4.40) using parts of the AdS₅ dressing phase

$$\sigma_{12}^+ = \frac{(\sigma_{12}^{\text{BES}})^2}{\sigma_{12}^{\text{HL}}}, \quad i.e. \quad \theta_{12}^+ = 2\theta_{12}^{\text{BES}} - \theta_{12}^{\text{HL}}. \quad (4.42)$$

To show that such a σ_{12}^+ satisfies equation (4.40) one need only use equations (4.33) and (4.38). It is convenient to express σ_{12}^+ in terms of a DHM-like double-integral representation, by defining $\chi^+(x, y)$ as

$$\begin{aligned} \chi^+(x, y) &= 2\chi^{\text{BES}}(x, y) - \chi^{\text{HL}}(x, y) \\ &= \oint \frac{dw}{2\pi i} \oint \frac{d\tilde{w}}{2\pi i} \frac{1}{x-w} \frac{1}{y-\tilde{w}} \left(2i \log \frac{\Gamma[1 + ih(w + 1/w - \tilde{w} - 1/\tilde{w})]}{\Gamma[1 - ih(w + 1/w - \tilde{w} - 1/\tilde{w})]} \right. \\ &\quad \left. - \frac{\pi}{2} \text{sign}(\tilde{w} + 1/\tilde{w} - w - 1/w) \right), \end{aligned} \quad (4.43)$$

in the physical region. Notice that the above expression is exact to all orders in the coupling h . We postpone the perturbative expansion of this and the following expressions to chapter 7, where we will also compare them with independent results.

Solution for the difference of the phases

Taking the ratio of the two crossing equations (4.29), we get

$$\frac{\sigma^-(z_1, z_2)^2}{\sigma^-(z_1 + \omega_2, z_2)^2} = \frac{\tilde{g}_{12}}{g_{12}}, \quad (4.44)$$

where

$$\frac{\tilde{g}_{12}}{g_{12}} = \frac{\ell^-(x_1^+, x_2^-) \ell^-(x_1^-, x_2^+)}{\ell^-(x_1^+, x_2^+) \ell^-(x_1^-, x_2^-)}, \quad \ell^-(x, y) \equiv (x - y) \left(1 - \frac{1}{xy} \right). \quad (4.45)$$

Notice that this equation involves the ratio rather than the product of the dressing factor with its analytic continuation. As we show in appendix A.4, defining $\chi^-(x, y)$ in the

physical region ($|x|, |y| > 1$) by the integral

$$\begin{aligned} \chi^-(x, y) &= \oint \frac{dw}{8\pi} \frac{1}{x-w} \log \left[(y-w) \left(1 - \frac{1}{yw} \right) \right] \text{sign}((w-1/w)/i) \quad -x \leftrightarrow y \\ &= \left(\oint - \oint \right) \frac{dw}{8\pi} \frac{1}{x-w} \log \left[(y-w) \left(1 - \frac{1}{yw} \right) \right] \quad -x \leftrightarrow y, \end{aligned} \quad (4.46)$$

solves the crossing equation (4.44). By construction, χ^- is antisymmetric. Note that the integrand of χ^- does not depend explicitly on the coupling h , in contrast to the solution of the crossing equation (4.40) which is solved by an integrand with an infinite series expansion in h . This is because equation (4.44) is “odd” in the sense of [65].

The all-loop expressions for χ and $\tilde{\chi}$ are then given by

$$\begin{aligned} \chi(x, y) &= \chi^{\text{BES}}(x, y) + \frac{1}{2} (-\chi^{\text{HL}}(x, y) + \chi^-(x, y)) , \\ \tilde{\chi}(x, y) &= \chi^{\text{BES}}(x, y) + \frac{1}{2} (-\chi^{\text{HL}}(x, y) - \chi^-(x, y)) . \end{aligned} \quad (4.47)$$

These solutions are expressed in terms of the non-perturbative BES phase plus terms at the HL order. These latter contributions to χ and $\tilde{\chi}$ are independent of h . As such, they can be added to the DHM representation of the BES phase without affecting the h -resummation.

4.3 Poles of the S matrix

Now that we have found a solution to the crossing equations, it is natural to ask whether this is *the* solution appearing in the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ S matrix. It is clear that our phases could be multiplied by any “CDD factor” [145], that is, any solution of the homogeneous crossing equations

$$\sigma_{pq}^{\text{CDD}} \tilde{\sigma}_{p^c q}^{\text{CDD}} = 1, \quad \sigma_{p^c q}^{\text{CDD}} \tilde{\sigma}_{pq}^{\text{CDD}} = 1. \quad (4.48)$$

Such solutions exist. The simplest ones are meromorphic functions on the torus, which can be defined in terms of

$$\chi_{pq}^{\text{CDD}} = \frac{i}{2} \log \frac{(x-y)^{n_1}}{(1-xy)^{n_2}}, \quad \tilde{\chi}_{pq}^{\text{CDD}} = \frac{i}{2} \log \frac{(x-y)^{n_2}}{(1-xy)^{n_1}}, \quad (4.49)$$

with n_1, n_2 integer constants. Such factors will modify the pole structure of the S matrix. However, there is a close connection between simple poles in the physical region of the S matrix and the bound states of the model. Therefore, by exploring the expected bound-state spectrum of the model we will be able to put stringent restriction on the CDD factors.

Bound states and short representations

Excitations transform in representations of $\mathfrak{psu}(1|1)_{c.e.}^4$. Furthermore, the representations satisfy a shortening condition which immediately follows from the one of $\mathfrak{su}(1|1)_{c.e.}^2$, see eq. (3.41)

$$\mathbb{H}^2 = \mathbb{M}^2 + \overline{\mathbb{C}\mathbb{C}}. \quad (4.50)$$

Bound states preserving some supersymmetry also transform in a short representation of the symmetry algebra. Let us consider a two-particle state containing two left-moving particles, *e.g.* $|\Phi_{++}^L \Phi_{++}^L\rangle$. For generic values of the momenta p and q the tensor product of two fundamental left representations is an irreducible long representation. However, at special points the tensor product becomes reducible. In particular, we find that the shortening condition (4.50) is satisfied for

$$x_p^+ = x_q^- \quad \text{or} \quad x_p^- = x_q^+. \quad (4.51)$$

Only at these points it is possible to construct short sub-representations. Therefore any pole in the S-matrix corresponding to a supersymmetric bound state will have to satisfy one of these conditions.

An interesting feature of the $\mathfrak{psu}(1|1)_{c.e.}^4$ algebra is that all short irreducible representations are two-dimensional while all long irreducible representations have dimension four. A two-particle bound state will therefore transform in a representation which has the same form as the fundamental representation, differing only in the values of the central charges. This should be contrasted with the centrally extended $\mathfrak{psu}(2|2)$ algebra appearing in $\text{AdS}_5 \times S^5$ [18, 26, 57], where the fundamental representation has dimension four while the M -particle bound state has dimension $4M$ [146].

At the points where the tensor product becomes reducible some of the elements of the S matrix become zero or develop poles. These singularities will appear both in the dressing factors, as we will investigate later, and in the ratio of S matrix elements. In fact, with an appropriate normalisation we will have

$$\mathbb{R}_{pq} \mathcal{V}_0 = 0, \quad \mathcal{V}_0 \subset \mathcal{V}_L(p) \otimes \mathcal{V}_L(q), \quad (4.52)$$

when p and q satisfy (4.51), while \mathbb{R}_{pq} is regular (finite) on the complement of \mathcal{V}_0 . The bound state representation is then the factor representation on the quotient space $\mathcal{V}_L(p) \otimes \mathcal{V}_L(q) / \mathcal{V}_0$ [147]. Using the explicit form of the S matrix, for the point $x_p^+ = x_q^-$ we find that the state $|\Phi_{++}^L \Phi_{++}^L\rangle$ belongs to the short representation, and we will therefore refer to it as a $\mathfrak{su}(2)$ bound state. In the case $x_p^- = x_q^+$ the short representation includes the state $|\Phi_{--}^L \Phi_{--}^L\rangle$, and is a potential $\mathfrak{sl}(2)$ bound state.⁷

⁷In $\text{AdS}_5 \times S^5$ the physical bound states correspond to “ $\mathfrak{su}(2)$ bound states”. The “ $\mathfrak{sl}(2)$ bound states” appear as bound states of the mirror theory that we alluded to in the introduction [74].

To decide which bound state belongs to the physical spectrum we need to impose additional constraints on the momenta of the fundamental excitations. In the region $s_1 \ll s_2$ the wave function of a scattering state takes the general form⁸

$$\Psi(s_1, s_2) \approx e^{i(ps_1+qs_2)} + S(p, q)e^{i(ps_2+qs_1)}, \quad (4.53)$$

where the first term describes the incoming wave and the second term the outgoing wave, or equivalently

$$\Psi(s_1, s_2) \approx \frac{1}{S(p, q)} e^{i(ps_1+qs_2)} + e^{i(ps_2+qs_1)}. \quad (4.54)$$

To find a bound state we analytically continue the wave function to complex values of the momenta

$$p = \frac{\tilde{p}}{2} + iv, \quad q = \frac{\tilde{p}}{2} - iv. \quad (4.55)$$

The wave function then behaves as

$$\Psi(s_1, s_2) \approx \frac{1}{S(p, q)} e^{v(s_2-s_1)} + e^{-v(s_2-s_1)}, \quad s_1 \ll s_2. \quad (4.56)$$

If there are bound states in the spectrum, we expect $S(p, q)$ to have a pole, as it can be understood in a diagrammatic expansion in terms of a propagator that goes on the mass shell. Additionally, for the bound-state wave function to be normalizable, the outgoing-wave exponential should be decaying. Hence we are interested in the solution where the momentum of the first particle has a positive imaginary part, $v > 0$. By imposing the condition (3.47) for x_p^\pm and x_q^\pm in the physical region, we find that for $x_p^+ = x_q^-$ the momentum p has a positive imaginary part, while $x_p^- = x_q^+$ leads to the imaginary part being negative. We hence conclude that only the $\mathfrak{su}(2)$ bound state can appear in the physical spectrum. Of course this will have to be confirmed by the presence of suitable poles in the S matrix, which will pose a constraint on the dressing factors

So far we have only considered bound states in the LL-sector. If we start with two right-moving excitations we again find an $\mathfrak{su}(2)$ bound state at $x_p^+ = x_q^-$, simply by left-right symmetry. It is interesting to consider a state consisting of one left- and one right-moving excitations such as $|\Phi_{++}^L \Phi_{--}^R\rangle$. In this case the shortening condition (4.50) is satisfied for $x_p^+ = 1/x_q^+$ and $x_p^- = 1/x_q^-$. Neither of these solutions lie in the physical region $|x_p^\pm| > 1$, $|x_q^\pm| > 1$ and hence *there are no supersymmetric bound states in the LR sector*.

In summary we find that, based on the shortening condition and the matrix form of \mathbb{R}_{pq} , physical two-particle $\mathfrak{su}(2)$ bound states exist in the LL and RR sectors. The LR sector, on the other hand, does not contain any bound states.

⁸In order to avoid confusion with the dressing phase we denote the world-sheet coordinate by s .

Semiclassical bound states from giant magnons

Owing to its integrability, the $\text{AdS}_5 \times \text{S}^5$ NLSM admits classical off-shell soliton solutions called *giant magnons* [20]. These can be thought of as a coherent superposition of several many one-particle excitations (magnons in the dual spin chain). The simplest giant magnon is a classical string solution living in a $\mathbb{R} \times \text{S}^2$ subspace of $\text{AdS}_5 \times \text{S}^5$, and having a definite momentum p . It can be extended to a solution in $\mathbb{R} \times \text{S}^3$, the *dyonic* giant magnon [148], which carries angular momentum M along the additional angle. This solution corresponds to the semiclassical limit of a bound state of $|M|$ fundamental magnons [149].

Since both the fundamental giant magnon and the dyonic extension live in $\mathbb{R} \times \text{S}^3$ they can be directly embedded in $\text{AdS}_3 \times \text{S}^3$ [150]. How does our discussion on the allowed bound states fit together with the giant magnon picture?

An important difference between our case and $\text{AdS}_5 \times \text{S}^5$ is that in the latter space a dyonic giant magnon with positive \mathbb{M} -charge $+M$ can be *continuously* rotated to the corresponding magnon with negative charge $-M$, due to the presence of an additional $\mathfrak{su}(2)$ symmetry. However, in the case of $\text{AdS}_3 \times \text{S}^3$ such a rotation is not possible since the intermediate states would not sit inside S^3 . Therefore, while in $\text{AdS}_5 \times \text{S}^5$ there is no notion of left and right giant magnon, in our case the two states with charges $+M$ and $-M$ are independent and can be distinguished by the sign of their eigenvalue under \mathbb{M} —their target-space chirality. Only configurations of the same chirality can be used to build a dyonic giant magnon, so that only the corresponding microscopic excitations will have bound states. Since the notion of left and right excitations was defined precisely in terms of their charge under \mathbb{M} , we must expect to have LL and RR $\mathfrak{su}(2)$ -bound states resembling the ones of $\text{AdS}_5 \times \text{S}^5$, but no LR or RL bound states. This is precisely the result of our representation-theoretical analysis of the previous subsection.

Simple poles of the S matrix

Scattering processes involving formation or exchange of bound states give rise to single poles in the S matrix for physical values of the spectral parameters [151]. Let us consider the s -channel diagram in figure 4.4a. The process involves two fundamental particles from the same sector, *e.g.* two left-movers, in the physical region $|x_i| > 1$, $i = p, q$, which form an on-shell boundstate and then split up again. Similarly to the case of the $\mathfrak{su}(2)$ sector in AdS_5 [142, 149], this should lead to a pole in the corresponding S-matrix element at $x_p^+ = x_q^-$. The relevant element is, up to inessential e^{ip} prefactors which we will always drop here,

$$\mathcal{A}_{pq} = \langle \Phi_{++}^L \Phi_{++}^L | \check{\mathbb{S}}_{pq} | \Phi_{++}^L \Phi_{++}^L \rangle = \frac{x_p^- - x_q^+}{x_p^+ - x_q^-} \frac{1 - \frac{1}{x_p^- x_q^+}}{1 - \frac{1}{x_p^+ x_q^-}} \sigma_{pq}^{-2}. \quad (4.57)$$




Figure 4.4: On the left two particles in the same sector form an $\mathfrak{su}(2)$ bound state in the s channel. Applying the crossing transformation to $\Phi_{++}^L(p)$ yields the t -channel diagram on the right, where one particle has unphysical momentum p^c (red dashed lines).

As discussed in appendix A.4 the dressing factor is regular at $x_p^+ = x_q^-$, so that \mathcal{A}_{pq} has a simple pole there.

This s channel process is related through crossing symmetry to the exchange of a bound state in the t channel, depicted in figure 4.4b. There the particle of momentum p has been crossed so that $x_{p^c}^\pm = 1/x_p^\pm$ are not in the physical region. Since the two processes are related by crossing symmetry, the poles in the s channel automatically fix the singularities in the t channel. In fact crossing symmetry implies [S3]

$$\mathcal{A}_{pq} \tilde{\mathcal{A}}_{p^c q} = 1, \quad \text{where } \tilde{\mathcal{A}}_{pq} = \langle \Phi_{++}^L \Phi_{++}^R | \check{\mathbb{S}}_{pq} | \Phi_{++}^R \Phi_{++}^L \rangle, \quad (4.58)$$

so that a pole of \mathcal{A}_{pq} corresponds to a pole of $\tilde{\mathcal{A}}_{p^c q}^{-1}$. We can check this explicitly by considering

$$\tilde{\mathcal{A}}_{p^c q} = \frac{1 - \frac{1}{x_p^- x_q^+}}{1 - \frac{1}{x_p^c x_q^-}} \frac{1 - \frac{1}{x_p^+ x_q^-}}{1 - \frac{1}{x_p^+ x_q^+}} \tilde{\sigma}_{p^c q}^{-2}. \quad (4.59)$$

Since $\tilde{\sigma}$ is regular when continued inside the unit circle (see appendix A.4), $\tilde{\mathcal{A}}_{p^c q}$ has a zero at $x_{p^c}^+ = 1/x_q^-$, as expected.

If we consider S-matrix elements involving one left- and one right-moving particle we expect no poles, since there are no corresponding bound states. Therefore a process such as the one depicted in figure 4.5a should not happen. Indeed, the S-matrix element

$$\tilde{\mathcal{B}}_{pq} = \langle \Phi_{++}^L \Phi_{--}^R | \check{\mathbb{S}}_{pq} | \Phi_{--}^R \Phi_{++}^L \rangle = \frac{1 - \frac{1}{x_p^- x_q^+}}{1 - \frac{1}{x_p^+ x_q^-}} \frac{1 - \frac{1}{x_p^+ x_q^+}}{1 - \frac{1}{x_p^- x_q^-}} \tilde{\sigma}_{pq}^{-2}, \quad (4.60)$$

is regular in the physical region, and in particular has no pole at $x_p^+ = x_q^-$. It is an interesting check that the same holds in the crossed channel, whose exchange diagram




Figure 4.5: On the left the would-be Landau diagram for one left- and one right-moving particle is depicted. This process should be absent. Similarly, the crossed process on the right should be absent, and the corresponding S-matrix element have no pole.

would be as in figure 4.5b, should it exist. Again crossing symmetry relates the two processes in a simple way (because we are scattering highest weight states), *i.e.*

$$\tilde{\mathcal{B}}_{pq}\mathcal{B}_{p^c q} = 1, \quad \text{where } \mathcal{B}_{pq} = \langle \Phi_{++}^L \Phi_{--}^L | \check{\mathcal{S}}_{pq} | \Phi_{--}^L \Phi_{++}^L \rangle, \quad (4.61)$$

which implies the first crossing equation in (4.26). Since $\tilde{\mathcal{B}}_{pq}$ has no singularity at $x_p^+ = x_q^-$ we expect $\mathcal{B}_{p^c q}$ to have no singularity at $x_{p^c}^+ = 1/x_q^-$. Explicitly we have

$$\mathcal{B}_{p^c q} = \frac{(x_{p^c}^- - x_q^-)^2}{(x_{p^c}^- - x_q^+)(x_{p^c}^+ - x_q^-)} \frac{1 - \frac{1}{x_{p^c}^- x_q^+}}{1 - \frac{1}{x_{p^c}^+ x_q^-}} \sigma_{p^c q}^{-2}. \quad (4.62)$$

The rational terms have a pole at $x_{p^c}^+ = 1/x_q^-$, but once the dressing factor is continued to the crossed region as in (A.75), this is canceled by a zero of σ^{-2} , so that the result is non-singular.

Conditions on CDD factors

The fact that we correctly match the structure of single poles in the physical region tells us that for any CDD factors of the form (4.49) we must set $n_1 = n_2 = 0$. However, in principle we could still allow for different solutions of the homogeneous crossing equations. If these are defined on a cover of the rapidity torus and feature no poles in the physical region, for instance, they would not be ruled out by our bound-state analysis. Since the dressing phases we propose live on such a cover, such solutions cannot be ruled out.

4.4 Chapter summary

In this chapter we have formalised the particle-to-antiparticle symmetry of the (non-relativistic) worldsheet theory in terms of a set of crossing equations, in the spirit of what was done in $\text{AdS}_5 \times \text{S}^5$ by Janik [97]. In our case, we find that the particle-to-antiparticle transformation exchanges the left and right representations. This is a qualitatively new feature, which tells us that for consistency of our 2-dimensional worldsheet theory we cannot restrict ourselves to particles of a single target-space chirality, even if they form an irreducible representation of the off-shell symmetry algebra.

The crossing equation couples the LL and LR dressing factors \mathcal{S}_{pq} and $\widetilde{\mathcal{S}}_{pq}$ or equivalently σ_{pq} and $\widetilde{\sigma}_{pq}$. Specifically, it relates one scalar factor evaluated in the physical region with another analytically continued to the crossed region. We see that, were it not for the dressing factors, the S matrix would be a rational function on a double complex torus—instead, it is defined on an infinite cover of it.

The main result of the chapter is an all-loop solution to the crossing equations. This features the well known Beisert-Eden-Staudacher [28] and Hernández-López [144] dressing factors that originally emerged in $\text{AdS}_5 \times \text{S}^5$, together with a novel factor defined by (4.46). This also shows that the BES phase, which essentially constitutes the dressing factors of integrable $\text{AdS}_5/\text{CFT}_4$ and $\text{AdS}_4/\text{CFT}_3$, is not universal to all AdS/CFT duals.

Besides being crossing-symmetric, the dressing factors we found reproduce the bound-state spectrum we expect to find in the physical region. Still, it is unclear whether they are the physical solution of the crossing equations, or should be modified by multiplying them by a solution of the homogeneous equations, which could be highly non-trivial. To check whether this is the case, we can compare the large- \hbar expansion of the dressing factors to independent perturbative calculations. We will successfully do so in chapter 7. Additionally, we could try to investigate further the analytic properties of the S matrix, especially beyond the physical region. For instance, its properties in the region where a mirror theory should be defined⁹ will be important to formulate the mirror thermodynamical Bethe ansatz description that we mentioned in the introduction. So far, such an investigation has not been performed.

Finally, we should remark that in principle the dressing factors of massive particles should also contain some information due to the presence of massless virtual particles. This could appear through a solution of the homogeneous crossing equations, too. At the moment, however, it is not clear what its contribution should be, or whether somehow it has already been accounted for.

⁹This is the region in the middle of the green curves in figure 4.3, corresponding to a shift of $\frac{1}{2}\omega_2$, which in fact reproduces the mirror transformation (1.28).

5 The $\mathfrak{psu}(1, 1|2)^2$ spin chain

In this chapter we will see how the all-loop S matrix can be found in a *spin-chain picture*. Inspired from what happens in the case $\text{AdS}_5 \times \text{S}^5$, which we briefly discussed in the introduction, it is rather natural to assume that strings on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ can be described in terms of a spin chain. If we restrict to the massive sector, the “spins” should be in modules of $\mathfrak{psu}(1, 1|2)^2$. By picking a ground state for the chain, and considering fluctuations around it, we will find an off-shell symmetry algebra. Out of this we will be able to fix a two-body S matrix. While the S matrix that we found out of the Zamolodchikov-Faddeev algebra in chapter 3 describes the reordering of two ZF creation operators, the one we study in this chapter acts on pairs of spin-chain sites. Consistently extending it to an M -sites S matrix will again require a sort of Yang-Baxter equation to hold. Equivalently, we can describe the S matrix as acting on plane-wave excitations (magnons), which will allow us to show that it is in fact equivalent to the worldsheet S matrix, in a sense that we will specify.

This spin-chain picture was the way in which the S matrix was originally derived [S3, S5]. To keep our presentation as homogeneous as possible, here we use slightly different conventions than in refs. [S3, S5].

5.1 The weakly-coupled spin chain

We want to describe the all-loop $\mathfrak{psu}(1, 1|2)^2$ spin-chain dual to free $\text{AdS}_3 \times \text{S}^3$ strings. At weak coupling—that is, for $h \ll 1$ —the spin chain was originally constructed in [112]. There one finds two copies of the superalgebra $\mathfrak{psu}(1, 1|2)$, describing the left- and right-moving sectors of the dual CFT. At leading order in a $h \rightarrow 0$ expansion, left- and right-movers decouple. The spectrum is then described by *two* homogeneous spin-chains with the sites of each transforming in the representation¹ $(\frac{1}{2}; \frac{1}{2})$ of $\mathfrak{psu}(1, 1|2)$ —one for the left and one for the right sector. At higher orders in h the two sectors couple to each other through local interactions. We will be able to account for all of these interactions from

¹To make contact with the coset construction, we take the $\mathfrak{su}(1, 1)$ spin to be positive. Still, the $\frac{1}{2}$ representation of $\mathfrak{su}(1, 1)$ is infinite-dimensional, while the $\frac{1}{2}$ representation of $\mathfrak{su}(2)$ is finite-dimensional. This is in contrast with ref. [S3] where we considered the $(-\frac{1}{2}; \frac{1}{2})$ representation of $\mathfrak{psu}(1, 1|2)$.

symmetry arguments.

The spin-chain algebra and representation

The sites of each $\mathfrak{psu}(1, 1|2)$ spin-chain transform in the infinite-dimensional representation $(\frac{1}{2}; \frac{1}{2})$, consisting of the bosonic $\mathfrak{su}(2)$ doublets $\phi_{\pm}^{(n)}$ and the two sets of fermionic $\mathfrak{su}(2)$ singlets $\psi_{\pm}^{(n)}$, where the index n indicates the $\mathfrak{sl}(2)$ quantum number. In the spin-chain picture it is convenient to consider a real form of $\mathfrak{psu}(1, 1|2)$ which differs from the one we saw in section 2.2.² Let us consider

$$\begin{aligned} [\mathbb{L}_3, \mathbb{L}_{\pm}] &= \pm \mathbb{L}_{\pm}, & [\mathbb{L}_+, \mathbb{L}_-] &= 2 \mathbb{L}_3, \\ [\mathbb{J}_3, \mathbb{J}_{\pm}] &= \pm \mathbb{J}_{\pm}, & [\mathbb{J}_+, \mathbb{J}_-] &= 2 \mathbb{J}_3, \end{aligned} \quad (5.1)$$

for the bosonic charges and

$$\begin{aligned} [\mathbb{L}_3, \mathbb{Q}_{\pm\kappa\iota}] &= \mp \frac{1}{2} \mathbb{Q}_{\pm\kappa\iota}, & [\mathbb{L}_{\pm}, \mathbb{Q}_{\pm\kappa\iota}] &= \mathbb{Q}_{\mp\kappa\iota}, \\ [\mathbb{J}_3, \mathbb{Q}_{a\pm\iota}] &= \pm \frac{1}{2} \mathbb{Q}_{a\pm\iota}, & [\mathbb{J}_{\pm}, \mathbb{Q}_{a\mp\iota}] &= \mathbb{Q}_{a\pm\iota}, \end{aligned} \quad (5.2)$$

with $\kappa = \pm, \iota = \pm$ and $a = \pm$. The anticommutators then are

$$\begin{aligned} \{\mathbb{Q}_{\pm++}, \mathbb{Q}_{\pm--}\} &= \pm \mathbb{L}_{\mp}, & \{\mathbb{Q}_{\pm+-}, \mathbb{Q}_{\pm-+}\} &= \mp \mathbb{L}_{\mp}, \\ \{\mathbb{Q}_{\pm++}, \mathbb{Q}_{-+-}\} &= \mp \mathbb{J}_{\pm}, & \{\mathbb{Q}_{\pm+-}, \mathbb{Q}_{-++}\} &= \pm \mathbb{J}_{\pm}, \\ \{\mathbb{Q}_{\pm\pm\pm}, \mathbb{Q}_{-\mp\mp}\} &= + \mathbb{L}_3 \pm \mathbb{J}_3, & \{\mathbb{Q}_{\pm\pm\mp}, \mathbb{Q}_{-\mp\pm}\} &= - \mathbb{L}_3 \mp \mathbb{J}_3. \end{aligned} \quad (5.3)$$

Further properties of $\mathfrak{psu}(1, 1|2)$ are given in appendix A.1.

Let us focus our attention on the action of the generators on a single site, which may be in any of the states of the $(\frac{1}{2}; \frac{1}{2})$ module. We have that bosonic states are charged under $\mathfrak{su}(2)$

$$\mathbb{J}_3 |\phi_{\pm}^{(n)}\rangle = \pm \frac{1}{2} |\phi_{\pm}^{(n)}\rangle, \quad \mathbb{J}_+ |\phi_{-}^{(n)}\rangle = |\phi_{+}^{(n)}\rangle, \quad \mathbb{J}_- |\phi_{+}^{(n)}\rangle = |\phi_{-}^{(n)}\rangle, \quad (5.4)$$

and all states are charged under $\mathfrak{sl}(2)$

$$\begin{aligned} \mathbb{L}_3 |\phi_{\kappa}^{(n)}\rangle &= \left(\frac{1}{2} + n\right) |\phi_{\kappa}^{(n)}\rangle, & \mathbb{L}_3 |\psi_{\iota}^{(n)}\rangle &= (1 + n) |\psi_{\iota}^{(n)}\rangle, \\ \mathbb{L}_- |\phi_{\kappa}^{(n)}\rangle &= +n |\phi_{\kappa}^{(n-1)}\rangle, & \mathbb{L}_- |\psi_{\iota}^{(n)}\rangle &= +\sqrt{(n+1)n} |\psi_{\iota}^{(n-1)}\rangle, \\ \mathbb{L}_+ |\phi_{\kappa}^{(n)}\rangle &= -(n+1) |\phi_{\kappa}^{(n+1)}\rangle, & \mathbb{L}_+ |\psi_{\iota}^{(n)}\rangle &= -\sqrt{(n+2)(n+1)} |\psi_{\iota}^{(n+1)}\rangle. \end{aligned} \quad (5.5)$$

Finally, the supercharges act as

$$\begin{aligned} \mathbb{Q}_{-\pm\iota} |\phi_{\mp}^{(n)}\rangle &= \pm \sqrt{n+1} |\psi_{\iota}^{(n)}\rangle, & \mathbb{Q}_{+\pm\iota} |\phi_{\mp}^{(n)}\rangle &= \pm \sqrt{n} |\psi_{\iota}^{(n-1)}\rangle, \\ \mathbb{Q}_{-\kappa\pm} |\psi_{\mp}^{(n)}\rangle &= \mp \sqrt{n+1} |\phi_{\kappa}^{(n+1)}\rangle, & \mathbb{Q}_{+\kappa\pm} |\psi_{\mp}^{(n)}\rangle &= \mp \sqrt{n+1} |\phi_{\kappa}^{(n)}\rangle. \end{aligned} \quad (5.6)$$

²This difference in the real form is familiar from $\text{AdS}_5/\text{CFT}_4$, see *e.g.* ref. [152].




Figure 5.1: An illustration of the short $(\frac{1}{2}; \frac{1}{2})$ module, where the action of the (super)charges is represented up to the numerical coefficients. These are given in eqs. (5.4–5.6).

Therefore, the highest weight state $|\phi_+^{(0)}\rangle$ is annihilated by the $\mathfrak{su}(2)$ grading raising operators $\mathbb{Q}_{+\pm\pm}$, and by the two generators $\mathbb{Q}_{-\pm\pm}$. Hence, the representation $(\frac{1}{2}; \frac{1}{2})$ is a short representation, satisfying the shortening conditions

$$\{\mathbb{Q}_{+-\mp}, \mathbb{Q}_{-+\pm}\} |\phi_+^{(0)}\rangle = \pm(\mathbb{L}_3 - \mathbb{J}_3) |\phi_+^{(0)}\rangle = 0. \quad (5.7)$$

The action of the generators on part of the module is depicted in figure 5.1. We will take the sites of the left and right spin-chains to transform in identical modules. At the very end of our construction, we will see that self-consistency dictates the left and right algebras to be in different gradings, as it was the case in the coset construction of chapter 2.

The ground state

The states of the left- and right-moving spin-chains of length ℓ transform in the ℓ -fold tensor product of the above representation. The ground state of the full spin-chain is given by

$$|0\rangle_\ell = \left| (\phi_+^{(0)})^\ell \right\rangle \otimes \left| (\phi_+^{(0)})^\ell \right\rangle. \quad (5.8)$$

This is the highest weight state of the short $(\frac{\ell}{2}; \frac{\ell}{2}) \otimes (\frac{\ell}{2}; \frac{\ell}{2})$ representation of the superalgebra $\mathfrak{psu}(1, 1|2)_L \oplus \mathfrak{psu}(1, 1|2)_R$. Such a choice preserves as much supersymmetry as possible. Specifically, the ground state is preserved by eight supercharges $\mathbb{Q}^{jL,R}$ and $\bar{\mathbb{Q}}^{jL,R}$, with $j = 1, 2$, as well as two central charges $\mathbb{H}^{L,R}$. In terms of the $\mathfrak{psu}(1, 1|2)$

generators they are given by

$$\begin{aligned} \mathbb{Q}^{1\text{L}} &= \mathbb{Q}^{\text{L}}_{-++}, & \mathbb{Q}^{2\text{L}} &= -\mathbb{Q}^{\text{L}}_{--+}, & \overline{\mathbb{Q}}^{1\text{L}} &= \mathbb{Q}^{\text{L}}_{+--}, & \overline{\mathbb{Q}}^{2\text{L}} &= \mathbb{Q}^{\text{L}}_{-++}, \\ \mathbb{Q}^{1\text{R}} &= \mathbb{Q}^{\text{R}}_{-++}, & \mathbb{Q}^{2\text{R}} &= -\mathbb{Q}^{\text{R}}_{--+}, & \overline{\mathbb{Q}}^{1\text{R}} &= \mathbb{Q}^{\text{R}}_{+--}, & \overline{\mathbb{Q}}^{2\text{R}} &= \mathbb{Q}^{\text{R}}_{-++}, \end{aligned} \quad (5.9)$$

and

$$\mathbb{H}^{\text{L}} = \mathbb{L}_3^{\text{L}} - \mathbb{J}_3^{\text{L}}, \quad \mathbb{H}^{\text{R}} = \mathbb{L}_3^{\text{R}} - \mathbb{J}_3^{\text{R}}. \quad (5.10)$$

This forms two (one left, one right) copies of the $(\mathfrak{su}(1|1)^2)/\mathfrak{u}(1)$ algebra, where the quotient is due to the fact that we have the same Hamiltonian for $j = 1, 2$:

$$\{\mathbb{Q}^{j\text{L}}, \overline{\mathbb{Q}}^{k\text{L}}\} = \delta^{jk} \mathbb{H}^{\text{L}}, \quad \{\mathbb{Q}^{j\text{R}}, \overline{\mathbb{Q}}^{k\text{R}}\} = \delta^{jk} \mathbb{H}^{\text{R}}. \quad (5.11)$$

The charges \mathbb{H}^{L} and \mathbb{H}^{R} are the left- and right-moving spin-chain Hamiltonians. Let us define

$$\mathbb{H} = \mathbb{H}^{\text{L}} + \mathbb{H}^{\text{R}}, \quad \mathbb{M} = \mathbb{H}^{\text{L}} - \mathbb{H}^{\text{R}}, \quad (5.12)$$

where the positive-definite combination \mathbb{H} has the interpretation of the spin-chain Hamiltonian, and may depend on the momenta of the spin-chain excitations. The central charge \mathbb{M} measures an angular momentum in $\text{AdS}_3 \times \text{S}^3$ and should be quantised.

This symmetry algebra appears *at small coupling*, when the L and R sectors are decoupled. It also coincides with the *on-shell symmetry algebra* of section 2.4. This picture is similar to what happens in $\text{AdS}_5 \times \text{S}^5$, where the small-coupling spin chain does not display dependence on the additional momentum-dependent central charges.

For later convenience, let us introduce two additional generators $\mathbb{V}_1^{\text{L,R}}$ and $\mathbb{V}_2^{\text{L,R}}$ acting as outer automorphisms. These can be constructed from the $\mathfrak{psu}(1, 1|2)$ generators $\mathbb{J}_3^{\text{L,R}}$ and the automorphisms $\mathbb{U}^{\text{L,R}}$ defined in eq. (A.7),

$$\begin{aligned} \mathbb{V}_1^{\text{L,R}} &= -\mathbb{U}^{\text{L,R}} - \mathbb{J}_3^{\text{L,R}}, & \mathbb{V}_2^{\text{L,R}} &= +\mathbb{U}^{\text{L,R}} - \mathbb{J}_3^{\text{L,R}}, \\ \mathbb{V}_1 &= +\mathbb{V}_1^{\text{L}} - \mathbb{V}_1^{\text{R}}, & \mathbb{V}_2 &= +\mathbb{V}_2^{\text{L}} - \mathbb{V}_2^{\text{R}}. \end{aligned} \quad (5.13)$$

While all four left and right generators are automorphisms of $\mathfrak{psu}(1, 1|2)^2$, only the combinations \mathbb{V}_1 and \mathbb{V}_2 annihilate the vacuum (5.8). Note also that \mathbb{V}_1 commutes with $\mathbb{Q}^{2\text{L,R}}$ and $\overline{\mathbb{Q}}^{2\text{L,R}}$, while \mathbb{V}_2 commutes with $\mathbb{Q}^{1\text{L,R}}$ and $\overline{\mathbb{Q}}^{1\text{L,R}}$. The commutation relations involving the supercharges then read

$$\begin{aligned} [\mathbb{V}_j, \mathbb{Q}^{k\text{L}}] &= -\delta_j^k \mathbb{Q}^{k\text{L}}, & [\mathbb{V}_j, \overline{\mathbb{Q}}^{k\text{L}}] &= +\delta_j^k \overline{\mathbb{Q}}^{k\text{L}}, \\ [\mathbb{V}_j, \mathbb{Q}^{k\text{R}}] &= +\delta_j^k \mathbb{Q}^{k\text{R}}, & [\mathbb{V}_j, \overline{\mathbb{Q}}^{k\text{R}}] &= -\delta_j^k \overline{\mathbb{Q}}^{k\text{R}}. \end{aligned} \quad (5.14)$$

The generators \mathbb{V}_j give useful restrictions on the allowed deformations of the weak-coupling representations. Taking them into account, we regroup the symmetry algebra into two copies of $\mathfrak{u}(1) \oplus \mathfrak{su}(1|1)^2$, with the generators given by

$$\{\mathbb{Q}^{1\text{L}}, \mathbb{Q}^{1\text{R}}, \overline{\mathbb{Q}}^{1\text{L}}, \overline{\mathbb{Q}}^{1\text{R}}, \mathbb{H}^{\text{L}}, \mathbb{H}^{\text{R}}, \mathbb{V}_1\} \quad \text{and} \quad \{\mathbb{Q}^{2\text{L}}, \mathbb{Q}^{2\text{R}}, \overline{\mathbb{Q}}^{2\text{L}}, \overline{\mathbb{Q}}^{2\text{R}}, \mathbb{H}^{\text{L}}, \mathbb{H}^{\text{R}}, \mathbb{V}_2\}, \quad (5.15)$$




Figure 5.2: The action of the supercharges $Q^j = Q^{jL,R}$ and $\bar{Q}^j = \bar{Q}^{jL,R}$ on either of the left- or right-bifundamental representation (5.17). It takes the same form of figure 2.2.

respectively. This splitting is familiar from section 2.4, where it corresponded to a tensor-product structure in the representation of the excitations. We will see that the same holds here.

Excitations at weak coupling

To construct excited spin-chain states we replace one or more of the ground state sites by any other state in the same module. We can classify these excitations by their eigenvalues under the left and right spin-chain Hamiltonians \mathbb{H}^L and \mathbb{H}^R at zero coupling. Let us consider excitations in the left sector. Replacing one of the highest weight states $\phi_{+}^{(0)}$ by the scalar $\phi_{-}^{(n)}$ or $\phi_{+}^{(n)}$ increases the eigenvalue of \mathbb{H}^L by n or $n+1$, respectively. Similarly, insertion of a fermion $\psi_{\pm}^{(n)}$ also adds n to the energy. The lightest excitations are therefore

$$\phi_{-}^{(0)}, \quad \psi_{+}^{(0)}, \quad \psi_{-}^{(0)}, \quad \phi_{+}^{(1)}. \quad (5.16)$$

It is easy to see that the charges of any heavier excitation can be reproduced by considering a combination of the four states above. These states form a four-dimensional bifundamental representation of either of the two $\mathfrak{psu}(1|1)^2$ algebras (5.11), as illustrated in figure 5.2. To emphasise this we introduce the notation

$$\begin{aligned} \Phi_{++}^L &= \phi_{-}^{(0)} \otimes \phi_{+}^{(0)}, & \Phi_{--}^L &= \phi_{+}^{(1)} \otimes \phi_{+}^{(0)}, & \Phi_{-+}^L &= \psi_{+}^{(0)} \otimes \phi_{+}^{(0)}, & \Phi_{+-}^L &= -\psi_{-}^{(0)} \otimes \phi_{+}^{(0)}, \\ \Phi_{++}^R &= \phi_{+}^{(0)} \otimes \phi_{-}^{(0)}, & \Phi_{--}^R &= \phi_{+}^{(0)} \otimes \phi_{+}^{(1)}, & \Phi_{-+}^R &= \phi_{+}^{(0)} \otimes \psi_{+}^{(0)}, & \Phi_{+-}^R &= -\phi_{+}^{(0)} \otimes \psi_{-}^{(0)}, \end{aligned} \quad (5.17)$$

where the tensor product is over left and right sites. As before, the excitations $\Phi_{\pm\pm}$ are bosons while $\Phi_{\pm\mp}$ are fermions in either the left and right sector. We will also define the short-hand for a vacuum site,

$$Z = \phi_{+}^{(0)} \otimes \phi_{+}^{(0)}. \quad (5.18)$$

In figure 5.3 the structure of the spin chain is represented pictorially.




Figure 5.3: A pictorial representation of the $\mathfrak{psu}(1,1|2)^2$ spin chain. We can think of it as two (left and right) spin chains, each with ground state $\phi_+^{(0)}$. Excitations of the whole chain are considered as left- or right- moving depending on whether we excite a spin in the left or right chain.

Fundamental representations

As we did in the string picture, we can make the bifundamental nature of the representation above more explicit. Let us consider the left module. We introduce a fundamental $\mathfrak{su}(1|1)$ representation with basis $(\phi^L|\psi^L)$, and the generators \mathfrak{q} , $\bar{\mathfrak{q}}$ and \mathfrak{h} acting as

$$\mathfrak{q}^L |\phi^L\rangle = a^L |\psi^L\rangle, \quad \bar{\mathfrak{q}}^L |\psi^L\rangle = \bar{a}^L |\phi^L\rangle, \quad \mathfrak{h}^L = |a^L|^2 \mathbb{I}^L. \quad (5.19)$$

where \mathbb{I}^L is the identity on the $(\phi^L|\psi^L)$ representation. The representation for right-movers is similar,

$$\mathfrak{q}^R |\phi^R\rangle = a^R |\psi^R\rangle, \quad \bar{\mathfrak{q}}^R |\psi^R\rangle = \bar{a}^R |\phi^R\rangle, \quad \mathfrak{h}^R = |a^R|^2 \mathbb{I}^R. \quad (5.20)$$

We can also write down the action of the automorphism generators $\mathfrak{v}^{L,R}$ and \mathfrak{v} , which correspond to $\mathbb{V}_j^{L,R}$ and \mathbb{V}_j respectively. This is found from the charges written in table 5.1. The parameter v appearing there is a label of the representation, and as we will see it is natural to take it to be the same in the left and right ones. Looking at the eigenvalues, we see that indeed \mathfrak{v} annihilates the vacuum, whereas each of the $\mathfrak{v}^{L,R}$ do not, and instead measure its length.

Back to the bifundamental representations

Focusing *e.g.* on the left module, we can identify the fields in the bifundamental representations by setting like in (2.104)

$$|\Phi_+^{L,R}\rangle = |\phi^{L,R}\rangle, \quad |\Phi_-^{L,R}\rangle = |\psi^{L,R}\rangle, \quad (5.21)$$

so that $\Phi_{\kappa\ell}^{L,R} = \Phi_{\kappa}^{L,R} \otimes \Phi_{\ell}^{L,R}$ once again. The supercharges \mathbb{Q}^{1L} and $\bar{\mathbb{Q}}^{1L}$ only act on the first index of $\Phi_{\kappa\ell}$, while \mathbb{Q}^{2L} and $\bar{\mathbb{Q}}^{2L}$ act on the second index. Hence they can be written as

$$\mathbb{Q}^{1L} = \mathfrak{q}^L \otimes \mathbb{I}, \quad \bar{\mathbb{Q}}^{1L} = \bar{\mathfrak{q}}^L \otimes \mathbb{I}, \quad \mathbb{Q}^{2L} = \mathbb{I} \otimes \mathfrak{q}^L, \quad \bar{\mathbb{Q}}^{2L} = \mathbb{I} \otimes \bar{\mathfrak{q}}^L. \quad (5.22)$$

	$2 \mathbb{V}^L$	$2 \mathbb{V}^R$	$2 \mathbb{V}$
$ 0\rangle$	$-\ell$	$-\ell$	0
$ \phi^L\rangle$	$-\ell + v$	$-\ell$	$+v$
$ \psi^L\rangle$	$-\ell + v - 1$	$-\ell$	$+v - 1$
$ \phi^R\rangle$	$-\ell$	$-\ell + v$	$-v$
$ \psi^R\rangle$	$-\ell$	$-\ell + v - 1$	$-v + 1$
\mathbb{Q}^L	-1	0	-1
$\bar{\mathbb{Q}}^L$	$+1$	0	$+1$
\mathbb{Q}^R	0	-1	$+1$
$\bar{\mathbb{Q}}^R$	0	$+1$	-1

Table 5.1: Charges under the automorphisms $\mathbb{V}^{L,R}$ and \mathbb{V} of the ground state, of the single excitation states in the left- and right-moving multiplets as well as of the supercharges. All the spin-chain states have length ℓ . To make the table less cluttered, the charges have been rescaled.

The action of $\mathbb{V}_j^{L,R}$ and \mathbb{V}_j in terms of $\mathbb{V}^{L,R}$ and \mathbb{V} takes the same form.

The tensor-product structure is compatible with the fermion grading. In particular this means that we get a minus sign from commuting a supercharge through a fermion when we act with the charges of the second type on a state with a fermion in the first part of the tensor product. Hence, the left-moving generators act as

$$\begin{aligned}
\mathbb{Q}^{1L} |\Phi_{++}^L\rangle &= a^L |\Phi_{-+}^L\rangle, & \mathbb{Q}^{1L} |\Phi_{+-}^L\rangle &= a^L |\Phi_{--}^L\rangle, \\
\bar{\mathbb{Q}}^{1L} |\Phi_{-+}^L\rangle &= \bar{a}^L |\Phi_{++}^L\rangle, & \bar{\mathbb{Q}}^{1L} |\Phi_{--}^L\rangle &= \bar{a}^L |\Phi_{+-}^L\rangle, \\
\mathbb{Q}^{2L} |\Phi_{++}^L\rangle &= a^L |\Phi_{+-}^L\rangle, & \mathbb{Q}^{2L} |\Phi_{-+}^L\rangle &= -a^L |\Phi_{--}^L\rangle, \\
\bar{\mathbb{Q}}^{2L} |\Phi_{+-}^L\rangle &= \bar{a}^L |\Phi_{++}^L\rangle, & \bar{\mathbb{Q}}^{2L} |\Phi_{--}^L\rangle &= -\bar{a}^L |\Phi_{-+}^L\rangle,
\end{aligned} \tag{5.23}$$

with the right-moving charges acting trivially. Comparing the above representation with (5.4) we find that the central charge \mathbb{H}^L has eigenvalue $|a^L|^2 = 1$. Once again, this is precisely what happens for the *on-shell* (zero-momentum) one-particle representation in string theory. In the same way, we can introduce the right-moving excitations $\Phi_{\pm\pm}^R$ and $\Phi_{\pm\mp}^R$, transforming in a bifundamental representation of the right-moving $\mathfrak{psu}(1|1)^2$ algebra. Again we find that at zero coupling it is $|a^R|^2 = 1$.

5.2 The dynamical spin chain

At non-vanishing coupling $h > 0$, the spin-chain Hamiltonian \mathbb{H} should depend on h and on the momentum of the excitations. This requires the bifundamental representations discussed above to be deformed—a procedure akin to going from the one-particle, on-shell $p = 0$ representation to a general *off-shell* one.

This deformation should be done in such a way that the angular momentum \mathbb{M} remains quantised, *i.e.* undeformed. This can be done if we allow the right generators to act nontrivially on the left-moving excitations, and *vice versa*, which is what we expect from chapters 2 and 3.

Constructing the central extension

Let us investigate how to centrally extend the algebra. For simplicity, let us focus on constructing the extension of $\mathfrak{u}(1) \oplus \mathfrak{su}(1|1)^2$ in (5.15). The generalisation to the full algebra and tensor-product representation will be straightforward.

We want the left representation to be charged under all of the supercharges. Focusing on the highest weight state $|\phi^L\rangle$, we have two possibilities

$$(I): \quad \mathfrak{q}^R |\phi^L\rangle \neq 0, \quad \text{or} \quad (II): \quad \bar{\mathfrak{q}}^R |\phi^L\rangle \neq 0. \quad (5.24)$$

In case (I), in the right-hand side there should be a fermion whose charge under \mathfrak{v} is

$$\mathfrak{v} \mathfrak{q}^R |\phi^L\rangle = \frac{1}{2}(v+1) \mathfrak{q}^R |\phi^L\rangle. \quad (5.25)$$

Looking back at table 5.1, we see that there are no such fermionic states on the vacuum $|0\rangle_\ell$ preserved by \mathfrak{v} . On the other hand, in case (II) we have

$$\mathfrak{v} \bar{\mathfrak{q}}^R |\phi^L\rangle = \frac{1}{2}(v-1) \bar{\mathfrak{q}}^R |\phi^L\rangle, \quad (5.26)$$

so that the state $|\psi^L\rangle$ is a good candidate to appear on the right hand side. Therefore, let us restrict to case (II), and further investigate the central extension by looking at the action of \mathfrak{v}^L and \mathfrak{v}^R . We have

$$\begin{aligned} \mathfrak{v}^L \bar{\mathfrak{q}}^R |\phi^L\rangle &= \frac{1}{2}(v-1-(\ell-1)) \bar{\mathfrak{q}}^R |\phi^L\rangle, \\ \mathfrak{v}^R \bar{\mathfrak{q}}^R |\phi^L\rangle &= \frac{1}{2}(-(\ell-1)) \bar{\mathfrak{q}}^R |\phi^L\rangle. \end{aligned} \quad (5.27)$$

We can interpret this by saying that $\bar{\mathfrak{q}}^R$ acts on a state in the left-representation of length ℓ by exchanging a boson with fermion and *reducing the length* of the chain by one. If we write separately the left and right sites of the chain in a tensor product form, we have

$$\bar{\mathfrak{q}}^R |\phi^L\rangle = \bar{\mathfrak{q}}^R (|\phi Z^{\ell-1}\rangle) \approx |\psi Z^{\ell-2}\rangle. \quad (5.28)$$

More compactly, we can denote this action by $\bar{\mathfrak{q}}^R |\phi\rangle \approx |\psi Z^-\rangle$, to indicate that the state on the right hand side has been constructed on a vacuum of a length shorter by one with respect to the one on the left hand side. Still, the vacuum is of the form (5.8), which is why this action preserves \mathfrak{v} .³ It is also convenient to introduce a symbol Z^+ corresponding to

³The central extension (I) can also be understood in terms of length-changing effects, that however do not preserve (5.8). In fact, they correspond to adding one site to the left spin chain and removing one to the right one, or *vice versa*. This would force us to consider a much larger set of vacua $|0\rangle = |Z^L\rangle \otimes |Z^{\bar{L}}\rangle$ which has no analogue in the string theory. This central extension is further discussed in [S5].

the *insertion* of an additional vacuum site. We can finally write the action of centrally extended algebra on the left representation as

$$\begin{aligned} \mathfrak{q}^L |\phi^L\rangle &= a^L |\psi^L\rangle, & \bar{\mathfrak{q}}^L |\psi^L\rangle &= \bar{a}^L |\phi^L\rangle, \\ \bar{\mathfrak{q}}^R |\phi^L\rangle &= \bar{b}^L |\psi^L Z^-\rangle, & \mathfrak{q}^R |\psi^L\rangle &= b^L |\phi^L Z^+\rangle, \end{aligned} \quad (5.29)$$

and similarly on the right one as

$$\begin{aligned} \mathfrak{q}^R |\phi^R\rangle &= a^R |\psi^R\rangle, & \bar{\mathfrak{q}}^R |\psi^R\rangle &= \bar{a}^R |\phi^R\rangle, \\ \bar{\mathfrak{q}}^L |\phi^R\rangle &= \bar{b}^R |\psi^R Z^-\rangle, & \mathfrak{q}^L |\psi^R\rangle &= b^R |\phi^R Z^+\rangle, \end{aligned} \quad (5.30)$$

This action is very similar to (3.44), but the algebra is now realised in terms of a *dynamical spin chain*, where the symmetry generators may add or remove sites. Dynamical effects are a common feature of the spin chains appearing in AdS/CFT [26, 153, 154], see also ref. [155]. In the one-particle representation we can ignore length-changing effects as long as we restrict to *asymptotic states* with $\ell \rightarrow \infty$, which we will always do in this chapter. Then, the algebra that we constructed closes to a central extension of $\mathfrak{su}(1|1)^2$

$$\begin{aligned} \{\mathfrak{q}^L, \bar{\mathfrak{q}}^L\} &= \mathfrak{h}^L, & \{\mathfrak{q}^R, \bar{\mathfrak{q}}^R\} &= \mathfrak{h}^R, \\ \{\mathfrak{q}^L, \mathfrak{q}^R\} &= \mathfrak{c}, & \{\bar{\mathfrak{q}}^L, \bar{\mathfrak{q}}^R\} &= \bar{\mathfrak{c}}, \end{aligned} \quad (5.31)$$

where on the one-particle representation

$$\mathfrak{c} = a^L b^L \mathbb{I}^L + a^R b^R \mathbb{I}^R, \quad \bar{\mathfrak{c}} = \bar{a}^L \bar{b}^L \mathbb{I}^L + \bar{a}^R \bar{b}^R \mathbb{I}^R. \quad (5.32)$$

Eq. (5.31) defines the same algebra we found from analysis of the off-shell symmetries of asymptotic string states (3.35).

The bifundamental representations

It is easy to reformulate our result in terms of the two bifundamental representations of $\mathfrak{su}(1|1)_{\text{c.e.}}^4$, which now features two additional central charges \mathbb{C} and $\bar{\mathbb{C}}$ appearing as

$$\{\mathbb{Q}^{jL}, \mathbb{Q}^{kR}\} = \delta^{jk} \mathbb{C}, \quad \{\bar{\mathbb{Q}}^{jL}, \bar{\mathbb{Q}}^{kR}\} = \delta^{jk} \bar{\mathbb{C}}. \quad (5.33)$$

The non-trivial action of the right generators on the left representation is then

$$\begin{aligned} \mathbb{Q}^{1R} |\Phi_{--}^L\rangle &= b^L |\Phi_{--}^L Z^+\rangle, & \mathbb{Q}^{1R} |\Phi_{-+}^L\rangle &= b^L |\Phi_{++}^L Z^+\rangle, \\ \bar{\mathbb{Q}}^{1R} |\Phi_{++}^L\rangle &= \bar{b}^L |\Phi_{-+}^L Z^-\rangle, & \bar{\mathbb{Q}}^{1R} |\Phi_{--}^L\rangle &= \bar{b}^L |\Phi_{--}^L Z^-\rangle, \\ \mathbb{Q}^{2R} |\Phi_{--}^L\rangle &= -b^L |\Phi_{-+}^L Z^+\rangle, & \mathbb{Q}^{2R} |\Phi_{-+}^L\rangle &= b^L |\Phi_{++}^L Z^+\rangle, \\ \bar{\mathbb{Q}}^{2R} |\Phi_{++}^L\rangle &= \bar{b}^L |\Phi_{-+}^L Z^-\rangle, & \bar{\mathbb{Q}}^{2R} |\Phi_{-+}^L\rangle &= -\bar{b}^L |\Phi_{--}^L Z^-\rangle. \end{aligned} \quad (5.34)$$

The action on the supercharges on the right representation can be found in a similar way.

In what follows, it will be usually be easier to work in terms of $\mathfrak{su}(1, 1)_{\text{c.e.}}^2$ keeping in mind that our results extend straightforwardly to $\mathfrak{psu}(1, 1)_{\text{c.e.}}^4$.

Magnons

Since the supercharges and in particular \mathbb{H} are momentum-dependent we will consider spin-chain states in which the excitations carry specific momenta—these are the objects that ultimately we will want to scatter. A one-excitation state can then be written as a plane wave⁴

$$|\mathcal{X}_p\rangle = \sum_{n=1}^{\ell} e^{-ipn} |Z^{n-1} \mathcal{X} Z^{\ell-n}\rangle, \quad (5.35)$$

where \mathcal{X} is any left or right excitation. Of course we can always think of localizing an excitation by constructing an appropriate wave packet. It is now straightforward to generalise this form to the case of multiple excitations. For two excitations we have *e.g.*

$$|\mathcal{X}_p \mathcal{Y}_q\rangle = \sum_{n_1 < n_2}^{\ell} e^{-i(pn_1 + qn_2)} |ZZ \dots ZZ \mathcal{X} ZZ \dots ZZ \mathcal{Y} ZZ \dots ZZ\rangle, \quad (5.36)$$

with $p > q$ and the excitations sitting at positions n_1 and n_2 . We restrict to *asymptotic* states, where the spin chain is considered to be very long, $\ell \rightarrow \infty$, and the excitations are well separated. The interactions are then described by the spin-chain S matrix $\mathcal{S}(p_1, p_2)$ permuting the order of excitations along the chain.

The *length-changing* action on the spin-chain excitations takes a simple form on the magnons. Adding or removing a vacuum site by Z^{\pm} in the plane-wave ansatz we get

$$|Z^{\pm} \mathcal{X}_p\rangle = e^{\pm ip} |\mathcal{X}_p Z^{\pm}\rangle, \quad (5.37)$$

i.e. commutation with the length-changing effects result in a momentum-dependent phase. By these relations we can always shift any insertions of Z^{\pm} through all excitations and collect them at the right end of the state, and since we are dealing with asymptotic states, identify $|\mathcal{X}_p Z^{\pm}\rangle = |\mathcal{X}_p\rangle$. Using the identification of length-changing effects with phase shifts we will obtain a non-trivial coproduct similar to the one that in the NLSM appeared due to the non-local field x_- .

Charge action on multiparticle states

Suppose now that we want to act with a (super)charge on an asymptotic state containing two magnons, for instance $|\phi_p^L \phi_q^L\rangle$. How to do this follows from the natural action of the charges in the spin-chain representation, whereby a charge acts separately on every site of the chain as a derivation. To take statistics into account, such action should be graded,

⁴We have the possibility of choosing the plane-wave coefficient to be $e^{\pm ipn}$. Here we pick the negative sign, in contrast with the original choice of ref. [S5], to more easily compare with the string theory results. The two resulting S matrices are related by a change of basis, as we will discuss in the summary to this chapter.

so that every time a supercharge is anticommutated (from the left, in our convention) past a fermionic site in the chain, we pick up a minus sign. Therefore, we find *e.g.*

$$\mathfrak{q}^L |\phi_p^L \phi_q^L\rangle = a_p^L |\psi_p^L \phi_q^L\rangle + a_q^L |\phi_p^L \psi_q^L\rangle, \quad (5.38)$$

where we made explicit the dependence of the representation coefficients on the momentum, while for two fermions it would be

$$\bar{\mathfrak{q}}^L |\psi_p^L \psi_q^L\rangle = \bar{a}_p^L |\phi_p^L \psi_q^L\rangle - \bar{a}_q^L |\psi_p^L \phi_q^L\rangle. \quad (5.39)$$

It is particularly interesting to look at the action of the supercharges that give rise to the central extension:

$$\begin{aligned} \mathfrak{q}^R |\psi_p^L \psi_q^L\rangle &= b_p^L |\phi_p^L Z^+ \psi_q^L\rangle - b_q^L |\psi_p^L \phi_q^L Z^+\rangle \\ &= b_p^L e^{iq} |\phi_p^L \psi_q^L Z^+\rangle - b_q^L |\psi_p^L \phi_q^L Z^+\rangle \\ &\approx b_p^L e^{iq} |\phi_p^L \psi_q^L\rangle - b_q^L |\psi_p^L \phi_q^L\rangle, \end{aligned} \quad (5.40)$$

where in the last line we used that we are dealing with asymptotic states. We have that *length-changing effects induce a non-trivial coproduct* on asymptotic multi-magnon states. This will dictate a specific form for the coefficient of the one-particle representation, as well as for the form of the two-particle one.

One-particle representation coefficients

Let us consider the left-moving representation. We already know that the coefficients of the undeformed algebra, which coincide with ones we evaluated at zero momentum, satisfy

$$|a_{p=0}^L|^2 = 1, \quad |b_{p=0}^L|^2 = 0. \quad (5.41)$$

The action of \mathfrak{c} and $\bar{\mathfrak{c}}$ on a one-particle state is given by $a_p^L b_p^L$ and $\bar{a}_p^L \bar{b}_p^L$ respectively, that should vanish at $p = 0$, too. The action on the two particle states is more interesting: following the discussion of the previous subsection we get that it must be

$$a_p^L b_p^L e^{iq} + a_q^L b_q^L = 0 \quad \text{if} \quad p + q = 0, \quad (5.42)$$

Similar equations hold for the right-moving excitations by exchanging $L \leftrightarrow R$. Finally, we must require that the angular momentum \mathfrak{m} remains quantised for any value of p , which gives

$$|a_p^L|^2 - |b_p^L|^2 = +1, \quad |b_p^R|^2 - |a_p^R|^2 = -1, \quad (5.43)$$

on the left and right representations respectively. Condition (5.42) can be solved by setting

$$a_p^L b_p^L = i \frac{\hbar}{2} (e^{ip} - 1) \quad \Rightarrow \quad \mathfrak{c} = i \frac{\hbar}{2} (e^{ip} - 1) \mathbb{I}, \quad (5.44)$$

and similarly for \bar{c} . We therefore find the non-linear form of the central charges from the length-changing effects, rather than from the presence of x_- . While in the string-theoretical description the presence of the complex exponential was due to the choice of the light-cone geodesics, here it arises from the plane-wave ansatz for the magnons.

Together with their conjugates, the parameters (5.44) can be used to fix the dispersion relation. In fact, as in section 3.2 we are dealing with short representations of $\mathfrak{su}(1|1)_{c.e.}^2$ or equivalently $\mathfrak{psu}(1|1)_{c.e.}^4$, so that we have the shortening condition

$$\mathbb{H}^2 = \mathbb{M}^2 + \bar{\mathbb{C}}\mathbb{C} = \mathbb{I} + \bar{\mathbb{C}}\mathbb{C}. \quad (5.45)$$

In terms of a_p and b_p we can find the dispersion relation by taking the positive branch of the square root of the shortening condition:

$$\omega(p) = |a_p^L|^2 + |b_p^L|^2 = \sqrt{1 + |a_p^L|^2 |b_p^L|^2} = \sqrt{1 + 4h^2 \sin^2\left(\frac{p}{2}\right)}. \quad (5.46)$$

Taking into account all of these conditions, we can parametrise the one-particle representation as we did in (3.48)

$$\begin{aligned} a_L = a_R &= e^{\frac{1}{4}ip} \eta_p, & \bar{a}_L = \bar{a}_R &= e^{-\frac{3}{4}ip} \eta_p, \\ b_L = b_R &= -\frac{e^{-\frac{3}{4}ip}}{x_p^-} \eta_p, & \bar{b}_L = \bar{b}_R &= -\frac{e^{\frac{1}{4}ip}}{x_p^+} \eta_p, \end{aligned} \quad (5.47)$$

where the Zhukovski parameters x_p^\pm and η_p are given by eqs. (3.47) and (3.49). Once again the symmetry between the left and right representations is reflected by having the same choice of parameters.

Two-particle representation

As we have seen, using the spin-chain picture we can automatically build the two-particle representation on the space of magnons. It is useful to work with matrices, and to this end we introduce a basis \mathcal{B} which takes the same form of (3.43), where now the excitations are interpreted as magnons. Then it is easy to check that the charges $q^{L,R}$ and their conjugates have the same form as in (3.44). What *is different*, however, is the action on the two-particle states. Before we had the expression (3.59) whereby the coproduct would be deformed by a factor of $e^{\pm\frac{1}{2}p}$ when the charge acts on the second-particle space. Here, the momentum-dependence by $e^{\pm iq}$ is on the first-magnon space, and *only on one of the two representations*, as in (5.40).

To express this, let us introduce two matrices \mathcal{L}_p and \mathcal{R}_p that act on the left and right representation exclusively,

$$\mathcal{L}_p = e^{ip} \mathbb{I}^L + \mathbb{I}^R, \quad \mathcal{R}_p = \mathbb{I}^L + e^{ip} \mathbb{I}^R. \quad (5.48)$$

Then, on a two-magnon state we have

$$\begin{aligned}
(q^L(p, q))_{(12)} &= (\mathcal{R}_q q^L(p)) \otimes \mathbb{I} + \Sigma \otimes q^L(q), \\
(q^R(p, q))_{(12)} &= (\mathcal{L}_q q^R(p)) \otimes \mathbb{I} + \Sigma \otimes q^R(q), \\
(\bar{q}^L(p, q))_{(12)} &= (\mathcal{R}_q^\dagger \bar{q}^L(p)) \otimes \mathbb{I} + \Sigma \otimes \bar{q}^L(q), \\
(\bar{q}^R(p, q))_{(12)} &= (\mathcal{L}_q^\dagger \bar{q}^R(p)) \otimes \mathbb{I} + \Sigma \otimes \bar{q}^R(q).
\end{aligned} \tag{5.49}$$

Here we made the fermion signs explicit by means of the matrix Σ , see eq. (3.60).

5.3 The spin-chain S matrix

Following our discussion, we are now in a position to derive the two-body spin-chain S matrix, which should be invariant under the dynamical symmetry algebra that we constructed. As in chapter 3, it is easier to first derive our results for $\mathfrak{su}(1|1)_{\text{c.e.}}^2$.

We have two interpretations for such an S matrix: on the one hand, we can think of it as an operator \mathcal{S} that acts on pairs of spin-chain sites, possibly inserting or removing vacuum sites. On the other hand, we reduce its action to the one of a 16×16 matrix \mathbb{S} (or $\check{\mathbb{S}}$, in the convention of chapter 3) on the vector space $\mathcal{V}_{\text{magn}}(p) \otimes \mathcal{V}_{\text{magn}}(q)$ where now the symmetries have a momentum-dependent coproduct (5.49). The former condition is useful to re-derive some properties of the spin chain that we first obtained from the ZF algebra, while the latter is more suitable for explicit calculations.

Properties of the S matrix

The two-body spin-chain S matrix has very similar properties to the two-particle QFT S matrix we investigated in chapter 3, and as before they will be useful to explicitly find it, up to the scalar factors.

Symmetries

The crucial ingredient to find the form of the S matrix is requiring that it respects the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ (or $\mathfrak{su}(1|1)_{\text{c.e.}}^4$) symmetry, *i.e.*

$$\mathcal{S}_{(12)} \mathbb{Q} = \mathbb{Q} \mathcal{S}_{(12)}. \tag{5.50}$$

This can be written for the magnon S matrix as

$$\check{\mathbb{S}}_{(12)}(p, q) \mathbb{Q}_{(12)}(p, q) = \mathbb{Q}_{(12)}(q, p) \check{\mathbb{S}}_{(12)}(p, q), \tag{5.51}$$

or equivalently

$$\mathbb{R}_{(12)}(p, q) \mathbb{Q}_{(12)}(p, q) = \mathbb{Q}_{(21)}(q, p) \mathbb{R}_{(12)}(p, q), \tag{5.52}$$

with the action of the supercharges on the two-magnon state is given by (5.49). Additionally, we will once again require left-right symmetry, *i.e.* that elements of the S matrix that differ only by relabeling $L \leftrightarrow R$ should be equal.

Braiding unitarity and physical unitarity

The concept of braiding unitarity is quite natural in the spin-chain formalism: before, the S matrix was exchanging two ZF creation operators, whence (3.66) followed by iterating the exchange twice. Here we have that the twofold exchange of two excitations should be inconsequential,

$$\mathcal{S}_{(12)} \mathcal{S}_{(12)} = \mathbb{I}, \quad (5.53)$$

where the subscript indices indicates the spaces where the matrix acts. This is depicted in figure 5.4a, and in terms of a matrix formulation reads

$$\check{\mathcal{S}}(p, q) \check{\mathcal{S}}(q, p) = \mathbb{I} = \mathbb{R}(p, q) \mathbb{R}(q, p). \quad (5.54)$$

Physical unitarity is once again just a natural reality condition on the scattering elements, which can be simply phrased in terms of a matrix representation of \mathcal{S} ,

$$\check{\mathcal{S}}^\dagger \check{\mathcal{S}} = \check{\mathcal{S}} \check{\mathcal{S}}^\dagger = \mathbb{I} = \mathbb{R}^\dagger \mathbb{R} = \mathbb{R} \mathbb{R}^\dagger. \quad (5.55)$$

Yang-Baxter equation and multiparticle scattering

We now want to extend the action of the two-body S matrix to M sites. Once again this can be done in several inequivalent ways, as illustrated in figure 5.4b in the case of $M = 3$. Requiring the equivalence of the two pictures we get again the Yang-Baxter equation

$$\mathcal{S}_{(12)} \mathcal{S}_{(23)} \mathcal{S}_{(12)} = \mathcal{S}_{(23)} \mathcal{S}_{(12)} \mathcal{S}_{(23)}. \quad (5.56)$$

We should not forget that the spin-chain S matrix will naturally feature length-changing processes, so that rewriting the above equation on the space of magnons requires some care. If we can restrict to processes where no length-changing effects arise, it is easy to rewrite the YB equation *e.g.* for $\check{\mathcal{S}}$

$$(\text{no length-changing}) \quad \check{\mathcal{S}}_{qr} \otimes \mathbb{I} \cdot \mathbb{I} \otimes \check{\mathcal{S}}_{pr} \cdot \check{\mathcal{S}}_{pq} \otimes \mathbb{I} = \mathbb{I} \otimes \check{\mathcal{S}}_{pq} \cdot \check{\mathcal{S}}_{pr} \otimes \mathbb{I} \cdot \mathbb{I} \otimes \check{\mathcal{S}}_{qr}, \quad (5.57)$$

where the subscripts indicate the dependence on the magnon momenta p, q and r . This has the same form as (3.87).

More generally, however, this is not the case. Let us assume that there exists a process where two magnons \mathcal{X}_p and \mathcal{Y}_q scatter giving $\tilde{\mathcal{Y}}_q$ and $\tilde{\mathcal{X}}_p$, and producing length-changing effects, *i.e.*

$$|\mathcal{X}_p \mathcal{Y}_q\rangle \mapsto \# |\tilde{\mathcal{Y}}_q \tilde{\mathcal{X}}_p Z^\pm\rangle. \quad (5.58)$$




Figure 5.4: Consistency conditions on the spin-chain S matrix. Unitarity (left panel) requires that acting twice with $S_{(12)}$ on a two-particle state gives back the original state. The Yang-Baxter equation (right panel) resolves the ambiguity in decomposing the scattering of three excitations. Note how, with respect to figure 3.2, this picture takes into account how S permutes the spin-chain excitations.

When we take equation (5.56) and project it on the asymptotic states, we want to write all of the vacuum sites *to the right* of the excitations. When the process (5.58) involves the two leftmost magnons, this means that the symbol Z^\pm should be commuted with the last magnon. If this rightmost magnon has momentum r , we have to account for this by writing an explicit factor of $e^{\pm ir}$ in the Yang-Baxter equation. Therefore *in presence of length-changing effects the Yang-Baxter equation is twisted* and reads

$$\mathbb{I} \otimes \check{S}_{pq} \cdot (\mathcal{F}_q \check{S}_{pr} \mathcal{F}_q^\dagger) \otimes \mathbb{I} \cdot \mathbb{I} \otimes \check{S}_{qr} = (\mathcal{F}_p \check{S}_{qr} \mathcal{F}_p^\dagger) \otimes \mathbb{I} \cdot \mathbb{I} \otimes \check{S}_{pr} \cdot (\mathcal{F}_r \check{S}_{pq} \mathcal{F}_r^\dagger) \otimes \mathbb{I}, \quad (5.59)$$

where the matrix \mathcal{F} implements a twist depending on the momentum of the rightmost magnon. Again, once the Yang-Baxter equation is established to resolve the case of $M = 3$ sites, any $M > 3$ case follows.

S-matrix elements

Using all of the properties listed above it is easy to find the whole S matrix up to two scalar factors \mathcal{S}_{pq} and $\check{\mathcal{S}}_{pq}$. Once again symmetry and unitarity requirements narrow down the solutions to two possibilities: a pure-transmission and a pure-reflection S matrix, in the sense discussed in section 3.3. As we saw there, the physical choice is the pure-transmission one. Even if the form (*i.e.* the non-vanishing elements) of such a matrix is the same as in chapter 3, the explicit expressions for the elements differ. Let us write down those expressions, splitting them in the same-chirality and opposite-chirality sectors.

Same-chirality scattering

Let us consider the scattering of two left magnons. The non-vanishing scattering processes are, as before,

$$\begin{aligned} \mathcal{S} |\phi_p^L \phi_q^L\rangle &= \mathbf{A}_{pq}^{\text{LL}} |\phi_q^L \phi_p^L\rangle, & \mathcal{S} |\phi_p^L \psi_q^L\rangle &= \mathbf{B}_{pq}^{\text{LL}} |\psi_q^L \phi_p^L\rangle + \mathbf{C}_{pq}^{\text{LL}} |\phi_q^L \psi_p^L\rangle, \\ \mathcal{S} |\psi_p^L \psi_q^L\rangle &= \mathbf{F}_{pq}^{\text{LL}} |\psi_q^L \psi_p^L\rangle, & \mathcal{S} |\psi_p^L \phi_q^L\rangle &= \mathbf{D}_{pq}^{\text{LL}} |\phi_q^L \psi_p^L\rangle + \mathbf{E}^{\text{LL}} |\psi_q^L \phi_p^L\rangle. \end{aligned} \quad (5.60)$$

The matrix elements now read

$$\begin{aligned} \mathbf{A}_{pq}^{\text{LL}} &= \frac{1}{\mathcal{S}_{pq}^{\text{LL}}} \frac{x_p^+ - x_q^-}{x_p^- - x_q^+}, & \mathbf{B}_{pq}^{\text{LL}} &= \frac{1}{\mathcal{S}_{pq}^{\text{LL}}} \frac{x_p^- - x_q^-}{x_p^- - x_q^+}, \\ \mathbf{C}_{pq}^{\text{LL}} &= \frac{e^{\frac{i}{4}(p-3q)} \frac{2i}{\hbar} \eta_p \eta_q}{\mathcal{S}_{pq}^{\text{LL}} \frac{x_p^- - x_q^+}{x_p^- - x_q^+}}, & \mathbf{D}_{pq}^{\text{LL}} &= \frac{1}{\mathcal{S}_{pq}^{\text{LL}}} \frac{x_p^+ - x_q^+}{x_p^- - x_q^+}, \\ \mathbf{E}_{pq}^{\text{LL}} &= \frac{e^{\frac{i}{4}(q-3p)} \frac{2i}{\hbar} \eta_p \eta_q}{\mathcal{S}_{pq}^{\text{LL}} \frac{x_p^- - x_q^+}{x_p^- - x_q^+}}, & \mathbf{F}_{pq}^{\text{LL}} &= \frac{-1}{\mathcal{S}_{pq}^{\text{LL}}}. \end{aligned} \quad (5.61)$$

Whether to insert the scalar factor $\mathcal{S}_{pq}^{\text{LL}}$ or its inverse is arbitrary. The present choice has the advantage that it makes the diagonal matrix element $\mathbf{A}_{pq}^{\text{LL}}$ precisely equal to *the inverse* of its string-theoretical counterpart, see eq. (3.73). Again, in the RR sector we find the same expressions with a scalar factor $\mathcal{S}_{pq}^{\text{RR}}$ and once again LR symmetry implies

$$\mathcal{S}_{pq}^{\text{LL}} = \mathcal{S}_{pq}^{\text{RR}} = \mathcal{S}_{pq}. \quad (5.62)$$

Opposite-chirality scattering

If we now consider processes of LR chirality we find

$$\begin{aligned} \mathcal{S} |\phi_p^L \phi_q^R\rangle &= \mathbf{A}_{pq}^{\text{LR}} |\phi_q^R \phi_p^L\rangle + \mathbf{B}_{pq}^{\text{LR}} |\psi_q^R \psi_p^L Z^-\rangle, & \mathcal{S} |\phi_p^L \psi_q^R\rangle &= \mathbf{C}_{pq}^{\text{LR}} |\psi_q^R \phi_p^L\rangle, \\ \mathcal{S} |\psi_p^L \psi_q^R\rangle &= \mathbf{E}_{pq}^{\text{LR}} |\psi_q^R \psi_p^L\rangle + \mathbf{F}_{pq}^{\text{LR}} |\phi_q^R \phi_p^L Z^+\rangle, & \mathcal{S} |\psi_p^L \phi_q^R\rangle &= \mathbf{D}_{pq}^{\text{LR}} |\phi_q^R \psi_p^L\rangle, \end{aligned} \quad (5.63)$$

while for RL

$$\begin{aligned} \mathcal{S} |\phi_p^R \phi_q^L\rangle &= \mathbf{A}_{pq}^{\text{RL}} |\phi_q^L \phi_p^R\rangle + \mathbf{B}_{pq}^{\text{RL}} |\psi_q^L \psi_p^R Z^-\rangle, & \mathcal{S} |\phi_p^R \psi_q^L\rangle &= \mathbf{C}_{pq}^{\text{RL}} |\psi_q^L \phi_p^R\rangle, \\ \mathcal{S} |\psi_p^R \psi_q^L\rangle &= \mathbf{E}_{pq}^{\text{RL}} |\psi_q^L \psi_p^R\rangle + \mathbf{F}_{pq}^{\text{RL}} |\phi_q^L \phi_p^R Z^+\rangle, & \mathcal{S} |\psi_p^R \phi_q^L\rangle &= \mathbf{D}_{pq}^{\text{RL}} |\phi_q^L \psi_p^R\rangle. \end{aligned} \quad (5.64)$$

Notice that we have highlighted the presence of length-changing effects. They are responsible for the presence of a twist \mathcal{F} in multi-magnon scattering events, and notably in the Yang-Baxter equation (5.56). In fact, from (5.63) we can write the explicit form of the matrix \mathcal{F}_p as

$$\mathcal{F}_p = \mathcal{U}_p \otimes \mathcal{U}_p, \quad (5.65)$$

where \mathcal{U}_p is a diagonal matrix

$$\mathcal{U}_p = \text{diag} \left(e^{\frac{i}{2}p}, 1, e^{\frac{i}{2}p}, 1 \right). \quad (5.66)$$

We then find that the twisted YB equation is satisfied, upon using the form of the matrix elements (5.61) and

$$\begin{aligned}
A_{pq}^{\text{LR}} &= \frac{e^{-iq} 1 - x_p^- x_q^+}{\mathcal{S}_{pq}^{\text{LR}} 1 - x_p^- x_q^-}, & B_{pq}^{\text{LR}} &= \frac{e^{-\frac{3i}{4}(p+q)} - \frac{2i}{h} \eta_p \eta_q}{\mathcal{S}_{pq}^{\text{LR}} 1 - x_p^- x_q^-}, \\
C_{pq}^{\text{LR}} &= \frac{e^{-i(p+q)} 1 - x_p^+ x_q^+}{\mathcal{S}_{pq}^{\text{LR}} 1 - x_p^- x_q^-}, & D_{pq}^{\text{LR}} &= \frac{1}{\mathcal{S}_{pq}^{\text{LR}}}, \\
E_{pq}^{\text{LR}} &= \frac{-e^{ip} 1 - x_p^+ x_q^-}{\mathcal{S}_{pq}^{\text{LR}} 1 - x_p^- x_q^-}, & F_{pq}^{\text{LR}} &= \frac{e^{-\frac{3i}{4}(p+q)} \frac{2i}{h} \eta_p \eta_q}{\mathcal{S}_{pq}^{\text{LR}} 1 - x_p^- x_q^-},
\end{aligned} \tag{5.67}$$

with the remaining ones following by left-right symmetry.

As before, we can use a single scalar factor for the LR and RL sectors by the definition

$$\begin{aligned}
\mathcal{S}_{pq}^{\text{LR}} &= \widetilde{\mathcal{S}}_{pq} e^{-\frac{i}{2}(p+q)} \left(\frac{1 - x_p^+ x_q^+}{1 - x_p^- x_q^-} \right)^{+1/2}, \\
\mathcal{S}_{pq}^{\text{RL}} &= \widetilde{\mathcal{S}}_{pq} e^{+\frac{i}{2}(p+q)} \left(\frac{1 - x_p^+ x_q^+}{1 - x_p^- x_q^-} \right)^{-1/2},
\end{aligned} \tag{5.68}$$

After which LR symmetry becomes manifest upon relabeling $L \leftrightarrow R$.

The full $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ -invariant S matrix

The full $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ -invariant S matrix can be found easily exploiting the tensor product structure. For instance, in the left-left sector we have

$$\begin{aligned}
\mathcal{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{++q}^L\rangle &= A_{pq}^{\text{LL}} A_{pq}^{\text{LL}} |\Phi_{++q}^L \Phi_{++p}^L\rangle, \\
\mathcal{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{--q}^L\rangle &= B_{pq}^{\text{LL}} B_{pq}^{\text{LL}} |\Phi_{--q}^L \Phi_{++p}^L\rangle + C_{pq}^{\text{LL}} C_{pq}^{\text{LL}} |\Phi_{++q}^L \Phi_{--p}^L\rangle \\
&\quad + B_{pq}^{\text{LL}} C_{pq}^{\text{LL}} (|\Phi_{-+q}^L \Phi_{+-p}^L\rangle - |\Phi_{+-q}^L \Phi_{-+p}^L\rangle),
\end{aligned} \tag{5.69}$$

where we used again the short-hand notation $|\Phi_{\pm\pm p}^L\rangle = |\Phi_{\pm\pm}^L(p)\rangle$. For mixed chiralities we have

$$\begin{aligned}
\mathcal{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{++q}^R\rangle &= A_{pq}^{\text{LR}} A_{pq}^{\text{LR}} |\Phi_{++q}^R \Phi_{++p}^L\rangle - B_{pq}^{\text{LR}} B_{pq}^{\text{LR}} |\Phi_{--q}^R \Phi_{--p}^L Z^- Z^- \rangle \\
&\quad + A_{pq}^{\text{LR}} B_{pq}^{\text{LR}} (|\Phi_{+-q}^R \Phi_{+-p}^L Z^- \rangle + |\Phi_{-+q}^R \Phi_{-+p}^L Z^- \rangle), \\
\mathcal{S}_{\mathfrak{psu}(1|1)_{\text{c.e.}}^4} |\Phi_{++p}^L \Phi_{--q}^R\rangle &= C_{pq}^{\text{LR}} C_{pq}^{\text{LR}} |\Phi_{--q}^R \Phi_{++p}^L\rangle.
\end{aligned} \tag{5.70}$$

where we wrote explicitly the length-changing effect. The remaining relations are straightforward. The S-matrix depend on the phases $\mathcal{S}^2, \widetilde{\mathcal{S}}^2$ that can be rewritten in terms of $\sigma^2, \widetilde{\sigma}^2$ by means of (4.24).

5.4 Comparing with the worksheet S matrix

The S-matrix elements we just found differ from the ones of chapter 3. Firstly, they seem to be related to the *inverse* scattering processes. This can be explained by our

ansatz (5.35) for the magnon wave functions. The choice of a negative sign in the phase factor is non-standard [127] and leads to a two-magnon scattering wave function of the form

$$\Psi(n_1, n_2) = e^{-i(pn_1 + qn_2)} + S(p, q) e^{-i(pn_2 + qn_1)}, \quad (5.71)$$

with $p > q$. We can relate this to the standard form (4.53) by flipping the sign of p and q which however reverse their order, yielding $S(q, p) \approx S(p, q)^{-1}$.

Still, the \check{S} that we found in the previous section cannot be just the inverse of the worldsheet one, as the latter satisfies Yang-Baxter equation rather than its twisted version (5.59). This discrepancy is due to the form of the spin-chain coproduct (5.49), as it was well understood already in the case of $\text{AdS}_5/\text{CFT}_4$ [58].

From the spin-chain to the worldsheet coproduct

Any coproduct can be modified by a non-local, momentum dependent change of basis in the two-excitation space [58, 137–139]. As it was shown in ref. [58], such transformations appear naturally from changes of basis of ZF algebra operators. In the case of the spin chain coproduct, let us consider the following change of basis on $\mathcal{V}_{\text{magn}}(p) \otimes \mathcal{V}_{\text{magn}}(q)$, acting on the charges as

$$\mathbb{Q}_{(12)}(p, q) \rightarrow \mathcal{W}_q^\dagger \otimes \mathbb{I} \cdot \mathbb{Q}_{(12)}(p, q) \cdot \mathcal{W}_q \otimes \mathbb{I}, \quad (5.72)$$

and, in the case of the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ spin-chain S matrix, let us take \mathcal{W}_p as in eq. (5.66). We then find a new form for the two-particle supercharges

$$\begin{aligned} \mathcal{W}_q^\dagger \otimes \mathbb{I} \cdot (\mathfrak{q}^{\text{L,R}}(p, q))_{(12)} \cdot \mathcal{W}_q \otimes \mathbb{I} &= \mathfrak{q}^{\text{L,R}}(p) \otimes \mathbb{I} e^{+\frac{i}{2}q} + \Sigma \otimes \mathfrak{q}^{\text{L,R}}(q), \\ \mathcal{W}_q^\dagger \otimes \mathbb{I} \cdot (\bar{\mathfrak{q}}^{\text{L,R}}(p, q))_{(12)} \cdot \mathcal{W}_q \otimes \mathbb{I} &= \bar{\mathfrak{q}}^{\text{L,R}}(p) \otimes \mathbb{I} e^{-\frac{i}{2}q} + \Sigma \otimes \bar{\mathfrak{q}}^{\text{L,R}}(q), \end{aligned} \quad (5.73)$$

which is now identical for L and R supercharges, but still different from the worldsheet one 3.61. The transformation (5.72) induces a change on the \mathbb{R} and \check{S} matrices,

$$\begin{aligned} \mathbb{R}(p, q) &\rightarrow \mathcal{W}_q^\dagger \otimes \mathbb{I} \cdot \mathbb{R}(p, q) \cdot \mathcal{W}_q \otimes \mathbb{I}, \\ \check{S}(p, q) &\rightarrow \mathcal{W}_p^\dagger \otimes \mathbb{I} \cdot \check{S}(p, q) \cdot \mathcal{W}_q \otimes \mathbb{I}, \end{aligned} \quad (5.74)$$

and the resulting matrices are precisely the ones that we would have found from the invariances

$$\begin{aligned} \mathbb{R}(p, q) \mathbb{Q}_{(12)}(p, q) &= \mathbb{Q}_{(21)}(q, p) \mathbb{R}(p, q), \\ \check{S}(p, q) \mathbb{Q}_{(12)}(p, q) &= \mathbb{Q}_{(12)}(q, p) \check{S}(p, q), \end{aligned} \quad (5.75)$$

had we considered the charges on the right hand side of (5.73). In fact, those transformed scattering matrices would obey the usual untwisted Yang-Baxter equation. To see this one can plug the right hand side of (5.75) in the twisted Yang-Baxter equation (5.59) and use the fact that the twist matrix \mathcal{F}_p is precisely given by (5.65).

It is worth exploring further the symmetry invariance condition in the new frame, which explicitly takes the form

$$\begin{aligned} & \mathcal{U}_q^\dagger \otimes \mathbb{I} \cdot \mathbb{R}_{pq} \cdot \mathcal{U}_q \otimes \mathbb{I} \cdot \left(\mathfrak{q}^{\text{L,R}}(p) \otimes \mathbb{I} e^{\frac{i}{2}q} + \Sigma \otimes \mathfrak{q}^{\text{L,R}}(q) \right) \\ &= \left(\mathfrak{q}^{\text{L,R}}(q) \otimes \Sigma + e^{\frac{i}{2}q} \mathbb{I} \otimes \mathfrak{q}^{\text{L,R}}(p) \right) \cdot \mathcal{U}_q^\dagger \otimes \mathbb{I} \cdot \mathbb{R}_{pq} \cdot \mathcal{U}_q \otimes \mathbb{I}, \end{aligned} \quad (5.76)$$

where we used the coproduct appearing in the right hand side of (5.73), and a similar equation holds for the conjugates $\bar{\mathfrak{q}}^{\text{L,R}}$. Using the graded permutation matrix Π^g and introducing the short-hand notation

$$\tilde{\mathbb{R}}_{pq} = \Pi^g \cdot \mathcal{U}_q^\dagger \otimes \mathbb{I} \cdot \mathbb{R}_{pq} \cdot \mathcal{U}_q \otimes \mathbb{I} \cdot \Pi^g, \quad (5.77)$$

we can finally write⁵

$$\tilde{\mathbb{R}}_{pq} \cdot \left(\mathfrak{q}^{\text{L,R}}(q) \otimes \mathbb{I} + e^{\frac{i}{2}q} \Sigma \otimes \mathfrak{q}^{\text{L,R}}(p) \right) = \left(\mathfrak{q}^{\text{L,R}}(p) \otimes \Sigma e^{\frac{i}{2}q} + \mathbb{I} \otimes \mathfrak{q}^{\text{L,R}}(q) \right) \cdot \tilde{\mathbb{R}}_{pq}. \quad (5.78)$$

Comparing this with the QFT invariance property of \mathbb{R} (4.19), we see that they coincide upon swapping the momenta $p \leftrightarrow q$. We have therefore established the relation between the worksheet and spin-chain (magnon) S matrices,

$$\mathbb{R}_{pq} \Big|_{\text{worksheet}} = \Pi^g \cdot \mathcal{U}_p^\dagger \otimes \mathbb{I} \cdot \mathbb{R}_{qp} \cdot \mathcal{U}_p \otimes \mathbb{I} \cdot \Pi^g \Big|_{\text{spin chain}}, \quad (5.79)$$

which confirms that the S matrix computed in this chapter and the one of chapter 3 describe scattering processes “inverse” to each other. It is easy to explicitly check that this identity holds using the explicit form of the S matrix elements given here and in the previous chapters.

Scalar factors and crossing

We have established that the matrix part of \mathbb{R}_{pq} can be computed equivalently in a spin-chain or worksheet picture. However, in our previous discussion we have also exploited crossing invariance of the scalar factors to constrain their form. Crossing symmetry is seemingly a purely field theoretical property, and is not obvious that anything similar can be imposed purely in a spin chain picture.⁶

An ingenious observation due to Beisert [26], see also ref. [143], shows that this is actually the case. Suppose that we can construct a two-magnon singlet state $|\mathbf{1}_{q_1 q_2}\rangle$ that is annihilated by all of the supercharges, *i.e.* in our case

$$\mathfrak{q}^{\text{L,R}} |\mathbf{1}_{q_1 q_2}\rangle = 0, \quad \bar{\mathfrak{q}}^{\text{L,R}} |\mathbf{1}_{q_1 q_2}\rangle = 0. \quad (5.80)$$

⁵We use that $(\Pi^g)^2 = \mathbb{I}$ and that for any supercharge \mathbb{Q} we have $\Pi^g \cdot \mathbb{I} \otimes \mathbb{Q} \cdot \Pi^g = \mathbb{Q} \otimes \Sigma$ and $\Pi^g \cdot \Sigma \otimes \mathbb{Q} \cdot \Pi^g = \mathbb{Q} \otimes \mathbb{I}$.

⁶However, once the spin-chain coproduct (5.49) has been established, this could be used to define a Hopf algebra structure where the notion of crossing symmetry is naturally related to the antipode operation, see appendix B in ref. [S3].




Figure 5.5: The scattering of a fundamental excitation with a singlet is trivial. We depict the singlet as composed of two excitations, one of which moves “backward” due to analytical continuation to the crossed region.

By $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ symmetry, scattering any magnon $|\mathcal{X}_p\rangle$ with it should have no consequences. On the other hand, we can think of scattering *its constituent magnons* separately with $|\mathcal{X}_p\rangle$, as illustrated in figure 5.5. This implies that the product of certain pairs of S-matrix elements should give one, when they are evaluated at momenta p, q_1 and p, q_2 . This is a constraint on the scalar factors. Moreover, since the state $|\mathbf{1}_{q_1 q_2}\rangle$ has zero momentum and energy, it must be

$$q_1 = -q_2, \quad \omega(q_1) = -\omega(q_2). \quad (5.81)$$

The only non-trivial solutions to these equations imply that one of the two momenta has been continued to the crossed region, *i.e.*

$$q_1 = q_2^c \quad \text{or} \quad q_2 = q_1^c. \quad (5.82)$$

Focusing for the moment on the latter choice, in the case of $\mathfrak{psu}(1|1)_{\text{c.e.}}^2$ we have that the singlet takes the form

$$\begin{aligned} |\mathbf{1}_{qq^c}\rangle &= \xi^{\text{LR}} |\mathbf{1}_{qq^c}^{\text{LR}}\rangle + \xi^{\text{RL}} |\mathbf{1}_{qq^c}^{\text{RL}}\rangle \\ &= \xi^{\text{LR}} \left(e^{-\frac{i}{2}q} |\phi_q^{\text{L}} \phi_{q^c}^{\text{R}}\rangle + i |\psi_q^{\text{L}} \psi_{q^c}^{\text{R}}\rangle \right) + \xi^{\text{RL}} \left(e^{-\frac{i}{2}q} |\phi_q^{\text{R}} \phi_{q^c}^{\text{L}}\rangle + i |\psi_q^{\text{R}} \psi_{q^c}^{\text{L}}\rangle \right). \end{aligned} \quad (5.83)$$

The fact that the singlet couples the L and R representations is not surprising, since in particular it should be annihilated by \mathfrak{m} . This whole discussion is more and more reminiscent of the one of crossing of section 4.1. In fact, by an argument of Arutyunov and Frolov [29] it is not hard to see that the two are equivalent. If we remove the length-changing effects we have that

$$|\mathbf{1}_{qq^c}\rangle \rightarrow \mathcal{U}_{q^c}^\dagger |\mathbf{1}_{qq^c}\rangle = \mathbb{A}^\dagger(q) \mathcal{C} \mathbb{A}^{\dagger t}(q^c), \quad (5.84)$$

where the last expression in terms of the ZF creation operation are contracted to form a scalar⁷. Then triviality of scattering with $|\mathcal{X}_p\rangle$ amounts to the statement

$$\mathbb{A}_{(1)}^\dagger(p) \left(\mathbb{A}_{(2)}^\dagger(q) \mathcal{C}_{(2)} \mathbb{A}_{(2)}^{\dagger t}(q^c) \right) = \left(\mathbb{A}_{(2)}^\dagger(q) \mathcal{C}_{(2)} \mathbb{A}_{(2)}^{\dagger t}(q^c) \right) \mathbb{A}_{(1)}^\dagger(p). \quad (5.85)$$

⁷ Recall that \mathbb{A}^\dagger is a row vector so that $\mathbb{A}^{\dagger t}$ is a column vector.

It is straightforward to use the ZF algebra relations to find

$$\begin{aligned}
& \mathbb{A}_{(1)}^\dagger(p) \mathbb{A}_{(2)}^\dagger(q) \mathcal{C}_{(2)} \mathbb{A}_{(2)}^{\dagger t}(q^c) \\
&= \mathbb{A}_{(2)}^\dagger(q) \mathbb{A}_{(1)}^\dagger(p) \mathbb{R}_{(12)}(p, q) \mathcal{C}_{(2)} \mathbb{A}_{(2)}^{\dagger t}(q^c) \\
&= \mathbb{A}_{(2)}^\dagger(q) \left(\mathbb{A}_{(1)}^\dagger(p) \mathbb{A}_{(2)}^\dagger(q^c) \mathcal{C}_{(2)} \mathbb{R}_{(12)}^{t_2}(p, q) \right)^{t_2} \\
&= \mathbb{A}_{(2)}^\dagger(q) \left(\mathbb{A}_{(2)}^\dagger(q^c) \mathbb{A}_{(1)}^\dagger(p) \mathbb{R}_{(12)}(p, q^c) \mathcal{C}_{(2)}^{t_2} \mathbb{R}_{(12)}^{t_2}(p, q) \right)^{t_2} \\
&= \mathbb{A}_{(2)}^\dagger(q) \left(\mathbb{R}_{(12)}(p, q) \mathcal{C}_{(2)} \mathbb{R}_{(12)}^{t_2}(p, q^c) \right) \mathbb{A}_{(2)}^{\dagger t}(q^c) \mathbb{A}_{(1)}^\dagger(p).
\end{aligned} \tag{5.86}$$

The expression in the last line equals the right hand side of (5.85) if

$$\mathbb{R}_{(12)}(p, q) \mathcal{C}_{(2)} \mathbb{R}_{(12)}^{t_2}(p, q^c) = \mathcal{C}_{(2)}. \tag{5.87}$$

This is precisely equivalent⁸ to the crossing equations (4.23). If we instead chose to set $q_1 = q_2^c$ we can derive the crossing equation in the first variable instead. In that case we have

$$\begin{aligned}
|\mathbf{1}_{q^c q}\rangle &= \xi^{\text{LR}} |\mathbf{1}_{q^c q}^{\text{LR}}\rangle + \xi^{\text{RL}} |\mathbf{1}_{q^c q}^{\text{RL}}\rangle \\
&= \xi^{\text{LR}} \left(e^{\frac{i}{2}q} |\phi_{q^c}^{\text{L}} \phi_q^{\text{R}}\rangle - i |\psi_{q^c}^{\text{L}} \psi_q^{\text{R}}\rangle \right) + \xi^{\text{RL}} \left(e^{\frac{i}{2}q} |\phi_{q^c}^{\text{R}} \phi_q^{\text{L}}\rangle - i |\psi_{q^c}^{\text{R}} \psi_q^{\text{L}}\rangle \right),
\end{aligned} \tag{5.88}$$

which we can rewrite as

$$|\mathbf{1}_{q^c q}\rangle \rightarrow \mathcal{U}_q^\dagger |\mathbf{1}_{q^c q}\rangle = \mathbb{A}^\dagger(q^c) \mathcal{C}^\dagger \mathbb{A}^{\dagger t}(q), \tag{5.89}$$

which leads to a similar discussion as above and reproduces (4.22).

Finally, to further confirm our derivation, it is easy to check explicitly that imposing

$$\mathcal{S}_{(23)} \mathcal{S}_{(12)} |\mathcal{X}_p \mathbf{1}_{q^c q}\rangle = |\mathbf{1}_{q^c q} \mathcal{X}_p\rangle, \quad \mathcal{S}_{(23)} \mathcal{S}_{(12)} |\mathcal{X}_p \mathbf{1}_{q^c q}\rangle = |\mathbf{1}_{q^c q} \mathcal{X}_p\rangle, \tag{5.90}$$

results precisely in the equations (4.26) for the dressing factors. As always, all of this can be straightforwardly extended to the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ S matrix, see also ref. [S3].

5.5 Chapter summary

In this chapter we have seen how the all-loop S matrix and dispersion relation of chapter 3 can be equivalently found from a spin-chain picture.

⁸ It may appear strange that, while matching the S matrices also required exchanging $p \leftrightarrow q$ and involved graded permutations, none of this is necessary to reproduce the crossing equation. This is simply due to the fact that crossing invariance must hold (or, can be imposed) for \mathbb{R} , its inverse, and its graded permutation.

Our derivation built on the weakly-coupled spin chain description of ref. [112]. For $\hbar \ll 1$, we can think of *two* spin chains, one containing left-moving excitations transforming under $\mathfrak{psu}(1, 1|2)_L$, and the other containing right-moving ones transforming under $\mathfrak{psu}(1, 1|2)_R$, as in figure 5.3. We have then shown how from the $\mathfrak{psu}(1, 1|2)^2$ representations it is possible to construct a $\frac{1}{2}$ -supersymmetric vacuum, and excitations above it that transform under a $\mathfrak{psu}(1|1)^4$ residual symmetry. In fact, this weak-coupling symmetry is precisely equal to the on-shell symmetry of worldsheet excitations.

Moreover, we have extended our construction to arbitrary values of the coupling. This results in the excitations of a given chirality being charged under both left and right supercharges, together with a central extension of the residual symmetries and a *dynamical* spin chain, where the number of sites can fluctuate. Using these symmetries, supplemented by a discrete left-right one, we have been able to determine the two-magnon S matrix, which in fact satisfies an analogue of the Yang-Baxter equation and can be used to construct many-magnon S matrices in a consistent way.

The construction, including the form of the $\mathfrak{psu}(1, 1)_{c.e.}^4$ symmetry, was strongly reminiscent of what we did in the worldsheet theory in chapter 3. Still, an apparent difference was in the form of the coproduct (5.49) that defines the action of the symmetries on several spin-chain sites. We have shown explicitly that, up to a change of the two-particle basis, such a difference can be reabsorbed and in fact the worldsheet and spin-chain S matrices are equivalent (5.79). This was expected based on what happens in $\text{AdS}_5/\text{CFT}_4$, see ref. [58]. The final ingredient to put the spin-chain and worldsheet pictures on the same footing was crossing symmetry, which would appear to be a genuinely field-theoretical feature. We show that an equivalent notion holds for our spin-chain, similarly to what was argued by Beisert in ref. [26].

Let us conclude by mentioning that the notations used here for the spin-chain symmetry algebra and S matrix differ from the ones of our original works [S3, S5]. The present choice was meant to further emphasise the similarities with the worldsheet analysis, in particular by taking the very same form of the supercharges on the one-magnon representation as it was for the one-particle representation in chapter 3. Relating this choice with [S3, S5] amounts to another change of the two-particle basis, see also ref. [58].

6 The all-loop Bethe ansatz equations

Once the two-body S matrix and the dispersion relation are known, it only remains to impose that the spatial dimension of the worldsheet, or equivalently the spin chain, are given by a circle of finite length ℓ . This results in the Bethe-Yang equations for the QFT, or in the asymptotic¹ Bethe ansatz equations (BAE) for the spin chain. As we will argue, the equations are equivalent in the two frameworks, so that the two theories will have the same spectrum of momenta and, given that the dispersion relation is the same, of energy. There are several ways of deriving the BAE. Here we will focus on what is perhaps the most intuitive way from the physical point of view, that is the coordinate Bethe ansatz. We will work it out for both the spin-chain and the worldsheet picture.

6.1 Bethe ansatz essentials

Before working out the BAE for S matrix that we found in the previous chapters, let us illustrate the idea behind the Bethe ansatz on the simplest possible example, and postpone the more complicated cases to the next sections. We consider here the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ S matrix and *restrict to a single type of excitation, e.g. $\mathcal{X} = \phi^L$* . Even if this truncation violates crossing symmetry, it is consistent from the scattering point of view, since ϕ_p^L scatters diagonally with itself. Then, the S matrix reduces to a number.

Imposing periodicity

The spin-chain picture is perhaps the easiest to visualise. Let us therefore start by considering the M -magnon asymptotic wave function for an integrable theory. Consider an asymptotic state

$$|\mathcal{X}_{p_1} \cdots \mathcal{X}_{p_M}\rangle = \sum_{n_1 \ll \cdots \ll n_M} e^{-i(p_1 n_1 + \cdots + p_M n_M)} |Z \cdots Z \mathcal{X}_{(n_1)} Z \cdots Z \mathcal{X}_{(n_M)} Z \cdots\rangle, \quad (6.1)$$

¹As we mentioned in the introduction, these equations are asymptotic because they ignore wrapping effects. With this *caveat* in mind, we will often interchangeably refer to (asymptotic) Bethe ansatz and Bethe-Yang equations when comparing the spin-chain and worldsheet approaches.




Figure 6.1: Illustrations of the periodicity condition leading to the Bethe ansatz equations. Above: the two ends of the chain have been identified, so that the first excitation \mathcal{X}_{p_1} in position n_1 can be thought as sitting at the end of the chain, in position $n_1 + \ell$. This can be realised by scattering \mathcal{X}_{p_1} through all of the \mathcal{X}_{p_j} . Below: equivalently, the wave-function should be ℓ -periodic as a function on \mathbb{R} . The box identifies the periodicity condition evaluated as in (6.5).

with $p_1 > \dots > p_M$, which is a natural generalisation of the two-magnon state (5.36). By the same reasoning that gave us factorised scattering in chapter 3, it is clear now that the magnons will undergo pairwise scattering along the one-dimensional chain. For instance, if the first and second magnon scatter, we will get to an ansatz where the magnon with momentum p_1 is in the region indexed by n_2 , and *vice versa*. In general, the multimagnon wave function will be a combination

$$|\Psi(p_1, \dots, p_M)\rangle = \sum_{\pi \in S_M} \chi(p_{\pi(1)}, \dots, p_{\pi(M)}) |\mathcal{X}_{p_{\pi(1)}} \cdots \mathcal{X}_{p_{\pi(M)}}\rangle, \quad (6.2)$$

where $\pi \in S_M$ is a permutation. This wave function is still asymptotic, in the sense that it does not explicitly depend on regions where two magnons come close, *i.e.* where scattering happens. Interactions are encoded in the coefficients χ , which will involve S-matrix elements. Comparing with the two-magnon case (5.36) and (5.71) it is immediate to see that if the permutation involves only two indices, it will be

$$\chi(p_1, \dots, p_{j+1}, p_j, \dots, p_M) = S(p_j, p_{j+1}) \chi(p_1, \dots, p_j, p_{j+1}, \dots, p_M). \quad (6.3)$$

If we now take into account factorised scattering, we can immediately extend this to an *arbitrary* permutation, so that we can rewrite

$$\begin{aligned} |\Psi(p_1, \dots, p_M)\rangle &= \sum_{\pi \in S_M} \mathcal{S}_\pi |\mathcal{X}_{p_1} \cdots \mathcal{X}_{p_M}\rangle, \\ \mathcal{S}_\pi |\mathcal{X}_{p_1} \cdots \mathcal{X}_{p_M}\rangle &= \prod_{(j,k) \in \pi} S(p_j, p_k) |\mathcal{X}_{p_{\pi(1)}} \cdots \mathcal{X}_{p_{\pi(M)}}\rangle, \end{aligned} \quad (6.4)$$

where we set $\chi(p_1, \dots, p_M) = 1$.

We now will impose that this wave function is ℓ -periodic. Let us shift e.g. the first coordinate n_1 by ℓ and bring it all the way around the chain:

$$(n_1, n_2, \dots, n_{M-1}, n_M) \rightarrow (n_2, n_3, \dots, n_M, n_1 + \ell), \quad (6.5)$$

as depicted in figure 6.1. Periodicity is the statement that $|\Psi\rangle$ is invariant under such a transformation. This is not granted by the ansatz (6.2). Using (6.3) we in fact have that shifting n_1 by ℓ gives the periodicity condition²

$$|\Psi(p_1, \dots, p_M)\rangle = \left(e^{-ip_1\ell} \prod_{j=2}^M S(p_1, p_j) \right) |\Psi(p_1, \dots, p_M)\rangle. \quad (6.6)$$

We then have to require the expression in the big brackets to be equal to 1. Physically, this is a quantisation condition for the momentum p_1 . There are several such conditions: suppose that we repeat the shift (6.5) for n_2 , then n_3 , until n_k . We then get to the configuration

$$(n_k, n_{k+1}, \dots, n_M, n_1 + \ell, \dots, n_{k-1} + \ell, n_k + \ell). \quad (6.7)$$

Imposing periodicity for any $k = 1, \dots, M$ results in k coupled equations for the magnon momenta

$$\text{spin chain: } e^{-ip_k\ell} \prod_{j \neq k}^M S(p_k, p_j) = 1, \quad k = 1, \dots, M. \quad (6.8)$$

These are the celebrated *Bethe ansatz equations*. More precisely, the above derivation is referred to as the *coordinate* Bethe ansatz.

Let us remark that in our derivation we assumed that \mathcal{X}_p has a single flavour, so that the scattering is diagonal and the S matrix $S(p, q)$ is just a number. Extending the discussion above to the case of several flavours that scatter diagonally, *i.e.* by pure transmission of *all* quantum numbers is completely straightforward—it only requires adding flavour indices to \mathcal{X}_p and $S(p, q)$. Our S matrix is not so simple, because the fermion number is not always transmitted, for instance in the process

$$\mathcal{S} |\phi_p^L \psi_q^L\rangle = \mathbf{B}_{pq}^{\text{LL}} |\psi_q^L \phi_p^L\rangle + \mathbf{C}_{pq}^{\text{LL}} |\phi_q^L \psi_p^L\rangle. \quad (6.9)$$

In order to impose periodicity we will first have to diagonalise the action of the S matrix by the so-called nested Bethe ansatz. This is a technical complication that we will address in the next section in the simple case of the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ S matrix.

²One may find the notation of eq. (6.6) confusing, since quantum-mechanical states are defined up to an overall phase. However here we normalised $\chi(p_1, \dots, p_M) = 1$. Eq. (6.6) should be then understood as a condition of spatial periodicity on the wave-function.

Bethe ansatz from the worldsheet

It is easy to see that the reasoning above can be repeated almost *verbatim* from the point of view of the ZF algebra and S matrix. In that case, the asymptotic wave function would be

$$|\Psi(p_1, \dots, p_M)\rangle = \sum_{\pi \in S_M} \chi(p_{\pi(1)}, \dots, p_{\pi(M)}) |p_{\pi(1)}, \dots, p_{\pi(M)}\rangle, \quad (6.10)$$

where each asymptotic state would read, in position space,

$$|p_1, \dots, p_M\rangle = \int_{\sigma_1 \ll \dots \ll \sigma_M} d\sigma_1 \dots d\sigma_M e^{i(p_1\sigma_1 + \dots + p_M\sigma_M)} \mathbb{A}^\dagger(\sigma_1) \dots \mathbb{A}^\dagger(\sigma_M) |0\rangle. \quad (6.11)$$

Note that we kept an ordering such that the particle of momentum p_1 is the leftmost one. We can now require periodicity under shifts of $\sigma_j \rightarrow \sigma_j + \ell$, obtaining the Bethe ansatz

$$\text{worldsheet: } e^{ip_k\ell} \prod_{j \neq k}^M S(p_k, p_j) = 1, \quad k = 1, \dots, M. \quad (6.12)$$

The different sign in the plane-wave coefficient resulted in different quantisation conditions. As discussed in the previous chapter, that same sign choice makes the spin chain S matrix be the inverse of the worldsheet one. Indeed, equations (6.12) and (6.8) are related by inverting $S(p, q)$.

The level-matching condition

In chapter 2 we have found that in string theory the only physical states are the ones satisfying the constraint

$$\mathbb{P} |\text{physical}\rangle = 0, \quad (6.13)$$

in the case of zero winding. This results in a condition on the momenta of the M -particle state

$$p_1 + \dots + p_M = 0. \quad (6.14)$$

If we allow for winding it is not hard to see that the right hand side of the constraint should not vanish, but rather equal $2\pi W$, where the integer W is the winding number.

This string-theoretical requirement has an interesting equivalent in the spin-chain picture. Consider an M -magnon asymptotic state in the spin chain, and imagine of shifting each magnon *e.g.* to the left by one site, as in figure 6.2. On the one hand, cyclicity of the chain requires this transformation to leave it invariant. On the other hand, by eq. (5.37) this results in M phase shifts of the form e^{-ip_j} . We then have an additional requirement

$$e^{i(p_1 + \dots + p_M)} = 1. \quad (6.15)$$




Figure 6.2: By shifting each magnon to the right by one site, we produce a shift of $e^{-i(p_1+\dots+p_M)}$. On the other hand, for a cyclic chain this produces no effect, so that consistency dictates the level-matching condition (6.15).

We then find that the spin chain and worldsheet pictures are completely equivalent, *mutatis mutandis*, from the Bethe ansatz point of view. It is then just a matter of convenience to derive our equations with reference to one or the other framework.

6.2 Nested Bethe ansatz

In this subsection we will illustrate how the nesting procedure works for the simple case of the $\mathfrak{su}(1|1)_{c.e.}^2$ S matrix. In what follows, for definiteness, we will work *from the spin-chain point of view*, and only at the end of the day comment on the worldsheet picture.

The strategy will be to split the action of the S matrix in several steps: first, consider a set of “level-I excitations” on the usual vacuum, that have the property of scattering diagonally among themselves. For these excitations, the coordinate Bethe ansatz can be straightforwardly implemented as we illustrated in the previous section. In order to incorporate the remaining excitations, we construct a new “level-II” vacuum. This is a state consisting of level-I excitations only. We can now consider level-II excitations on this vacuum that scatter trivially among themselves. If needed, we can use those to construct a level-III vacuum, and so on. In this way, level by level, the scattering is very simple. Of course, we will have to require that *e.g.* the level-II excitations propagate on the level-II vacuum in a way that is compatible with the dynamics of the fundamental S matrix. This will ensure that different levels are “glued” in a consistent way.

Level-I excitations

For definiteness, here we work with the spin-chain S matrix. Looking at the scattering elements in section 5.3, we see that there are several non-diagonal scattering processes. The level-I vacuum is the usual one,

$$|0\rangle^I = |Z^\ell\rangle. \quad (6.16)$$

We have two possible choices of processes that scatter diagonally among themselves:

$$V_A^I = \{\phi^L, \psi^R\}, \quad V_B^I = \{\phi^R, \psi^L\}. \quad (6.17)$$

We can think of this as a choice of the highest weight states in the left and right representations, or equivalently of the grading of the superalgebra. Clearly the choice A corresponds to having as lowering operators q^L and \bar{q}^R , while B corresponds to picking \bar{q}^L and q^R . Let us for the moment pick the choice A .

A level-I state is a collection of excitations of V_A^I , *e.g.*

$$|\mathcal{X}_{p_1}^{\alpha_1}, \dots, \mathcal{X}_{p_M}^{\alpha_M}\rangle^I = |ZZ\mathcal{X}_{p_1}^{\alpha_1}ZZ \dots ZZ\mathcal{X}_{p_M}^{\alpha_M}ZZ \dots\rangle, \quad (6.18)$$

like in (6.1). There will be M_L^I left excitations and M_R^I right ones, distinguished by the flavour label α , so that

$$M = M_L^I + M_R^I. \quad (6.19)$$

Out of such states we can as before construct the asymptotic level-one wave function by acting with the multiparticle S matrix \mathcal{S}_π

$$|\Psi(p_1, \dots, p_M)\rangle_{\alpha_1, \dots, \alpha_M}^I = \sum_{\pi \in S_M} \mathcal{S}_\pi |\mathcal{X}_{p_1}^{\alpha_1}, \dots, \mathcal{X}_{p_M}^{\alpha_M}\rangle^I. \quad (6.20)$$

Now also the S matrix will carry flavour indices α that can take values L and R. Still, by construction \mathbb{S}_π acts diagonally on the level-one states,

$$\mathcal{S}_\pi |\mathcal{X}_{p_1}^{\alpha_1}, \dots, \mathcal{X}_{p_M}^{\alpha_M}\rangle^I = S_\pi^I(p_1, \dots, p_M) |\mathcal{X}_{p_{\pi(1)}}^{\alpha_{\pi(1)}}, \dots, \mathcal{X}_{p_{\pi(M)}}^{\alpha_{\pi(M)}}\rangle^I, \quad (6.21)$$

where the phase S_π factorises

$$S_\pi(p_1, \dots, p_M) = \prod_{(j,k) \in \pi} S_{\alpha_j \alpha_k}^{I,I}(p_j, p_k). \quad (6.22)$$

The level-one S-matrix elements $S_{\alpha_j \alpha_k}^{I,I}$ can be immediately read off the fundamental S matrix elements of section 5.3,

$$S_{LL}^{I,I} = A^{LL}, \quad S_{RR}^{I,I} = F^{RR}, \quad S_{LR}^{I,I} = C^{LR}, \quad S_{RL}^{I,I} = D^{RL}. \quad (6.23)$$

Level-II excitations

The level-II vacuum is a collection of level-I excitations

$$|0\rangle^{II} = |\mathcal{X}_{p_1}^{\alpha_1} \dots \mathcal{X}_{p_M}^{\alpha_M}\rangle. \quad (6.24)$$

Level-II excitations can be constructed by acting on $|0\rangle^{II}$ with the lowering operators. Acting by q^L turns ϕ^L and ψ^R into ψ^L and ϕ^R respectively³. However we want to treat $|0\rangle^{II}$ as a vacuum, so that the scattering properties its the level-II excitations should be blind

³Since we are working in the spin chain picture, we should not forget to insert the appropriate length-changing effects.

to the underlying level-I excitations. The effect of q^R would be similar, up to exchanging the left and right modules.

The generic form of a level-II state containing a single excitation is

$$\begin{aligned} |\mathcal{Y}_y^L\rangle^{\text{II}} &= \sum_{k=1}^M \chi_k(y^L) |\mathcal{X}_{p_1}^{\alpha_1} \cdots \mathcal{Y}_y^L \cdots \mathcal{X}_{p_M}^{\alpha_M}\rangle, \\ \chi_k(y^L) &= f(y^L, p_k) \prod_{j=1}^{k-1} S_{\alpha_k \alpha_j}^{\text{II,I}}(y^L, p_j), \end{aligned} \quad (6.25)$$

where for brevity we wrote $\mathcal{Y}^L = q^L \mathcal{X}_{p_k}^{\alpha_k}$. The superscript L in \mathcal{Y}^L is meant to remind us that we are acting on the vacuum by a left lowering operator. The coefficient $f(y^L, p_k)$ stands for the creation of a level-II excitation on top of a level-I one which had momentum p_k , while $S_{\alpha_k \alpha_j}^{\text{II,I}}(y^L, p_j)$ accounts for the scattering of the level-II excitation with level-I ones, that is needed to position the former at site k . These functions cannot be arbitrary if we want this description to be compatible with the one at level I. To this end, we require that creating a level-II excitation and scattering the underlying level-I excitation are commuting operations:

$$\mathcal{S}_\pi |y^L\rangle^{\text{II}} = S_\pi^{\text{I}}(p_1, \dots, p_M) |y^L\rangle_\pi^{\text{II}}, \quad (6.26)$$

where $|y^L\rangle_\pi^{\text{II}}$ is the level-II state constructed on the permuted vacuum,

$$\begin{aligned} |y^L\rangle_\pi^{\text{II}} &= \sum_{k=1}^M \chi_{k,\pi}(y^L) |\mathcal{X}_{p_{\pi(1)}}^{\alpha_{\pi(1)}} \cdots \mathcal{Y}_y^L \cdots \mathcal{X}_{p_{\pi(M)}}^{\alpha_{\pi(M)}}\rangle, \\ \chi_{k,\pi}(y^L) &= f(y^L, p_{\pi(k)}) \prod_{j=1}^{k-1} S_{\alpha_{\pi(k)} \alpha_{\pi(j)}}^{\text{II,I}}(y^L, p_{\pi(j)}). \end{aligned} \quad (6.27)$$

On the other hand, $\mathcal{S}_\pi |y\rangle^{\text{II}}$ can be computed just by acting on the excitations with the S matrix of section 5.3, regardless of whether they are in the first or second level.

Level-II excitations: propagation

To be more specific, let us consider a level-II vacuum consisting of two ϕ^L bosons, and one level-II excitation on it,

$$|y^L\rangle_{\text{LL}}^{\text{II}} = f_L(y^L, p) |\psi_p^L \phi_q^L\rangle + f_L(y^L, q) S_{\text{LL}}^{\text{II,I}}(y^L, p) |\phi_p^L \psi_q^L\rangle. \quad (6.28)$$

The generalisation to a longer vacuum will be straightforward owing to the factorisation property. Note how both of the expressions on the right hand side of the non-diagonal scattering processes appear in the ansatz. After scattering, the compatibility condition (6.26) mandates that this should be proportional to

$$|y\rangle_{\text{LL},\pi}^{\text{II}} = f_L(y^L, q) |\psi_q^L \phi_p^L\rangle + f_L(y^L, p) S_{\text{LL}}^{\text{II,I}}(y^L, q) |\phi_q^L \psi_p^L\rangle, \quad (6.29)$$

with a proportionality constant equal to $S_{LL}^{I,I}(p, q) = A_{pq}^{LL}$. On the other hand, we can act with \mathcal{S}_{pq} on $|\psi_p^L \phi_q^L\rangle$ and $|\phi_p^L \psi_q^L\rangle$, as in (5.60). The resulting expression can be proportional to $|y^L\rangle_{LL,\pi}^{II}$ and in fact equate $A_{pq}^{LL} |y\rangle_{LL,\pi}^{II}$, provided that the level-II coefficients satisfy

$$\begin{aligned} S_{LL}^{II,I}(y^L, q) f_L(y^L, p) A_{pq}^{LL} &= f_L(y^L, p) D_{pq}^{LL} + f_L(y^L, q) S_{LL}^{II,I}(y^L, p) C_{pq}^{LL}, \\ f_L(y^L, q) A_{pq}^{LL} &= f_L(y^L, p) E_{pq}^{LL} + f_L(y^L, q) S_{LL}^{II,I}(y^L, p) B_{pq}^{LL}. \end{aligned} \quad (6.30)$$

Using the explicit form of the S-matrix elements (5.61), we find that it must be

$$f_L(y^L, p) = g_L(y^L) \frac{\eta_p e^{\frac{i}{4}p}}{k_L(y^L) - x_p^-}, \quad S_{LL}^{II,I}(y^L, p) = \frac{k_L(y^L) - x_p^+}{k_L(y^L) - x_p^-}, \quad (6.31)$$

where $g_L(y)$ and $k_L(y)$ are arbitrary functions of y . Note that at this nesting level none of these functions can depend on the dressing factors.

A similar calculation in the right sector gives the ansatz

$$\begin{aligned} |y^L\rangle_{RR}^{II} &= f_R(y^L, p) |\phi_p^R Z^+ \psi_q^R\rangle - f_R(y^L, q) S_{RR}^{II,I}(y^L, p) |\psi_p^R \phi_q^R Z^+\rangle, \\ |y^L\rangle_{RR,\pi}^{II} &= f_R(y^L, q) |\phi_q^R Z^+ \psi_p^R\rangle - f_R(y^L, p) S_{RR}^{II,I}(y^L, q) |\psi_q^R \phi_p^R Z^+\rangle, \end{aligned} \quad (6.32)$$

where now ψ^R is the level-I excitation, and length-changing effects appear. The minus signs take into account the fact that the underlying vacuum is fermionic. With this choice it will be easier to impose the periodicity condition at the end of the day, since all fermion signs will be accounted for. The consistency condition now results in the equations

$$\begin{aligned} S_{RR}^{II,I}(y^L, q) f_R(y^L, p) F_{pq}^{RR} &= -f_R(y^L, p) e^{iq} C_{pq}^{RR} + f_R(y^L, q) S_{RR}^{II,I}(y, p) D_{pq}^{RR}, \\ f_R(y^L, q) e^{ip} F_{pq}^{RR} &= +f_R(y^L, p) e^{iq} B_{pq}^{RR} - f_R(y^L, q) S_{RR}^{II,I}(y, p) E_{pq}^{RR}. \end{aligned} \quad (6.33)$$

which are solved by

$$f_R(y^L, p) = -i \frac{g_R}{x_p^-}(y) \frac{\eta_p e^{-\frac{3}{4}ip}}{1 - \frac{1}{k_R(y^L) x_p^+}}, \quad S_{RR}^{II,I}(y^L, p) = \frac{1 - \frac{1}{k_R(y^L) x_p^-}}{1 - \frac{1}{k_R(y^L) x_p^+}}. \quad (6.34)$$

There is one last consistency condition the we should impose, which is the one arising when the vacuum contains states of the left *and* right modules. For two states, this gives a level-II excitation

$$\begin{aligned} |y^L\rangle_{LR}^{II} &= +f_L(y^L, p) |\psi_p^L \psi_q^R\rangle + f_R(y^L, q) S_{RL}^{II,I}(y^L, p) |\phi_p^L \phi_q^R Z^+\rangle, \\ |y^L\rangle_{LR,\pi}^{II} &= -f_R(y^L, q) |\psi_q^R \psi_p^L\rangle + f_L(y^L, p) S_{LR}^{II,I}(y^L, q) |\phi_q^R Z^+ \phi_p^L\rangle, \end{aligned} \quad (6.35)$$

when we take a left and right state in this order. This results in the equations

$$\begin{aligned} S_{LR}^{II,I}(y^L, q) f_L(y^L, p) e^{ip} C_{pq}^{LR} &= f_L(y^L, p) E_{pq}^{LR} + f_R(y^L, q) S_{RL}^{II,I}(y^L, p) B_{pq}^{LR}, \\ -f_R(y^L, q) C_{pq}^{LR} &= f_L(y^L, p) F_{pq}^{LR} + f_R(y^L, q) S_{RL}^{II,I}(y^L, p) A_{pq}^{LR}. \end{aligned} \quad (6.36)$$

These conditions can be solved by imposing

$$\begin{aligned} k_L(y^L) &= k_R(y^L) = y^L, & g_L(y^L) &= -i y g_R(y^L), \\ S_{RL}^{II,I}(y^L, p) &= S_{LL}^{II,I}(y^L, p), & S_{LR}^{II,I}(y^L, p) &= S_{RR}^{II,I}(y^L, p). \end{aligned} \quad (6.37)$$

The two last equations can be interpreted by saying that when we act with the lowering operator \mathfrak{q}^L in the plane-wave ansatz (6.25) we create a well-defined excitation. Its scattering does not depend on the underlying vacuum that it is created out of, but only on what it scatters with. This confirms that the lowering operators are blind to the constituents of $|0\rangle^{\text{II}}$.

So far we have exclusively considered excitations created by \mathfrak{q}^L . We can create different level-II excitations by acting with $\bar{\mathfrak{q}}^R$ instead. We will denote the corresponding one-particle excitation by $|y_R\rangle^{\text{II}}$. The computations follow the same pattern, up to appropriately taking into account the form of the S-matrix elements and the length-changing effects. We have, in particular

$$\begin{aligned} S_{LL}^{II,I}(y^R, p) &= \frac{1 - \frac{1}{y^R x_p^+}}{1 - \frac{1}{y^R x_p^-}}, & S_{RR}^{II,I}(y^R, p) &= \frac{y^R - x_p^-}{y^R - x_p^+}, \\ S_{RL}^{II,I}(y^R, p) &= S_{LL}^{II,I}(y^R, p), & S_{LR}^{II,I}(y^R, p) &= S_{RR}^{II,I}(y^R, p), \end{aligned} \quad (6.38)$$

where as usual fermion signs have been accounted for.

Level-II excitations: scattering

The preceding discussion covers all of the dynamics involving a single level-II excitation on an arbitrary vacuum, owing to factorisation of scattering. However, as soon as we include another excitation, a new dynamics arises: scattering of two level-II excitations.

A state containing two level-II excitations obtained by acting with \mathfrak{q}^L takes the form

$$|y_1^L, y_2^L\rangle^{\text{II}} = \sum_{k < l}^M \chi_k(y_1^L) \chi_l(y_2^L) |\mathcal{X}_{p_1}^{\alpha_1} \cdots \mathcal{Y}_{y_1}^L \cdots \mathcal{Y}_{y_2}^L \cdots \mathcal{X}_{p_M}^{\alpha_M}\rangle, \quad (6.39)$$

where $\mathcal{Y}_{y_1}^L$ sits at position k and $\mathcal{Y}_{y_2}^L$ at position l . Once again, this ansatz is subject to a condition similar to (6.26), reading

$$\check{S}_\pi |y_1^L, y_2^L\rangle^{\text{II}} = S_\pi^I(p_1, \dots, p_M) |y_1^L, y_2^L\rangle_\pi^{\text{II}}, \quad (6.40)$$

Clearly the only problems may arise from the scattering of y_1^L and y_2^L . Let us consider the case where we have two left-excitations on a length-two vacuum. We make an ansatz

$$\begin{aligned} |y_1^L y_2^L\rangle_{LL}^{\text{II}} &= f_L(y_1^L, p) f_L(y_2^L, q) S_{LL}^{II,I}(y_2^L, p) |\psi_p^L \psi_q^L\rangle \\ &\quad - f_L(y_2^L, p) f_L(y_1^L, q) S_{LL}^{II,I}(y_1^L, p) S_{LL}^{II,II}(y_1^L, y_2^L) |\psi_p^L \psi_q^L\rangle, \\ |y_1^L y_2^L\rangle_{LL, \pi}^{\text{II}} &= f_L(y_1^L, q) f_L(y_2^L, p) S_{LL}^{II,I}(y_2^L, q) |\psi_q^L \psi_p^L\rangle \\ &\quad - f_L(y_2^L, q) f_L(y_1^L, p) S_{LL}^{II,I}(y_1^L, q) S_{LL}^{II,II}(y_1^L, y_2^L) |\psi_q^L \psi_p^L\rangle, \end{aligned} \quad (6.41)$$

where again we accounted for the minus sign. The only new ingredient is the undetermined factor $S_{LL}^{\text{II,II}}(y_1^L, y_2^L)$ that represents the scattering of the two level-II excitations. It is easy to see that the consistency condition is solved by

$$S_{LL}^{\text{II,II}}(y_1^L, y_2^L) = 1, \quad (6.42)$$

i.e. the scattering is trivial. In fact, it is immediate to check that all of the level-II scattering matrices are trivial, including the ones scattering $|y^R\rangle$ with $|y^L\rangle$ or with itself. Using factorisation of scattering, all our considerations extend to any level-II state with M_L^{II} left and M_R^{II} right excitations.

Since we have taken into account all of the fundamental excitations and diagonalised all scattering processes, the nesting procedure terminates here. Had we to include additional non-diagonal processes, it would be necessary to construct a level-III vacuum, and so on.

Bethe ansatz equations

The Bethe ansatz equations arise now out of imposing periodicity for each set of excitations. Looking at (6.40) we see that we chose an ordering of the level-II magnons. As discussed in the previous section, the description should be unchanged if we rearrange them using the fact that the chain is periodic. It is easy to see that this gives the conditions

$$\begin{aligned} 1 &= \prod_{j=1}^{M_L^{\text{I}}} S_{LL}^{\text{II,I}}(y_k^L, p_j) \prod_{j=1}^{M_R^{\text{I}}} S_{RR}^{\text{II,I}}(y_k^L, p_j), \quad k = 1, \dots, M_L^{\text{II}}, \\ 1 &= \prod_{j=1}^{M_L^{\text{I}}} S_{LL}^{\text{II,I}}(y_k^R, p_j) \prod_{j=1}^{M_R^{\text{I}}} S_{RR}^{\text{II,I}}(y_k^R, p_j), \quad k = 1, \dots, M_R^{\text{II}}, \end{aligned} \quad (6.43)$$

where we used that $S^{\text{II,II}} = 1$. Note that we do not have to take into account any additional fermion sign, due to our convention. One pictorial way to interpret this equation is to take one level-II excitation and carry it along the chain: by periodicity, the phase arising from the whole scattering sequence should be one.

Fundamental excitations obey a slightly modified version of (6.8), which now includes also higher level excitations:

$$\begin{aligned} e^{ipL} &= \prod_{j \neq k}^{M_L^{\text{I}}} S_{LL}^{\text{I,I}}(p_k^L, p_j^L) \prod_{j=1}^{M_R^{\text{I}}} S_{LR}^{\text{I,I}}(p_k^L, p_j^R) \prod_{j=1}^{M_L^{\text{II}}} S_{LL}^{\text{I,II}}(p_k^L, y_j^L) \prod_{j=1}^{M_R^{\text{II}}} S_{LR}^{\text{I,II}}(p_k^L, y_j^R), \\ e^{ipL} &= \prod_{j \neq k}^{M_R^{\text{I}}} S_{RR}^{\text{I,I}}(p_k^R, p_j^R) \prod_{j=1}^{M_L^{\text{I}}} S_{RL}^{\text{I,I}}(p_k^R, p_j^L) \prod_{j=1}^{M_R^{\text{II}}} S_{RR}^{\text{I,II}}(p_k^R, y_j^R) \prod_{j=1}^{M_L^{\text{II}}} S_{RL}^{\text{I,II}}(p_k^R, y_j^L), \end{aligned} \quad (6.44)$$

for $k = 1, \dots, M_{L,R}^{\text{I}}$. Note that we have introduced explicit labels for the momenta of left and right excitations. For brevity, in terms of the Zukhovski variables, we set

$$x_j^\pm = x_j^\pm(p^L), \quad \tilde{x}_j^\pm = x_j^\pm(p^R), \quad y = y^L, \quad \tilde{y} = y^R. \quad (6.45)$$

The we can write the full set of Bethe equations as

$$\left(\frac{x_k^+}{x_k^-}\right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{M_L^I} \frac{x_k^+ - x_j^-}{x_k^- - x_j^+} \mathcal{S}_{kj}^{LL} \prod_{j=1}^{M_L^{II}} \frac{x_k^- - y_j}{x_k^+ - y_j} \prod_{j=1}^{M_R^I} \sqrt{\frac{1 - \frac{1}{x_k^+ \tilde{x}_j^+}}{1 - \frac{1}{x_k^- \tilde{x}_j^-}}} \mathcal{S}_{kj}^{LR} \prod_{j=1}^{M_R^{II}} \frac{1 - \frac{1}{x_k^- \tilde{y}_j}}{1 - \frac{1}{x_k^+ \tilde{y}_j}}, \quad (6.46)$$

$$1 = \prod_{j=1}^{K_L^I} \frac{y_k - x_j^+}{y_k - x_j^-} \prod_{j=1}^{K_R^I} \frac{1 - \frac{1}{y_k \tilde{x}_j^-}}{1 - \frac{1}{y_k \tilde{x}_j^+}}, \quad (6.47)$$

$$\left(\frac{\tilde{x}_k^+}{\tilde{x}_k^-}\right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{M_R^I} \mathcal{S}_{kj}^{RR} \prod_{j=1}^{M_R^{II}} \frac{\tilde{x}_k^+ - \tilde{y}_j}{\tilde{x}_k^- - \tilde{y}_j} \prod_{j=1}^{M_L^I} \sqrt{\frac{1 - \frac{1}{\tilde{x}_k^- x_j^-}}{1 - \frac{1}{\tilde{x}_k^+ x_j^+}}} \mathcal{S}_{kj}^{RL} \prod_{j=1}^{M_L^{II}} \frac{1 - \frac{1}{\tilde{x}_k^+ y_j}}{1 - \frac{1}{\tilde{x}_k^- y_j}}, \quad (6.48)$$

$$1 = \prod_{j=1}^{M_R^{II}} \frac{\tilde{y}_k - \tilde{x}_j^-}{\tilde{y}_k - \tilde{x}_j^+} \prod_{j=1}^{M_L^{II}} \frac{1 - \frac{1}{\tilde{y}_k x_j^-}}{1 - \frac{1}{\tilde{y}_k x_j^+}}. \quad (6.49)$$

These equations should be supplemented by the level matching condition

$$1 = \prod_{j=1}^{M_L^I} \frac{x_j^+}{x_j^-} \prod_{j=1}^{M_R^I} \frac{\tilde{x}_j^+}{\tilde{x}_j^-}. \quad (6.50)$$

By these equations, we can find the asymptotic spectrum of the theory. In fact, once we specify a state by its classical dimension and angular momenta, the labels $M_{L,R}^I$, $M_{L,R}^{II}$ and ℓ will be fixed. Then it only remains to solve the above equations: first for the auxiliary roots y, \tilde{y} and then for the momenta. Using those, we can compute the energy by the dispersion relation

$$E(\{x^\pm, \tilde{x}^\pm\}) = \sum_{j=1}^{M_L^I} \sqrt{1 + 4h^2 \sin^2 \frac{p_j}{2}} + \sum_{j=1}^{M_R^I} \sqrt{1 + 4h^2 \sin^2 \frac{\tilde{p}_j}{2}}. \quad (6.51)$$

Since the S matrix and Bethe ansatz that we just considered now does not correspond to the physical theory we are interested in, we will postpone the identification of the excitation numbers with physical charges to the next section, where we will consider the full $\mathfrak{psu}(1,1|2)^2$ Bethe ansatz. Before moving to that, remark that at the beginning of the nesting procedure we choose the set of excitations V_A^I in (6.17). The other choice quite clearly would lead to a set of equation differing by $L \leftrightarrow R$, and hopefully this will leave the spectrum unchanged. This is actually the case, as we show in appendix A.5 by a *duality transformation* of the equations [27, 156].

Worldsheet picture

In the worldsheet picture, the derivation of the Bethe ansatz equations (more properly, of the Bethe-Yang equations) follows the same logic as above. The nesting procedure is

exactly the same, since the form of the S matrix (*i.e.* its non-vanishing entries) is the same. Again, we have two choices of level-I excitations, given by V_A^I, V_B^I as in (6.17). For the choice V_A^I , the level-I S matrices are as before

$$S_{LL}^{I,I} = A^{LL}, \quad S_{RR}^{I,I} = F^{RR}, \quad S_{LR}^{I,I} = C^{LR}, \quad S_{RL}^{I,I} = D^{RL}. \quad (6.52)$$

Now, however, these elements should be read off the *worldsheet* fundamental S matrix of section 3.3. As we discussed at length in the previous chapter, those elements differ from their spin-chain counterparts for being *their inverse*⁴ and featuring additional factors of the form e^{ip} .

The diagonalisation works in the same way. In fact, even the compatibility equations that fix the higher-level S-matrix elements $S^{II,I}$, take essentially the same form as above, *e.g.* as in (6.30). In the worldsheet picture however we have no length-changing effects, and the non-trivial coproduct is entirely encoded in the form of the S matrix elements of section 3.3. Unsurprisingly, we find the inverse of the spin-chain results up to some factors of e^{ip} . More specifically, we have for the level-II excitations coming from q^L

$$\begin{aligned} S_{LL}^{II,I}(y^L, p) &= \frac{y^L - x_p^-}{y^L - x_p^+} e^{+\frac{i}{2}p}, & S_{RR}^{II,I}(y^L, p) &= \frac{1 - \frac{1}{y^L x_p^+}}{1 - \frac{1}{y^L x_p^-}} e^{+\frac{i}{2}p}, \\ S_{RL}^{II,I}(y^L, p) &= S_{LL}^{II,I}(y^L, p), & S_{LR}^{II,I}(y^L, p) &= S_{RR}^{II,I}(y^L, p), \end{aligned} \quad (6.53)$$

and for the ones coming from \bar{q}^R

$$\begin{aligned} S_{LL}^{II,I}(y^R, p) &= \frac{1 - \frac{1}{y^R x_p^-}}{1 - \frac{1}{y^R x_p^+}} e^{-\frac{i}{2}p}, & S_{RR}^{II,I}(y^R, p) &= \frac{y^R - x_p^+}{y^R - x_p^-} e^{-\frac{i}{2}p}, \\ S_{RL}^{II,I}(y^R, p) &= S_{LL}^{II,I}(y^R, p), & S_{LR}^{II,I}(y^R, p) &= S_{RR}^{II,I}(y^R, p), \end{aligned} \quad (6.54)$$

while again for all the excitations it is

$$S^{II,II} = 1. \quad (6.55)$$

Using as before the short-hand notation (6.45), we can write the Bethe-Yang equations

⁴Looking at formula (5.79) we see that a graded permutation appears in the map between the spin-chain and worldsheet frame. Since we are dealing with diagonal elements and due to left-right symmetry, this plays no role.

as

$$\left(\frac{x_k^+}{x_k^-}\right)^{\ell+\delta} = e^{\frac{i}{2}P_{\text{tot.}}} \prod_{\substack{j=1 \\ j \neq k}}^{M_L^I} \frac{x_k^+ - x_j^-}{x_k^- - x_j^+} \mathcal{S}_{kj}^{LL} \prod_{j=1}^{M_L^{II}} \frac{x_k^- - y_j}{x_k^+ - y_j} \prod_{j=1}^{M_R^I} \sqrt{\frac{1 - \frac{1}{x_k^+ \tilde{x}_j^+}}{1 - \frac{1}{x_k^- \tilde{x}_j^-}}} \mathcal{S}_{kj}^{LR} \prod_{j=1}^{M_R^{II}} \frac{1 - \frac{1}{x_k^- \tilde{y}_j}}{1 - \frac{1}{x_k^+ \tilde{y}_j}}, \quad (6.56)$$

$$1 = e^{-\frac{i}{2}P_{\text{tot.}}} \prod_{j=1}^{M_L^I} \frac{y_k - x_j^+}{y_k - x_j^-} \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{y_k \tilde{x}_j^-}}{1 - \frac{1}{y_k \tilde{x}_j^+}}, \quad (6.57)$$

$$\left(\frac{\tilde{x}_k^+}{\tilde{x}_k^-}\right)^{\ell+\delta} = \prod_{\substack{j=1 \\ j \neq k}}^{M_R^I} \mathcal{S}_{kj}^{RR} \prod_{j=1}^{M_R^{II}} \frac{\tilde{x}_k^+ - \tilde{y}_j}{\tilde{x}_k^- - \tilde{y}_j} \prod_{j=1}^{M_L^I} \sqrt{\frac{1 - \frac{1}{\tilde{x}_k^- x_j^-}}{1 - \frac{1}{\tilde{x}_k^+ x_j^+}}} \mathcal{S}_{kj}^{RL} \prod_{j=1}^{M_L^{II}} \frac{1 - \frac{1}{\tilde{x}_k^+ y_j}}{1 - \frac{1}{\tilde{x}_k^- y_j}}, \quad (6.58)$$

$$1 = e^{+\frac{i}{2}P_{\text{tot.}}} \prod_{j=1}^{M_R^{II}} \frac{\tilde{y}_k - \tilde{x}_j^-}{\tilde{y}_k - \tilde{x}_j^+} \prod_{j=1}^{M_L^{II}} \frac{1 - \frac{1}{\tilde{y}_k x_j^+}}{1 - \frac{1}{\tilde{y}_k x_j^-}}, \quad (6.59)$$

where now ℓ has the interpretation of the worldsheet size⁵ and we introduced the shift

$$\delta = -\frac{1}{2}M_L^I + \frac{1}{2}M_L^{II} - \frac{1}{2}M_R^{II}. \quad (6.60)$$

In order to rewrite the equations in terms of this quantity and the total momentum $P_{\text{tot.}}$, we observed that whenever we multiply the momentum-carrying terms by an expression such as $e^{i(p f(q) - q f(p))}$, we can collect

$$\prod_{\substack{j=1 \\ j \neq k}}^{M_L^I} e^{i(p_k f(p_j) - p_j f(p_k))} \prod_{j=1}^{M_R^I} e^{i(p_k f(p_j) - p_j f(p_k))} = e^{i p_k F_{\text{tot.}}} e^{-i P_{\text{tot.}} f(p_k)}, \quad (6.61)$$

where $F_{\text{tot.}}$ is the sum of $f(p_j)$ on all momentum-carrying excitations. In this simple case, we can use this identity with $f(p) = 1$. We then have that the only new features of the worldsheet Bethe ansatz are a shift in the notion of length with respect to the spin-chain one, and some (fractionary) powers of the total momentum.

6.3 The $\mathfrak{psu}(1, 1|2)^2$ Bethe ansatz

Let us finally work out the Bethe equations for the full $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ S matrix, again in a spin-chain picture. The derivation will be similar to the one we worked out in detail in the previous section, and we will only sketch the most conceptual points, referring the reader to ref. [S3] for more details.

⁵Strictly speaking, in this simple exercise the Bethe equations are not supposed to be matched with a genuine worldsheet theory—we will use the full $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ S matrix for that purpose.

	\mathbb{Q}^{1L}	\mathbb{Q}^{2L}	$\overline{\mathbb{Q}}^{1R}$	$\overline{\mathbb{Q}}^{2R}$
Φ_{++}^L	Φ_{-+}^L	Φ_{+-}^L	$\Phi_{-+}^L Z^-$	$\Phi_{+-}^L Z^-$
Φ_{--}^R	$\Phi_{+-}^R Z^+$	$\Phi_{-+}^R Z^+$	Φ_{+-}^R	Φ_{-+}^R

Table 6.1: Action of the lowering operators on the states of the level-II vacuum V_A^{II} , including length-changing effects in the spin-chain picture.

Sketch of the nesting procedure

Again, we want to construct asymptotic eigenstates of the multi-magnon S matrix. We start from the level-I vacuum, which again is just $|0\rangle^{\text{I}} \equiv |Z^\ell\rangle$. To proceed with nesting, we need to choose a maximal set of excitations that scatter by pure transmission among each other. The structure of \mathcal{S} leads to four possible choices

$$\begin{aligned}
 V_A^{\text{II}} &= \{\Phi_{++}^L, \Phi_{--}^R\}, & V_B^{\text{II}} &= \{\Phi_{--}^L, \Phi_{++}^R\}, \\
 V_C^{\text{II}} &= \{\Phi_{+-}^L, \Phi_{-+}^R\}, & V_D^{\text{II}} &= \{\Phi_{-+}^L, \Phi_{+-}^R\}.
 \end{aligned} \tag{6.62}$$

Each candidate level-II vacuum is composed of one left and one right excitation, that are either both bosonic or both fermionic. Correspondingly, we will have different choices of the lowering operators. In the following we will choose the set V_A^{II} to construct the level-II vacuum. As we expect, the other possible choices are related by dualities, allowing us to write all-loop Bethe equations in four different ways, as we discuss in appendix A.5.

Level-II excitations can be found as before by acting with the lowering operators on the level-II vacuum. Now we have four supercharges at our disposal, *i.e.* $\mathbb{Q}^{1L}, \mathbb{Q}^{2L}$ and $\overline{\mathbb{Q}}^{1R}, \overline{\mathbb{Q}}^{2R}$. In table 6.1 we collect their action on the fields in V_A^{II} , writing also the length-changing effect.

The presence of additional fields and charges produces a new feature: the fields Φ_{--}^L, Φ_{++}^R can be created by acting twice with the supercharges (*i.e.*, respectively by consecutively applying \mathbb{Q}^{1L} and \mathbb{Q}^{2L} on Φ_{++}^L and $\overline{\mathbb{Q}}^{1R}$ and $\overline{\mathbb{Q}}^{2R}$ on $\overline{\Phi}_{--}^R$) and therefore can be considered as composite excitations. As such, they will not explicitly appear in the Bethe ansatz, but rather be represented by pair of suitable excitations.

Apart from this, the calculations to diagonalise the S matrix are exactly the same as the ones of the previous section. This is not surprising, since the lowering operators we should consider now are tensor products of the previous ones, and we deal again with doublets of $\mathfrak{su}(1|1)$ —now we have *e.g.* $(\overline{\Phi}_{++}^L | \Phi_{-+}^L)$ instead of $(\phi^L | \psi^L)$. In particular, we find once again non-trivial matrices $S^{\text{II,I}}$, with the same functional form as before, and trivial $S^{\text{II,II}}$ matrix for all level-II excitations.

Bethe equations

As before, to obtain the Bethe ansatz equations we impose periodic boundary conditions on a spin-chain of finite length ℓ and use the S matrix in its diagonal form. We have again two momentum-carrying excitations, *i.e.* the ones in V_A^I . We denote as before the corresponding Zukhovski variables by x^\pm and \tilde{x}^\pm for left and right excitations respectively. Their number is denoted by M_L^I and M_R^I . We have two auxiliary ‘‘left’’ roots denoted by y_1, y_2 , corresponding respectively to the action of the supercharges $\mathbb{Q}^{1L}, \mathbb{Q}^{2L}$. The two auxiliary ‘‘right’’ roots are denoted by $y_{\bar{1}}, y_{\bar{2}}$ and they correspond respectively to the action of the supercharges $\mathbb{Q}^{1R}, \mathbb{Q}^{2R}$. The number of the corresponding excitations is denoted by $M_{1L}^{II}, M_{2L}^{II}, M_{1R}^{II}$, and M_{2R}^{II} . The Bethe equations then read

$$1 = \prod_{j=1}^{M_L^I} \frac{y_{1,k} - x_j^+}{y_{1,k} - x_j^-} \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{y_{1,k} \tilde{x}_j^-}}{1 - \frac{1}{y_{1,k} \tilde{x}_j^+}}, \quad (6.63)$$

$$\begin{aligned} \left(\frac{x_k^+}{x_k^-} \right)^\ell &= \prod_{\substack{j=1 \\ j \neq k}}^{K_L^I} \frac{x_k^+ - x_j^+}{x_k^- - x_j^-} \frac{1 - \frac{1}{x_k^+ x_j^-}}{1 - \frac{1}{x_k^- x_j^+}} \sigma^2(x_k, x_j) \prod_{j=1}^{M_{1L}^{II}} \frac{x_k^- - y_{1,j}}{x_k^+ - y_{1,j}} \prod_{j=1}^{M_{2L}^{II}} \frac{x_k^- - y_{2,j}}{x_k^+ - y_{2,j}} \\ &\quad \times \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{x_k^+ \tilde{x}_j^+}}{1 - \frac{1}{x_k^- \tilde{x}_j^-}} \frac{1 - \frac{1}{x_k^+ \tilde{x}_j^-}}{1 - \frac{1}{x_k^- \tilde{x}_j^+}} \tilde{\sigma}^2(x_k, \tilde{x}_j) \prod_{j=1}^{M_{1R}^{II}} \frac{1 - \frac{1}{x_k^- y_{1,j}}}{1 - \frac{1}{x_k^+ y_{1,j}}} \prod_{j=1}^{M_{2R}^{II}} \frac{1 - \frac{1}{x_k^- y_{2,j}}}{1 - \frac{1}{x_k^+ y_{2,j}}}, \end{aligned} \quad (6.64)$$

$$1 = \prod_{j=1}^{M_L^I} \frac{y_{2,k} - x_j^+}{y_{2,k} - x_j^-} \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{y_{2,k} \tilde{x}_j^-}}{1 - \frac{1}{y_{2,k} \tilde{x}_j^+}}, \quad (6.65)$$

$$1 = \prod_{j=1}^{M_R^I} \frac{y_{\bar{1},k} - \tilde{x}_j^-}{y_{\bar{1},k} - \tilde{x}_j^+} \prod_{j=1}^{M_L^I} \frac{1 - \frac{1}{y_{\bar{1},k} x_j^+}}{1 - \frac{1}{y_{\bar{1},k} x_j^-}}, \quad (6.66)$$

$$\begin{aligned} \left(\frac{\tilde{x}_k^+}{\tilde{x}_k^-} \right)^\ell &= \prod_{\substack{j=1 \\ j \neq k}}^{M_R^I} \frac{\tilde{x}_k^- - \tilde{x}_j^+}{\tilde{x}_k^+ - \tilde{x}_j^-} \frac{1 - \frac{1}{\tilde{x}_k^- \tilde{x}_j^+}}{1 - \frac{1}{\tilde{x}_k^+ \tilde{x}_j^-}} \sigma^2(\tilde{x}_k, \tilde{x}_j) \prod_{j=1}^{M_{1R}^{II}} \frac{\tilde{x}_k^+ - y_{\bar{1},j}}{\tilde{x}_k^- - y_{\bar{1},j}} \prod_{j=1}^{M_{2R}^{II}} \frac{\tilde{x}_k^+ - y_{\bar{2},j}}{\tilde{x}_k^- - y_{\bar{2},j}} \\ &\quad \times \prod_{j=1}^{M_L^I} \frac{1 - \frac{1}{\tilde{x}_k^- x_j^-}}{1 - \frac{1}{\tilde{x}_k^+ x_j^+}} \frac{1 - \frac{1}{\tilde{x}_k^- x_j^+}}{1 - \frac{1}{\tilde{x}_k^+ x_j^-}} \tilde{\sigma}^2(\tilde{x}_k, x_j) \prod_{j=1}^{M_{1L}^{II}} \frac{1 - \frac{1}{\tilde{x}_k^+ y_{1,j}}}{1 - \frac{1}{\tilde{x}_k^- y_{1,j}}} \prod_{j=1}^{M_{2L}^{II}} \frac{1 - \frac{1}{\tilde{x}_k^+ y_{2,j}}}{1 - \frac{1}{\tilde{x}_k^- y_{2,j}}}, \end{aligned} \quad (6.67)$$

$$1 = \prod_{j=1}^{M_R^I} \frac{y_{\bar{2},k} - \tilde{x}_j^-}{y_{\bar{2},k} - \tilde{x}_j^+} \prod_{j=1}^{M_L^I} \frac{1 - \frac{1}{y_{\bar{2},k} x_j^+}}{1 - \frac{1}{y_{\bar{2},k} x_j^-}}. \quad (6.68)$$

Note that, since we are dealing with the physical S matrix for the massive modes, we normalised the level-I scattering factors in terms of the canonical dressing factors σ_{pq}

and $\tilde{\sigma}_{pq}$. The level matching condition is once again

$$\prod_j^{M_L^I} \frac{x_j^+}{x_j^-} \prod_j^{M_R^I} \frac{\tilde{x}_j^+}{\tilde{x}_j^-} = 1. \quad (6.69)$$

The total energy of a multi-excitation state that satisfies the Bethe equations and the level matching condition is given by

$$E(\{x^\pm, \tilde{x}^\pm\}) = \sum_{j=1}^{M_L^I} \sqrt{1 + 4h^2 \sin^2 \frac{p_j}{2}} + \sum_{j=1}^{M_R^I} \sqrt{1 + 4h^2 \sin^2 \frac{\tilde{p}_j}{2}}. \quad (6.70)$$

Let us now analyse the small- h limit, which in the spin-chain picture is the most natural, since there the left and right chains decouple and the dynamical length-changing effects are suppressed. We will come back to the large- h limit in section 7.2, when we will compare it with an independent result, the “finite-gap” equations.

Small- h limit and Cartan matrix

When we are dealing with non-dynamical spin chains having a Lie (super)algebra structure, the Bethe equations can immediately be written down using data from the algebra and its representations [15, 157]. In particular, in the case of $\mathfrak{psu}(1, 1|2)^2$, we should write

$$\left(\frac{u_{l,k} + \frac{i}{2} w_l}{u_{l,k} - \frac{i}{2} w_l} \right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{K_l} \frac{u_{l,k} - u_{l,j} + \frac{i}{2} A_{ll}}{u_{l,k} - u_{l,j} - \frac{i}{2} A_{ll}} \prod_{l' \neq l} \left(\prod_{j=1}^{K_{l'}} \frac{u_{l,k} - u_{l',j} + \frac{i}{2} A_{ll'}}{u_{l,k} - u_{l',j} - \frac{i}{2} A_{ll'}} \right), \quad (6.71)$$

where w_l are weights, $A_{ll'}$ are elements of the Cartan matrix of $\mathfrak{psu}(1, 1|2)^2$, and K_l are excitation numbers pertaining to each Cartan element. Since, as discussed in appendix A.1, each copy of $\mathfrak{psu}(1, 1|2)$ has rank 3, we expect to find 6 equations from this construction.

On the other hand, when $h \ll 1$, we can expand the Zukhovski variables in terms of the spin-chain rapidity u as

$$x^\pm \approx \frac{u_x \pm i/2}{h/2}, \quad y \approx \frac{u_y}{h/2}, \quad (6.72)$$

in the left sector, where u_i are finite as $h \rightarrow 0$, and similarly in the right sector. In these terms, we have the familiar formulae (see *e.g.* [129])

$$p \approx \frac{u + \frac{i}{2}}{u - \frac{i}{2}}, \quad E = 1 + \delta D \approx 1 + \frac{2h^2}{1 + 4u^2}, \quad (6.73)$$

where in the last formula we split the energy into a classical contribution and the correction due to the anomalous dimension, that for M excitations is

$$\delta D(p) = E(p) - M = -ih \sum_{k=1}^M \left(\frac{1}{x_k^-} - \frac{1}{x_k^+} \right), \quad (6.74)$$




Figure 6.3: Three Dynkin diagrams for $\mathfrak{psu}(1,1|2)$. From left to right, we depict the $\mathfrak{su}(2)$ grading, two fully fermionic gradings and the $\mathfrak{sl}(2)$ grading.

where the sum is over all (left and right) momentum-carrying excitations.

If we assume that the dressing phases σ and $\tilde{\sigma}$ expand trivially in this limit⁶, we indeed find that the Bethe ansatz takes the form (6.71) and we can read off the resulting Cartan matrix

$$A = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & +2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & +1 & 0 \\ 0 & 0 & 0 & +1 & -2 & +1 \\ 0 & 0 & 0 & 0 & +1 & 0 \end{pmatrix}. \quad (6.75)$$

Comparing this with the Cartan matrices in appendix A.1, in particular equations (A.3) and (A.5), we see that this does correspond to $\mathfrak{psu}(1,1|2)^2$, with different gradings for the two factors of the algebra. The left copy is in the $\mathfrak{su}(2)$ grading, while the right one is in the $\mathfrak{sl}(2)$ grading, represented by the Dynkin diagrams of figure 6.3a and 6.3c. This shows that there is a strict connection between the type of particles appearing in the Bethe ansatz and the nodes of the Dynkin diagrams. In the Cartan basis is natural to identify the nodes by 1, 2, 3 for the left algebra and $\tilde{1}, \tilde{2}, \tilde{3}$ for the right one. We can identify the momentum-carrying excitations with the middle nodes,

$$M_{\text{L}}^{\text{I}} = K_2, \quad M_{\text{R}}^{\text{I}} = K_{\tilde{2}}, \quad (6.76)$$

and the remaining ones with the peripheral nodes

$$M_{\text{1L}}^{\text{II}} = K_1, \quad M_{\text{2L}}^{\text{II}} = K_3, \quad M_{\text{1R}}^{\text{II}} = K_{\tilde{1}}, \quad M_{\text{2R}}^{\text{II}} = K_{\tilde{3}}. \quad (6.77)$$

In this way it is easy to represent the Bethe equations pictorially, as in figure 6.4. There, the solid links are the ones that survive the $\hbar \rightarrow 0$ limit (together with the self interaction of the momentum-carrying nodes), and therefore can be read off directly from the Dynkin diagram. The dashed ones can be interpreted as arising from the \mathbb{Z}_4 -graded structure of

⁶ In the next chapter we will return on this issue and see that this is not exactly the case.




Figure 6.4: The Dynkin diagram for $\mathfrak{psu}(1,1|2)^2$ with the various interaction terms appearing in the Bethe ansatz indicated. The label ± 1 inside the middle Dynkin nodes indicate the $\mathfrak{su}(2)$ and $\mathfrak{sl}(2)$ gradings of the left- and right-moving sectors.

the model, and finally the curly lines are couplings involving the dressing phases, which as discussed can appear only between momentum-carrying nodes.

Not surprisingly, different Dynkin diagrams would appear, should we choose different level-I excitations in nesting procedure. As we discuss in appendix A.5, this amounts to dualisation of the Bethe equations, *i.e.* to replacing a set of roots with an equivalent one. In particular, after dualisation of the nodes 1 and $\tilde{1}$ the Bethe equations are written in a different grading, where all the nodes of the Dynkin diagrams are fermionic. From the weak coupling expansion we get the Cartan matrix

$$\tilde{A} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \quad (6.78)$$

corresponding to the fermionic gradings in (A.6). If we had dualised the nodes 3 and $\tilde{3}$ instead we would have found the Cartan matrix $-\tilde{A}$. The consecutive dualisation of 1, $\tilde{1}$ and 3, $\tilde{3}$ gives the Cartan matrix $-A$. These last two choices are once again a manifestation of left-right symmetry.

Global charges

By expanding the Bethe equations around large values of the spectral parameter u we should obtain the global charges of the symmetry algebra [27, 112]. In doing so we will assume again that the dressing factors σ_{pq} and $\tilde{\sigma}_{pq}$ do not contribute to the charges.⁷

As we have seen above, the left- and right-moving sectors of the Bethe equations are naturally written using different gradings of the $\mathfrak{psu}(1,1|2)$ algebra. The Dynkin labels

⁷ In the next chapter we will see that the dressing factors do contribute, and in a troubling way.

r_1, r_2 and r_3 for the left-movers therefore give the eigenvalues of the Cartan generators \mathfrak{h}_j given in (A.2), while the labels $r_{\bar{1}}, r_{\bar{2}}$ and $r_{\bar{3}}$ for the right-movers correspond to the generators $\tilde{\mathfrak{h}}_j$ in (A.4). Expanding the Bethe equations around small values of the momentum (or equivalently large values of the rapidity u) we find

$$\begin{aligned} r_1 = r_3 &= +K_2 + \frac{1}{2}\delta D, & r_2 &= \ell + K_{\bar{1}} - 2K_{\bar{2}} + K_{\bar{3}}, \\ r_{\bar{1}} = r_{\bar{3}} &= -K_{\bar{2}} - \frac{1}{2}\delta D, & r_{\bar{2}} &= \ell - K_{\bar{1}} + 2K_{\bar{2}} - K_{\bar{3}} + \delta D, \end{aligned} \quad (6.79)$$

where the anomalous dimension δD is given by (6.74).

A representation of $\mathfrak{psu}(1,1|2)^2$ can be labeled by the eigenvalues of the highest weight state under the four generators $\mathbb{L}_3^L, \mathbb{L}_3^R, \mathbb{J}_3^L$ and \mathbb{J}_3^R . It is useful to combine them into the charges

$$\begin{aligned} \mathbb{D} &= \mathbb{L}_3^L + \mathbb{L}_3^R, & \mathbb{J} &= \mathbb{J}_3^L + \mathbb{J}_3^R, \\ \mathbb{L} &= \mathbb{L}_3^L - \mathbb{L}_3^R, & \mathbb{K} &= \mathbb{J}_3^L - \mathbb{J}_3^R. \end{aligned} \quad (6.80)$$

The most important of these is the generator of dilatations \mathbb{D} , that is the dual of the target-space energy. It is related to the spin-chain Hamiltonian \mathbb{H} by

$$\mathbb{H} = \mathbb{D} - \mathbb{J}, \quad (6.81)$$

a relation that we have already seen in (2.17) in terms of the worldsheet theory.

We can now express the eigenvalues of the generators (6.80) in terms of the excitation numbers K_j as

$$\begin{aligned} D &= +K_{\bar{2}} + \frac{1}{2}(K_1 + K_3 - K_{\bar{1}} - K_{\bar{3}}) + \ell + \delta D, \\ L &= -K_{\bar{2}} + \frac{1}{2}(K_1 + K_3 + K_{\bar{1}} + K_{\bar{3}}), \\ J &= -K_2 + \frac{1}{2}(K_1 + K_3 - K_{\bar{1}} - K_{\bar{3}}) + \ell, \\ K &= -K_2 + \frac{1}{2}(K_1 + K_3 + K_{\bar{1}} + K_{\bar{3}}). \end{aligned} \quad (6.82)$$

Note that the anomalous dimension δD only contributes to the eigenvalue Δ of the dilatation operator. The eigenvalue of the Hamiltonian now takes the form

$$E = \Delta - J = K_2 + K_{\bar{2}} + \delta D, \quad (6.83)$$

as expected from equation (6.74).

Worksheet picture

As we have seen in the previous section, the Bethe ansatz in the worldsheet picture is not very different from the spin-chain one, and in fact can be found in an analogous way.

The resulting equations read

$$1 = e^{-\frac{i}{2}P_{\text{tot.}}} \prod_{j=1}^{M_L^I} \frac{y_{1,k} - x_j^+}{y_{1,k} - x_j^-} \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{y_{1,k} \tilde{x}_j^-}}{1 - \frac{1}{y_{1,k} \tilde{x}_j^+}}, \quad (6.84)$$

$$\begin{aligned} \left(\frac{x_k^+}{x_k^-} \right)^{\ell+\delta} &= e^{iP_{\text{tot.}}} \prod_{\substack{j=1 \\ j \neq k}}^{K_L^I} \frac{x_k^+ - x_j^-}{x_k^- - x_j^+} \frac{1 - \frac{1}{x_k^+ x_j^-}}{1 - \frac{1}{x_k^- x_j^+}} \sigma^2(x_k, x_j) \prod_{j=1}^{M_{1L}^{II}} \frac{x_k^- - y_{1,j}}{x_k^+ - y_{1,j}} \prod_{j=1}^{M_{2L}^{II}} \frac{x_k^- - y_{2,j}}{x_k^+ - y_{2,j}} \\ &\times \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{x_k^+ \tilde{x}_j^-}}{1 - \frac{1}{x_k^- \tilde{x}_j^+}} \frac{1 - \frac{1}{x_k^+ \tilde{x}_j^-}}{1 - \frac{1}{x_k^- \tilde{x}_j^+}} \tilde{\sigma}^2(x_k, \tilde{x}_j) \prod_{j=1}^{M_{1R}^{II}} \frac{1 - \frac{1}{x_k^- y_{1,j}}}{1 - \frac{1}{x_k^+ y_{1,j}}} \prod_{j=1}^{M_{2R}^{II}} \frac{1 - \frac{1}{x_k^- y_{2,j}}}{1 - \frac{1}{x_k^+ y_{2,j}}}, \end{aligned} \quad (6.85)$$

$$1 = e^{-\frac{i}{2}P_{\text{tot.}}} \prod_{j=1}^{M_L^I} \frac{y_{2,k} - x_j^+}{y_{2,k} - x_j^-} \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{y_{2,k} \tilde{x}_j^-}}{1 - \frac{1}{y_{2,k} \tilde{x}_j^+}}, \quad (6.86)$$

$$1 = e^{+\frac{i}{2}P_{\text{tot.}}} \prod_{j=1}^{M_R^I} \frac{y_{\bar{1},k} - \tilde{x}_j^-}{y_{\bar{1},k} - \tilde{x}_j^+} \prod_{j=1}^{M_L^I} \frac{1 - \frac{1}{y_{\bar{1},k} x_j^+}}{1 - \frac{1}{y_{\bar{1},k} x_j^-}}, \quad (6.87)$$

$$\begin{aligned} \left(\frac{\tilde{x}_k^+}{\tilde{x}_k^-} \right)^{\ell+\delta} &= \prod_{\substack{j=1 \\ j \neq k}}^{M_R^I} \frac{\tilde{x}_k^- - \tilde{x}_j^+}{\tilde{x}_k^+ - \tilde{x}_j^-} \frac{1 - \frac{1}{\tilde{x}_k^- \tilde{x}_j^+}}{1 - \frac{1}{\tilde{x}_k^+ \tilde{x}_j^-}} \sigma^2(\tilde{x}_k, \tilde{x}_j) \prod_{j=1}^{M_{1R}^{II}} \frac{\tilde{x}_k^+ - y_{\bar{1},j}}{\tilde{x}_k^- - y_{\bar{1},j}} \prod_{j=1}^{M_{2R}^{II}} \frac{\tilde{x}_k^+ - y_{\bar{2},j}}{\tilde{x}_k^- - y_{\bar{2},j}} \\ &\times \prod_{j=1}^{M_L^I} \frac{1 - \frac{1}{\tilde{x}_k^- x_j^-}}{1 - \frac{1}{\tilde{x}_k^+ x_j^+}} \frac{1 - \frac{1}{\tilde{x}_k^- x_j^-}}{1 - \frac{1}{\tilde{x}_k^+ x_j^+}} \tilde{\sigma}^2(\tilde{x}_k, x_j) \prod_{j=1}^{M_{1L}^{II}} \frac{1 - \frac{1}{\tilde{x}_k^- y_{\bar{1},j}}}{1 - \frac{1}{\tilde{x}_k^+ y_{\bar{1},j}}} \prod_{j=1}^{M_{2L}^{II}} \frac{1 - \frac{1}{\tilde{x}_k^- y_{\bar{3},j}}}{1 - \frac{1}{\tilde{x}_k^+ y_{\bar{2},j}}}, \end{aligned} \quad (6.88)$$

$$1 = e^{+\frac{i}{2}P_{\text{tot.}}} \prod_{j=1}^{M_R^I} \frac{y_{\bar{2},k} - \tilde{x}_j^-}{y_{\bar{2},k} - \tilde{x}_j^+} \prod_{j=1}^{M_L^I} \frac{1 - \frac{1}{y_{\bar{2},k} x_j^+}}{1 - \frac{1}{y_{\bar{2},k} x_j^-}}, \quad (6.89)$$

where now

$$\delta = -M_L^I + \frac{1}{2}M_{1L}^{II} + \frac{1}{2}M_{1L}^{II} - \frac{1}{2}M_{1R}^{II} - \frac{1}{2}M_{2R}^{II}. \quad (6.90)$$

By comparing with (6.82), we see that the difference between the spin-chain and worldsheet notion of length is given by the charge J , which is the S^3 angular momentum. Moreover, the total-momentum contributions drop out of the equations upon imposition of the level-matching constraint—the spin-chain and worldsheet construction are equivalent.

Let us remark that ℓ is not a gauge-invariant quantity. Going back to the (*e.g.* bosonic) string theory discussion of chapter 2, we have seen that $\ell = P_+$ depends on the choice of lighcone gauge-fixing. In a more general a -dependent gauge fixing [55, 117, 118], we would have

$$\ell = P_+ = a E^{\text{t.s.}} + (1-a) J = J + a E. \quad (6.91)$$

Since the spectrum is supposed to be gauge invariant, the dependence on a should drop from the Bethe-Yang equation. This can be understood in terms of an overall prefactor

tor for the S matrix, which in particular has a non-trivial momentum dependence in the form $e^{i a (p \delta D_q - q \delta D_p)}$. It is easy to check that this factor drops out of the crossing equations, and therefore cannot be fixed based on our symmetry arguments. Moreover, from (6.61) we see that this produces exactly the energy-dependent shift that can cancel the a -dependent part of ℓ . A tree-level perturbative calculation [100–102] immediately shows that the factor appears in such a way as to cancel the gauge dependence, as it is expected from what happens in $\text{AdS}_5 \times S^5$, see *e.g.* ref. [158].

6.4 Chapter summary

In this chapter we have worked out the asymptotic Bethe ansatz equations for the all-loop S matrix we found earlier. We outlined the procedure for doing so in a spin-chain and worldsheet picture, that turned out to be equivalent.

The Bethe equations can be nicely summarised in figure 6.4, where the excitations are represented by the corresponding nodes of the $\mathfrak{psu}(1, 1|2)^2$ Dynkin diagram. This is a manifestation of the $\mathfrak{psu}(1, 1|2)^2$ symmetry of the spectrum, which in the Bethe ansatz construction is expected [86], see also the discussion in chapter 8. As we have already seen a number of times, such a symmetry requires the two copies of the superalgebra to be in opposite gradings.

As we stressed in the introduction, these equations describe only the *asymptotic* spectrum, *i.e.* the spectrum when ℓ is finite but large. This is due to wrapping effects: either particles wrapping around the worldsheet or the spin-chain non-local interactions wrapping around the chain. While these effects are exponentially suppressed in ℓ , a complete treatment of the spectral problem requires accounting for them. This can be done by the mirror thermodynamical Bethe ansatz (Mirror TBA), as we briefly sketched in section 1.2. As we mentioned there, this requires having a handle over the whole asymptotic spectrum of the theory, including massless excitations and bound states. For this reason, such a description is for the moment beyond our reach.

7 Comparison with perturbative calculations

It is finally time to compare our results with perturbative ones. This will give us some insight on whether our assumptions of integrability were appropriate.

Since the dual CFT_2 is quite hard to evaluate perturbatively [109], all the calculations so far available are from the string side. Therefore, they all require the limit $h \rightarrow \infty$. This can be taken in several ways, depending on how the magnon momentum scales. We have already encountered in chapter 2 the BMN or near-plane-wave limit [119], whereby the momentum scales as $p \sim \rho/h$ and the Zhukovsky variables expand as

$$x_p^\pm = \left(1 \pm \frac{i\rho}{2h}\right) \frac{(1 + \omega_p)}{p} + \mathcal{O}(1/h^2), \quad \omega_p = \sqrt{1 + \rho^2}. \quad (7.1)$$

In this approximation, the theory becomes relativistic as the dispersion ω_p shows. Another useful limit is the Maldacena-Swanson or near-flat-space (NFS) limit [159], whereby one uses light-cone kinematics and scales $p \sim p_-/\sqrt{h}$. In this regime, loop calculations on the worldsheet are more feasible. Finally, it will be useful to consider the semiclassical limit, whereby we also take the number of excitations to infinity and scale the spectral parameters as

$$x^\pm = x \pm \frac{i}{h} \frac{x^2}{x^2 - 1} + \mathcal{O}\left(\frac{1}{h^3}\right). \quad (7.2)$$

The expansion of the S-matrix elements would be straightforward were not for the dressing factors, that require particular care.

7.1 Expansions of the dressing factors

In chapter 4 we solved the AdS_3 crossing equations (4.29) in terms of (4.47). In what follows we give the strong- and weak-coupling expansions of these all-loop phases.

Strong-coupling expansion

The dressing phases admit an expansion in terms of local conserved charges $q_r(p)$ [25]

$$\theta(p_1, p_2) = \sum_{r=1}^{\infty} \sum_{\substack{s>r \\ r+s=\text{odd}}}^{\infty} c_{r,s}(h) [q_r(p_1)q_s(p_2) - q_r(p_2)q_s(p_1)], \quad (7.3)$$

where $c_{r,s}(h)$ are functions of the coupling constant h with expansion

$$c_{r,s}(h) = hc_{r,s}^{(0)} + c_{r,s}^{(1)} + c_{r,s}^{(2)}h^{-1} + \dots \quad (7.4)$$

and are antisymmetric in r, s . The phase $\tilde{\theta}(p_1, p_2)$ has a similar expansion where the coefficients will be denoted $\tilde{c}_{r,s}(h)$. The expression above is similar to the corresponding one in AdS₅, but unlike what happens in that case, we will need to include the $r = 1$ terms. This new feature was first noted in ref. [103]. For $r \geq 2$ the conserved charges are given by

$$q_r(p_k) = \mathcal{Q}_r(x_k^+) - \mathcal{Q}_r(x_k^-) = \frac{i}{r-1} \left[\frac{1}{(x_k^+)^{r-1}} - \frac{1}{(x_k^-)^{r-1}} \right], \quad (7.5)$$

$$\mathcal{Q}_r(x) = \frac{i}{r-1} \frac{1}{x^{r-1}},$$

where we introduced the function $\mathcal{Q}_r(x_k)$ for later convenience. For $r = 1$ the charge is just the momentum

$$q_1(p_k) = \mathcal{Q}_1(x_k^+) - \mathcal{Q}_1(x_k^-) = -i \log \left(\frac{x_k^+}{x_k^-} \right), \quad \mathcal{Q}_1(x) \equiv i \log \left(\frac{1}{x} \right). \quad (7.6)$$

Expressing $\theta(p_1, p_2)$ in terms of χ (see equation (4.32)), we obtain the expansion

$$\chi(x, y) = \sum_{r=1}^{\infty} \sum_{\substack{s>r \\ r+s=\text{odd}}}^{\infty} c_{r,s}(h) [\mathcal{Q}_r(x)\mathcal{Q}_s(y) - \mathcal{Q}_r(y)\mathcal{Q}_s(x)]. \quad (7.7)$$

with a corresponding expression for $\tilde{\chi}$. The coefficients $c_{r,s}$ and $\tilde{c}_{r,s}$ can be obtained by expanding the integrands through which χ and $\tilde{\chi}$ are defined. This expansion is in $h \gg 1$ and then in $x, y \gg 1$. Recall that our phases involve the Hernández-López [144] and Beisert-Eden-Staudacher phases [28], as well as the phase χ^- of eq. (4.46). The expansions for χ^{BES} and χ^{HL} are well known in the literature, and in particular we have

$$\chi^{\text{HL}}(x, y) = \frac{2}{\pi} \sum_{r=2}^{\infty} \sum_{\substack{s>r \\ r+s=\text{odd}}}^{\infty} \frac{(r-1)(s-1)}{(r-s)(r+s-2)} [\mathcal{Q}_r(x)\mathcal{Q}_s(y) - \mathcal{Q}_r(y)\mathcal{Q}_s(x)]. \quad (7.8)$$

The expansion for χ^- (see equation (4.46)) is

$$\chi^-(x, y) = -\frac{1}{\pi} \sum_{r=2}^{\infty} \sum_{\substack{s>r \\ r+s=\text{odd}}}^{\infty} \frac{(r-1)^2 + (s-1)^2}{(r-s)(r+s-2)} [\mathcal{Q}_r(x)\mathcal{Q}_s(y) - \mathcal{Q}_r(y)\mathcal{Q}_s(x)]$$

$$+ \frac{1}{2\pi} \sum_{\substack{s>1 \\ s=\text{even}}}^{\infty} [\mathcal{Q}_1(x)\mathcal{Q}_s(y) - \mathcal{Q}_1(y)\mathcal{Q}_s(x)]. \quad (7.9)$$

Expanding (4.47) at large h we find

$$\chi(x, y) = h\chi^{\text{AFS}}(x, y) + \frac{1}{2}(\chi^{\text{HL}}(x, y) + \chi^-(x, y)) + \mathcal{O}\left(\frac{1}{h}\right),$$

$$\tilde{\chi}(x, y) = h\chi^{\text{AFS}}(x, y) + \frac{1}{2}(\chi^{\text{HL}}(x, y) - \chi^-(x, y)) + \mathcal{O}\left(\frac{1}{h}\right), \quad (7.10)$$

where we have extracted the h -scaling of each phase. At leading order both phases reduce to the Arutyunov-Frolov-Staudacher one [25]. However, at HL-order all three terms on the right hand side of equation (4.47) contribute and we find

$$\begin{aligned} c_{r,s}^{(1)} &= +\frac{1}{2\pi} \frac{1 - (-1)^{s+r}}{2} \left[\frac{s-r}{s+r-2} - \frac{1}{2} (\delta_{r,1} - \delta_{1,s}) \right], \\ \tilde{c}_{r,s}^{(1)} &= -\frac{1}{2\pi} \frac{1 - (-1)^{s+r}}{2} \left[\frac{s+r-2}{s-r} - \frac{1}{2} (\delta_{r,1} - \delta_{1,s}) \right], \end{aligned} \quad (7.11)$$

for $s > r > 0$. Finally, the higher order coefficients $c_{r,s}^{(n)} = \tilde{c}_{r,s}^{(n)}$ with $n > 1$ are exactly the same as in the expansion of the BES phase.

Semiclassical and near-flat-space limits

In order to compare with perturbative results, it is convenient to write explicit expressions for our phases in the semiclassical limit (7.2). Such an expansion for the BES phase is well known: the leading order $O(1/h)$ is given by the AFS phase (4.35), which in our normalisation reads

$$\theta^{\text{AFS}}(x, y) = \frac{2}{h} \frac{x-y}{(x^2-1)(xy-1)(y^2-1)} + O\left(\frac{1}{h^3}\right), \quad (7.12)$$

whereas the next-to-leading-order is given by the HL phase which can be found by expanding (4.36) under the integral. Doing so also for (4.46), we get to the expressions

$$\begin{aligned} \theta(x, y) &= \theta^{\text{AFS}}(x, y) + \frac{1}{\pi h^2} \frac{x^2}{x^2-1} \frac{y^2}{y^2-1} \left[\frac{(x+y)^2(1-\frac{1}{xy})}{(x^2-1)(x-y)(y^2-1)} \right. \\ &\quad \left. + \frac{2}{(x-y)^2} \log\left(\frac{x+1}{x-1} \frac{y-1}{y+1}\right) \right] + O\left(\frac{1}{h^3}\right), \\ \tilde{\theta}(x, y) &= \theta^{\text{AFS}}(x, y) + \frac{1}{\pi h^2} \frac{x^2}{x^2-1} \frac{y^2}{y^2-1} \left[\frac{(xy+1)^2(\frac{1}{x}-\frac{1}{y})}{(x^2-1)(xy-1)(y^2-1)} \right. \\ &\quad \left. + \frac{2}{(xy-1)^2} \log\left(\frac{x+1}{x-1} \frac{y-1}{y+1}\right) \right] + O\left(\frac{1}{h^3}\right). \end{aligned} \quad (7.13)$$

Let us also evaluate the dressing factors in the near-flat-space limit [159]

$$\begin{aligned} \theta(p_-, q_-) &= \frac{p_- q_- (p_- - q_-)}{8h(p_- + q_-)} + \frac{p_-^2 q_-^2 \left(p_-^2 + 2p_- q_- \log \frac{q_-}{p_-} - q_-^2 \right)}{64\pi h^2 (p_- - q_-)^2} + O\left(\frac{1}{h^3}\right), \\ \tilde{\theta}(p_-, q_-) &= \frac{p_- q_- (p_- - q_-)}{8h(p_- + q_-)} - \frac{p_-^2 q_-^2 \left(p_-^2 - 2p_- q_- \log \frac{q_-}{p_-} - q_-^2 \right)}{64\pi h^2 (p_- + q_-)^2} + O\left(\frac{1}{h^3}\right). \end{aligned} \quad (7.14)$$

Weak-coupling expansion

In this subsection we compute the weak-coupling expansion of the dressing phases. While this will not be comparable with any direct calculation, it naturally enters our spin-chain Bethe ansatz, in particular in the identification of the global charges.

The results for σ^{BES} are well known from AdS₅/CFT₄. The leading-order contribution to the dressing phase starts at $\mathcal{O}(h^6)$ [28], and comes from the $r = 2, s = 3$ terms in the expansion of χ^{BES} .¹ The AdS₃ dressing phases (4.47) contain extra terms besides the BES phase. The coefficients $c_{r,s}$ and $\tilde{c}_{r,s}$ that come from these extra contributions are all order h^0 (see equation (7.9) and (7.8)). The coupling constant dependence comes only from the charges q_r and q_s in equation (7.7) through x^\pm . In fact, the leading contribution comes from the $r = 1$ and $s = 2$ term, and can be written

$$\begin{aligned}\theta(p, q) &= +i h c_{1,2}^{(1)} \left(p \delta D(q) - q \delta D(p) \right) + \mathcal{O}(h^3), \\ \tilde{\theta}(p, q) &= -i h c_{1,2}^{(1)} \left(p \delta D(q) - q \delta D(p) \right) + \mathcal{O}(h^3),\end{aligned}\tag{7.15}$$

where $\delta D(p) = E(p) - 1$ is the anomalous part of the dispersion relation. Note that the $\mathcal{O}(h^2)$ terms vanish.

The above result shows that the $r = 1$ terms, which are novel to AdS₃, contribute at order h to the BA, and so should modify the energy of states in the weakly-coupled spin-chain at order h^3 . Let us remark that *a priori* we do not know how $h(\lambda)$ behaves at weak coupling, where λ is the genuine CFT₂ coupling constant.² This prevents us from determining whether the $\mathcal{O}(h^1)$ contribution to $\theta(p, q)$ in the equation above comes with an integral power of λ as one would expect in a weakly-coupled planar limit.

Integrable spin chains with long-range interactions have been systematically considered in ref. [160] where, among other things, the weak-coupling expansion of the dressing factor and of some conserved charges was analysed. Interestingly, it appears that the weak-coupling properties of the AdS₃ spin chain differ from those found in that general analysis. This is not entirely surprising, since in contrast to ref. [160], the $\mathfrak{psu}(1,1|2)^2$ spin chain consists of both a left-moving and a right-moving sector. Something similar happens in the study of alternating spin chains [161], where novel operators that do not exist for the homogeneous spin chains of ref. [160] modify the structure of the dressing factor found there. We can then conclude that the AdS₃/CFT₂ weakly-coupled spin-chain dynamics is substantially different from the ones we are familiar with.

Let us now investigate how these contributions modify the Bethe ansatz picture. The form of the right hand side of eq. (7.15) is reminiscent of a gauge-dependent shift of the length, as discussed at the end of section 6.2. Indeed, if in the Bethe equations appeared *e.g.* only left excitations and therefore only $\theta(p, q)$ we could use the level-matching to reabsorb such factor in a shift of ℓ as in (6.61), at least in absence of winding. However, as soon as left *and* right excitations are present such a procedure is impossible, because the correction due to $\tilde{\theta}(p, q)$ has opposite sign than the one of $\theta(p, q)$, see (7.15). What is more, such terms are non-integer, and it is easy to see that they would appear in the identification of the global charges (6.79). In particular, the Dynkin label r_2 corresponds

¹See equation (7.7), and recall that for the BES phase there is no $r = 1$ term.

²Recall, for example, that in AdS₅/CFT₄ $h \sim \sqrt{\lambda}$ it is while in AdS₄/CFT₃ it is $h \sim \lambda$.

to $\mathfrak{su}(2)$ and as such should be an integer—but the phase contribution would contain anomalous terms.

At the moment we have no resolution for such a puzzle. It is tempting to say that perhaps the dressing factors should be modified by a solution of the homogeneous crossing equations, in such a way to remove such contributions.³ However one should bear in mind that these $r = 1$ terms appear also in independent string calculations [103, 105], as we will discuss in the next section. This once again points to some subtlety in interpreting these results, and in fact our whole construction, at weak coupling.

7.2 Comparisons

In this section we will look first at explicit S-matrix elements obtained from string theory. These include tree-level, one-loop and two-loop results. While the tree-level terms can be found directly in full generality, evaluating the one-loop ones requires either going to particular kinematic regions or using additional tricks such as unitarity techniques.⁴

For this reason, to investigate the one-loop structure of the dressing factors sometimes a different approach is more useful: one can focus on a specific classical solution of the string NLSM and compute the one-loop corrections to its energy, whence the dressing factors can be reverse engineered. This can be done using the classical integrability properties the we described in section 3.1, and in particular the spectral curve. In the same way, one can obtain *finite-gap equations* that capture part of the semiclassical spectrum.

Tree-level S matrix

To expand the S matrix of section 3.3 at tree level it suffices to use the AFS phase (4.35) and to plug in (7.1). As discussed in section 6.3, we should take into account gauge terms of the form $e^{i a (p \delta D_q - q \delta D_p)}$.

In ref. [102] Hoare and Tseytlin (HT), among other things, wrote down a tree level S matrix for all the massive excitations of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, from a suitable truncation of the $\text{AdS}_5 \times \text{S}^5$ S matrix⁵. It is immediate to see that the scattering processes allowed in the HT S matrix are those that are non-zero in the one we derived in section 3.3. In particular, both S matrices are reflectionless, and both come from the tensor product of two $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ S matrices. In fact we can match our results both with the full S matrix (4.1) and with

³In the case of $\text{AdS}_5/\text{CFT}_4$, for instance, there exist a crossing-symmetric phase that reproduces the leading large- h behaviour, but fails to match the weak-coupling structure of the CFT_4 [65].

⁴At a late stage of the preparation of this thesis, ref. [108] appeared. There, fully-fledged one-loop calculations are performed in the near-BMN limit of $\text{AdS}_3/\text{CFT}_2$. Just like the ones that we will discuss more in detail hereafter, those results agree with our proposals for S matrix and dressing factors.

⁵Such a truncation is possible at tree level, but as we have seen the two matrices should be genuinely different at higher orders in a loop expansion.

each factor (4.8) of [102]. To do so, we identify the fundamental fields there with $\phi^{L,R}$ and $\psi^{L,R}$ as follows

$$|\phi^L\rangle = |\phi_+\rangle, \quad |\psi^L\rangle = |\psi_+\rangle, \quad |\phi^R\rangle = |\phi_-\rangle, \quad |\psi^R\rangle = -|\psi_-\rangle, \quad (7.16)$$

and (consistently with the tensor product structure) we identify the composite fields as

$$\begin{aligned} |\Phi_{++}^L\rangle &= |y_+\rangle, & |\Phi_{+-}^L\rangle &= -|\zeta_+\rangle, & |\Phi_{-+}^L\rangle &= |\chi_+\rangle, & |\Phi_{--}^L\rangle &= -|z_+\rangle, \\ |\Phi_{++}^R\rangle &= |y_-\rangle, & |\Phi_{+-}^R\rangle &= -|\zeta_-\rangle, & |\Phi_{-+}^R\rangle &= |\chi_-\rangle, & |\Phi_{--}^R\rangle &= -|z_-\rangle. \end{aligned} \quad (7.17)$$

If we observe that the HT S matrix is written in terms of \mathbb{R} rather than \mathbb{S} or $\tilde{\mathbb{S}}$ and correctly account for the fermion signs, we find perfect agreement with our results.

In ref. [100, 101] the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ and $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ S matrices have been investigated perturbatively by Rughoonauth, Sundin and Wulff (RSW) and by Sundin and Wulff (SW). At tree-level, the comparison is similar to the one above. Since in SW some computations are performed for the more general $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background, we should take the parameter $\alpha = 1$ everywhere to recover the T^4 background at tree level. Only a subset of the S matrix elements has been computed. Still, this probes the left-left and left-right diagonal and non-diagonal scattering. The necessary field identifications are

$$\begin{aligned} |\Phi_{++}^L\rangle &= |y_2\rangle, & |\Phi_{+-}^L\rangle &= |\chi_2\rangle, & |\Phi_{-+}^L\rangle &= |\chi_1\rangle, & |\Phi_{--}^L\rangle &= |y_1\rangle, \\ |\Phi_{++}^R\rangle &= |\bar{y}_2\rangle, & |\Phi_{+-}^R\rangle &= |\bar{\chi}_2\rangle, & |\Phi_{-+}^R\rangle &= |\bar{\chi}_1\rangle, & |\Phi_{--}^R\rangle &= |\bar{y}_1\rangle. \end{aligned} \quad (7.18)$$

Then, our results match perfectly the ones of RSW and SW.

One- and two-loop S-matrix elements

Sundin and Wulff [101] also computed certain one-loop elements in the near-flat-space limit. These correspond to certain elements $\mathcal{A}, \tilde{\mathcal{A}}, \mathcal{C}, \tilde{\mathcal{C}}$, some of which we encountered in chapter 4. They are defined by

$$\begin{aligned} \mathcal{A}_{pq} &= \langle \Phi_{++q}^L \Phi_{++p}^L | \tilde{\mathbb{S}}_{pq} | \Phi_{++p}^L \Phi_{++q}^L \rangle, & \tilde{\mathcal{A}}_{pq} &= \langle \Phi_{++q}^R \Phi_{++p}^L | \tilde{\mathbb{S}}_{pq} | \Phi_{++p}^L \Phi_{++q}^R \rangle, \\ \mathcal{C}_{pq} &= \langle \Phi_{--q}^L \Phi_{++p}^L | \tilde{\mathbb{S}}_{pq} | \Phi_{++p}^L \Phi_{--q}^L \rangle, & \tilde{\mathcal{C}}_{pq} &= \langle \Phi_{--q}^R \Phi_{++p}^L | \tilde{\mathbb{S}}_{pq} | \Phi_{++p}^L \Phi_{--q}^R \rangle. \end{aligned} \quad (7.19)$$

and read⁶

$$\begin{aligned} \mathcal{A}_{pq} &= 1 - \frac{i}{4h} \frac{p_- q_- (p_- + q_-)}{p_- - q_-} \\ &+ \frac{1}{32h^2} \frac{p_-^2 q_-^2}{q_-^2 - p_-^2} \left(\frac{i}{\pi} (p_- + q_-)^2 - \frac{2i}{\pi} \frac{q_- p_- (q_- + p_-)}{q_- - p_-} \log \frac{q_-}{p_-} - \frac{(q_- + p_-)^3}{q_- - p_-} \right), \end{aligned} \quad (7.20)$$

⁶The tree level expressions for \mathcal{C} and $\tilde{\mathcal{C}}$ were not given in [101], but were communicated privately to us by the authors.

$$\tilde{\mathcal{A}}_{pq} = 1 - \frac{i}{4h} \frac{p_- q_- (p_- - q_-)}{p_- + q_-} \quad (7.21)$$

$$- \frac{1}{32h^2} \frac{p_-^2 q_-^2}{q_-^2 - p_-^2} \left(\frac{i}{\pi} (p_- - q_-)^2 + \frac{2i}{\pi} \frac{q_- p_- (q_- - p_-)}{q_- + p_-} \log \frac{q_-}{p_-} + \frac{(q_-^2 + p_-^2)(q_- - p_-)}{q_- + p_-} \right),$$

$$\mathcal{C}_{pq} = 1 - \frac{i}{4h} p_- q_- \quad (7.22)$$

$$+ \frac{1}{32h^2} \frac{p_-^2 q_-^2}{q_-^2 - p_-^2} \left(\frac{i}{\pi} (p_- + q_-)^2 - \frac{2i}{\pi} \frac{q_- p_- (q_- + p_-)}{q_- - p_-} \log \frac{q_-}{p_-} - \frac{(q_-^2 + p_-^2)(q_- + p_-)}{q_- - p_-} \right),$$

$$\tilde{\mathcal{C}}_{pq} = 1 - \frac{i}{4h} p_- q_- \quad (7.23)$$

$$- \frac{1}{32h^2} \frac{p_-^2 q_-^2}{q_-^2 - p_-^2} \left(\frac{i}{\pi} (p_- - q_-)^2 + \frac{2i}{\pi} \frac{q_- p_- (q_- - p_-)}{q_- + p_-} \log \frac{q_-}{p_-} + (q_-^2 - p_-^2) \right),$$

where in contrast with [101] we explicitly used the coupling constant h as a loop-counting parameter. A first nontrivial requirement of our construction is that these elements satisfy the crossing equations of chapter 4, that read simply

$$\mathcal{A}_{pq} \tilde{\mathcal{A}}_{p^c q^c} = 1, \quad \mathcal{C}_{p^c q^c} \tilde{\mathcal{C}}_{pq} = 1. \quad (7.24)$$

It is easy to check that this is actually the case, observing that in light-cone coordinate crossing the second variable $q \rightarrow q^c$ amounts to $q_- \rightarrow q_-^c = -q_-$, and taking everywhere the upper branch of the logarithm [S2]. Using the near-flat-space expansion of the dressing phases (7.14) we can check that we perfectly match these elements. This is a first direct validation of our dressing factors.

A different approach to obtain higher-loop expressions is to use unitarity techniques (akin to the optical theorem) and exploit two-dimensional kinematics [106, 107]. In particular, Engelund, McKeown and Roiban (EKR) have applied such techniques to $\text{AdS}^3 \times \text{S}^3 \times \text{T}^4$ in ref. [107], obtaining near-BMN one- and two-loop results for the diagonal S-matrix elements. However, in such an approach only the *logarithmic* part of the elements is predicted unambiguously. Moreover, an l -loop calculation requires as input the matrix structure of \mathbb{S} at $(l-1)$ -loop. At one-loop one can rely on the tree-level perturbative results above, but the two-loop calculation requires using the symmetries of the model to constrain \mathbb{S} as we did. Despite not being completely independent, the two-loop result is a very non-trivial consistency check of the S matrix and dressing factors. Once again, our proposal satisfies it.

Yet another approach to computing the dressing factors is to obtain them from the semiclassical energy shifts to classical NSLM solutions. This approach was pioneered in [162, 163] in the case of $\text{AdS}_5 \times \text{S}^5$ and was later applied to $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ as well [103–105, 150]. In ref. [103], Beccaria, Levkovic-Maslyuk, Macorini and Tseytlin (BLMMT) have computed the one-loop dressing phase. Their calculation pre-dated our proposal [S2] based on crossing symmetry. Up to different normalisation due to the choice of the expansion parameters resulting in a factor of 4π , their results are given in terms of the

expansion (7.3) by

$$\begin{aligned} c_{\text{BLMMT } r,s}^{(1)} &= c_{r,s}^{(1)}, & \tilde{c}_{\text{BLMMT } r,s}^{(1)} &= \tilde{c}_{r,s}^{(1)}, \\ c_{\text{BLMMT } 1,s}^{(1)} &= 2c_{1,s}^{(1)}, & \tilde{c}_{\text{BLMMT } 1,s}^{(1)} &= 2\tilde{c}_{1,s}^{(1)}. \end{aligned} \quad (7.25)$$

The coefficients $c_{r,s}^{(1)}$ and $\tilde{c}_{r,s}^{(1)}$ do not match our proposal when $r = 1$. In fact, it is not hard to see [S3] that the resulting phases do not even satisfy our crossing equations.

A resolution of this discrepancy was proposed by Abbott [105], who highlighted a subtlety in the definition of the charges \mathcal{Q}_r in the semiclassical derivation, as in refs. [164, 165]. This results in a factor of $\frac{1}{2}$ in the normalisation of \mathcal{Q}_1 , and does not affect any higher charge. Taking this into account, Abbot showed that the semiclassical calculation perfectly matches our proposal for the dressing factors.

Finite-gap equations

The spectral curve $\Gamma(x)$ (3.5) encodes all the information about the conserved charges of the classical NLSM. The same is true for its eigenvalues $\{\gamma_j(x)\}$, or equivalently the “quasi-momenta” $\{p_j(x)\}$, given by $\gamma_j(x) = e^{ip_j(x)}$. In the case where $\Gamma(x)$ is an algebraic curve, *i.e.* it has finite genus, it is possible to write down integral equations for the discontinuities at its cuts to describe $\Gamma(x)$ —the *finite-gap equations*. By solving those equations we can find the classical energy of a solution having a given set of charges. The finite gap equations for $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, were proposed in the seminal paper [50] by Babichenko, Stefański and Zarembo, and we expect that they describe a suitable classical limit of our spectrum.

It is natural to expect that the finite-gap equations are a limit of the Bethe equations where the magnon momenta condense to form cuts [24]. We introduce the densities⁷

$$\rho_j(x) = \sum_{k=1}^{K_j} \frac{x^2}{x^2 - 1} \delta(x - x_{j,k}), \quad j = 1, 2, 3, \tilde{1}, \tilde{2}, \tilde{3}, \quad (7.26)$$

and take the excitation numbers to be large, $K_j \gg 1$. By making use of the expansion (7.2), we find the following finite gap-equations

$$2\pi n_1 = - \int \frac{\rho_2(y)}{x-y} dy - \int \frac{\rho_{\tilde{2}}(y)}{x-1/y} \frac{dy}{y^2} \quad (7.27)$$

$$\begin{aligned} 2\pi n_2 &= - \frac{x}{x^2-1} 2\pi \mathcal{E} - \int \frac{\rho_1(y)}{x-y} dy + 2 \oint \frac{\rho_2(y)}{x-y} dy - \int \frac{\rho_3(y)}{x-y} dy \\ &+ \int \frac{\rho_{\tilde{1}}(y)}{x-1/y} \frac{dy}{y^2} + \int \frac{\rho_{\tilde{3}}(y)}{x-1/y} \frac{dy}{y^2} + \frac{1}{x^2-1} \mathcal{M}, \end{aligned} \quad (7.28)$$

$$2\pi n_3 = - \int \frac{\rho_2(y)}{x-y} dy - \int \frac{\rho_{\tilde{2}}(y)}{x-1/y} \frac{dy}{y^2} \quad (7.29)$$

⁷Here we find it convenient to work in terms of excitation numbers K_j rather than $M_{jL,R}^{\text{I,II}}$. The relation between the two sets is given in eqs. (6.76–6.76).

$$2\pi n_{\bar{1}} = \int \frac{\rho_2(y)}{x-1/y} \frac{dy}{y^2} + \int \frac{\rho_{\bar{2}}(y)}{x-y} dy \quad (7.30)$$

$$2\pi n_{\bar{2}} = -\frac{x}{x^2-1} 2\pi \mathcal{E} - \int \frac{\rho_1(y)}{x-1/y} \frac{dy}{y^2} - \int \frac{\rho_3(y)}{x-1/y} \frac{dy}{y^2} \\ + \int \frac{\rho_{\bar{1}}(y)}{x-y} dy - 2 \int \frac{\rho_{\bar{2}}(y)}{x-y} dy + \int \frac{\rho_{\bar{3}}(y)}{x-y} dy + \frac{1}{x^2-1} \mathcal{M}, \quad (7.31)$$

$$2\pi n_{\bar{3}} = \int \frac{\rho_2(y)}{x-1/y} \frac{dy}{y^2} + \int \frac{\rho_{\bar{2}}(y)}{x-y} dy, \quad (7.32)$$

where \int denotes the principal-value integral. Here \mathcal{E} corresponds to the residue of the quasi-momentum and it is given by

$$\mathcal{E} = \frac{1}{2\pi} (L - \epsilon_1 + 2\epsilon_2 - \epsilon_3 + \epsilon_{\bar{1}} + \epsilon_{\bar{3}}), \quad (7.33)$$

where

$$\epsilon_j = \int \frac{\rho_j(x)}{x^2} dx. \quad (7.34)$$

The quantity \mathcal{M} has the meaning of winding of the corresponding solutions and it is given by

$$\mathcal{M} = \mathcal{P}_1 + \mathcal{P}_3 - \mathcal{P}_{\bar{1}} + 2\mathcal{P}_{\bar{2}} - \mathcal{P}_{\bar{3}} = \mathcal{P}_1 - \mathcal{P}_2 + \mathcal{P}_3 - \mathcal{P}_{\bar{1}} + \mathcal{P}_{\bar{2}} - \mathcal{P}_{\bar{3}}, \quad (7.35)$$

where

$$\mathcal{P}_j = \int \frac{\rho_j(x)}{x} dx. \quad (7.36)$$

The last equality in (7.35) is possible thanks to the level matching condition that reads

$$\mathcal{P}_2 + \mathcal{P}_{\bar{2}} = 0. \quad (7.37)$$

The finite-gap equations that we derived are apparently different but equivalent to the ones in [50]. In fact, if we repeated the same construction performed there (see also refs. [98, 99]) with a different choice of the grading, such as (6.75), we would have found precisely (7.27–7.32). While for our Bethe equations we had only four choices of grading, all of them involving different Cartan matrices in the left and right sectors, at the classical level there are more choices—one can change grading to each copy of $\mathfrak{psu}(1,1|2)$ independently. We can then conclude that our proposal is compatible with semiclassical integrability calculations too.

7.3 Chapter summary

In this chapter we have compared our results, that stemmed from the assumption of quantum integrability, with independent ones. This comparison is a check of quantum integrability, but is also an important test for the dressing factors, that were not completely fixed by crossing invariance. All of the independent calculations that we checked were

performed on the *string theory side*, that is at large values of h . Our proposal, including the dressing factors, reproduces all of them. This non-trivial matching holds at one-loop and puts strong requirements on the two-loop S matrix elements too. Based on that, we are confident to say that we expect a quantum-integrable dynamics for the massive modes of the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ superstring, and we expect the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ symmetry and a discrete left-right symmetry to play a crucial role in it.

Still, it is highly desirable to test even further our results. In particular, we only have limited understanding of the weakly-coupled $h \ll 1$ regime, and of how $h(\lambda)$ scales in terms of the CFT coupling λ . Moreover, as we discussed in section 7.1, the dressing factors have a different structure compared to the ones of $\text{AdS}_5/\text{CFT}_4$ or $\text{AdS}_4/\text{CFT}_3$, that yields new puzzling features at weak coupling.

Let us conclude this chapter by reviewing some different proposals for the S matrix and dressing factors of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$. An early proposal due to David and Sahoo [122, 166] was derived from a study of the giant magnons and their symmetry properties. That proposal also relied on a $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ symmetry, but focused exclusively on the left-left sector. As a consequence, the presence of a substantially different LR S matrix was not accounted for, and the crossing invariance proposed there was different from the one we discussed, and resulted in a single dressing phase—a result incompatible with later calculations [103, 105].

In ref. [50], Babichenko, Stefański and Zarembo (BSZ) conjectured a set of all-loop Bethe equations from the finite-gap equations. Those presented two discrepancies with respect to the ones we proposed. Firstly, as discussed in section 7.2, they were written in a different grading. Secondly, they overlooked some terms⁸ that are subleading in the finite-gap limit but are important for unitarity of the all-loop S matrix.

After that, in ref. [167], Ahn and Bombardelli (AB) proposed an all-loop S matrix that reproduced the BSZ conjecture for the Bethe equations. As discussed more in detail in ref. [S5], such an S matrix was not invariant under the *extended* symmetry $\mathfrak{su}(1|1)_{\text{c.e.}}^2$, but only under $\mathfrak{su}(1|1)^2$, and its form differs from ours in particular in the LR sector. The proposal of AB can be ruled out by checking what are the non-vanishing processes in the LR sector against perturbative calculations.

⁸Namely, the symmetric phases that appear in the couplings between left and right sectors in front of $\tilde{\sigma}_{pq}^2$, see eqs. (6.64) and (6.67). These are required for the unitarity of the S matrix.

8 Recent developments and new directions

Now that we have a good handle on the simplest aspects of $\text{AdS}_3/\text{CFT}_2$ integrability, in this chapter we will describe some directions that are now being investigated, which go beyond those aspects. The most natural one in order to complete our treatment is the inclusion of massless excitations in the integrability description.

8.1 Massless modes in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$

We have seen in section 2.1 that the bosonic string spectrum on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ features four fundamental massless excitations. By supersymmetry, these must be supplemented by four fermionic ones. Extending our S-matrix treatment to such excitations appears problematic. Massless excitations are characterised by the scaling of their dispersion relation at small momentum, of the form¹

$$\omega(p)^2 = c^2 p^2 + O(p^4), \quad (8.1)$$

which results in $\omega(p)$ being a non-analytic function of p around zero. Consequently it is natural to distinguish between left- and right-movers *on the worldsheet*, having respectively $p > 0$ and $p < 0$, and energy

$$E_{\text{left}} = +cp + O(p^3), \quad E_{\text{right}} = -cp + O(p^3). \quad (8.2)$$

This should not be confused with the notion of left- and right-movers *in the dual CFT*, *i.e.* the “L” and “R” labels that were ubiquitous in the previous chapters.

In the familiar relativistic case, higher orders in p are absent, and the dispersion relation is linear. Therefore, the group velocity of a wave-packet is

$$v_{\text{rel}} = \frac{\partial \omega}{\partial p} = \pm c, \quad (8.3)$$

i.e., massless relativistic particles move at the speed of light. Particles with the same worldsheet chirality then cannot scatter, regardless of the value of their momentum. Still,

¹More general definitions of massless (quasi)particles may be given, but this one will suffice for our purposes.




Figure 8.1: The action of the supercharges on the massive excitations is supplemented by \mathbb{K}^α from $\mathfrak{su}(2)_K$, acting on the fermions. Note that for clarity we only depict the action of some of the supercharges, *i.e.* the ones that remain non-trivial on-shell.

a formal treatment of relativistic massless theories in terms of factorised scattering is possible [168–171]. To this end it is however necessary to introduce appropriate rapidity variables and take suitable limits.

In the case of our interest, however, factorised scattering appears to be *simpler* than in the relativistic case. In fact, as argued in [S1], the dispersion relation of massless worldsheet particles should take the form

$$\omega(p)^2 = 4h^2 \sin^2\left(\frac{p}{2}\right), \quad (8.4)$$

from which we find the group velocity

$$v_{\text{non-rel}} = \pm h \cos\left(\frac{p}{2}\right). \quad (8.5)$$

We see then that massless excitations that have different momenta also have different velocities, so that we can expect them to scatter in a way similar to massive excitations. Starting from this intuition, in ref. [S1] the fundamental S matrix of massive and massless excitations was computed. The approach is similar to the one used in chapters 2 and 3, relying on the off-shell symmetry algebra of the theory. As we have mentioned in chapter 2, the coset formalism is not suitable for treating massive and massless excitations at the same time. The issue is that the coset κ -gauge fixing—whereby all physical fermions are taken to live in the coset—is not compatible with light-cone gauge. To overcome this, it is possible to use the Green-Schwarz superstring formalism, taking a κ gauge where the massless fermions live on the torus. Then one can compute the Noether charges and Killing spinors of the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ geometry to find the off-shell symmetries.

One then finds that the $\mathfrak{psu}(1|1)^4$ is supplemented by a bosonic $\mathfrak{o}(4)$ symmetry coming from the torus, with in particular an $\mathfrak{so}(4) = \mathfrak{su}(2)_K \oplus \mathfrak{su}(2)_Y$ symmetry. This may appear surprising, as one would naïvely expect only $\mathfrak{u}(1)^4$ isometries from the torus. However, the presence of periodic boundary conditions on T^4 is probed only by winding modes, that in the decompactification limit, where the S matrix is defined, decouple. The




Figure 8.2: The massless excitations sit in two $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ modules, with fermionic highest weight states. The two modules transform as a doublet under $\mathfrak{su}(2)_Y$, and additionally the bosons $T^{\alpha a}$ also carry a Greek index of $\mathfrak{su}(2)_K$.

massive fundamental excitations are charged under some of the torus isometries. In fact, the fermions $\Phi_{\pm\mp}^L$ and $\Phi_{\pm\mp}^R$ form two doublets under $\mathfrak{su}(2)_K$, as depicted in figure 8.1. The massless excitations are also charged under $\mathfrak{so}(4)$, and we label them accordingly. Using Greek letters for fundamental and antifundamental $\mathfrak{su}(2)_K$ indices, and Latin ones for $\mathfrak{su}(2)_Y$, we can write the eight massless excitations as

$$\text{bosons: } T^{\alpha a}, \quad \text{fermions: } \chi^a, \tilde{\chi}^a, \quad (8.6)$$

see figure 8.2. The fermions χ^a and $\tilde{\chi}^a$ are distinguished by having opposite eigenvalues under the bosonic charge \mathbb{N} of eq. (2.86). The $\mathfrak{psu}(1|1)^4$ supercharges are also charged under $\mathfrak{su}(2)_K$, forming four doublets

$$\mathbb{Q}^{\alpha L}, \quad \bar{\mathbb{Q}}_{\alpha}{}^L, \quad \mathbb{Q}_{\alpha}{}^R, \quad \bar{\mathbb{Q}}^{\alpha R}. \quad (8.7)$$

The final ingredient for finding the S matrix is understanding if and how the non-trivial central extension appears when going from the on-shell to the off-shell symmetry algebra. In the coset formulation we saw this easily because the light-cone coordinate x_- was² neatly packaged into $\Lambda(x_{\pm})$ as an exponential. Without reference to the coset representatives, the same terms naturally arise in the Green-Schwarz formulation, by requiring that the fermions are not charged under the light-cone isometries [S1], see also *e.g.* ref. [172]. In fact one can see that the central extension takes the same form as in the massive sector. With these ingredients it is then straightforward to find the $2 \rightarrow 2$ S matrix, which for massless excitations is expressed in terms of modified Zukhovskiy variables satisfying

$$\frac{x_p^+}{x_p^-} = e^{ip}, \quad x_p^+ x_p^- = 1, \quad x_p^{\pm} = e^{\pm \frac{i}{2}p} \text{sign}\left(\sin \frac{p}{2}\right), \quad (8.8)$$

whereby we get the dispersion relation (8.4) by the usual formula (3.51). Unsurprisingly, since we are dealing with massless particles, a non-analyticity arises at $p = 0$, whence a

²Recall that x_- is related to the total worldsheet momentum by $\mathbb{P} = \int d\sigma x'_-$.

splitting into left- and right-movers of the worldsheet can be introduced. It is also interesting to observe that on symmetry grounds, unless the $\mathfrak{so}(4)$ invariance of the S matrix is broken, quantum effects cannot give a mass the massless particles—unlike what happens in other integrable theories such as the Gross-Neveu model [173, 174].

The worldsheet S matrix again factors in a tensor-product structure, and depends on five distinct dressing phases,

$$\mathcal{S}_{pq}^{\bullet\bullet}, \quad \widetilde{\mathcal{S}}_{pq}^{\bullet\bullet}, \quad \mathcal{S}_{pq}^{\bullet\circ}, \quad \mathcal{S}_{pq}^{\circ\bullet}, \quad \mathcal{S}_{pq}^{\circ\circ}, \quad (8.9)$$

where the first two phases are the massive-massive ones that we already studied in chapter 4, while the remaining three pertain to the massive-massless, massless-massive and massless-massless sectors, respectively. The requirement of crossing symmetry for all these phases can be written down straightforwardly, but a proposal for the latter three and a study of their properties is still lacking. To this end it may be useful to first obtain some insight in their perturbative form by performing worldsheet calculations, following ref. [101] and perhaps taking advantage of the recent advances in unitarity techniques [106, 107].

The proposal of the all-loop S matrix for all fundamental excitations is a crucial step forward in $\text{AdS}_3/\text{CFT}_2$ integrability, but should be supplemented by several related investigations. Among these, finding the Bethe-Yang equations and studying how the full $\mathcal{N} = (4, 4)$ infinite-dimensional symmetry is realised at the level of the asymptotic spectrum is probably the most important one. Once that has been understood, it should also be possible to formulate a string hypothesis and mirror TBA equations for this model, and to start to tackle finite-size physics, “spectroscopy”, comparison with other approaches, and many other issues, some of which we will mention in the conclusions. We foresee many of these studies to be completed in the near future, rapidly advancing our understanding of this model.

8.2 Integrability for the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background

There is another AdS_3 background (in fact, a family of backgrounds) that preserves the maximal amount of supersymmetry, *i.e.* 16 supercharges. It is given by $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ provided that the curvature radii of the two spheres R_+ and R_- satisfy

$$\frac{1}{R_+^2} + \frac{1}{R_-^2} = \frac{1}{R_{\text{AdS}}^2}, \quad (8.10)$$

where R_{AdS} is the radius of AdS_3 . This gives an one-parameter family that can be labelled by α

$$\alpha = \frac{R_{\text{AdS}}^2}{R_+^2} = 1 - \frac{R_{\text{AdS}}^2}{R_-^2}, \quad (8.11)$$



Figure 8.3: Two Dynkin diagrams for $\mathfrak{d}(2, 1; \alpha)$. Diagram (b) corresponds to a completely fermionic grading.

with $0 < \alpha < 1$. Clearly, up to exchanging the role of the two spheres, we can restrict to $\frac{1}{2} \leq \alpha < 1$, where the point $\alpha = \frac{1}{2}$ is special in that the two spheres become identical there. Another interesting configuration is the one where $\alpha \rightarrow 1$, when R_+ equates the AdS radius and R_- blows up. In this limit the second sphere becomes flat, and decompactifies to \mathbb{R}^3 . Up to the fact that \mathbb{R}^3 should be compactified back to T^3 , which can be done in several ways, we have basically obtained again the $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ background. However, since this limit appears quite delicate, let us for the moment take $\frac{1}{2} \leq \alpha < 1$ and come back to it later.

Once again the background has flat directions that give rise to massless excitations in its spectrum in light-cone gauge. Instead of the four massless modes from T^4 , here we have two: one resides on the circle S^1 , and one is shared by the two spheres³. Once again, up to discarding the massless excitations, the string action can be written in terms of a coset [50, 98, 99]

$$\frac{D(2, 1; \alpha) \times D(2, 1; \alpha)}{SU(1, 1) \times SU(2) \times SU(2)}, \quad (8.12)$$

where $D(2, 1; \alpha)$ is the supergroup corresponding to the exceptional basic simple Lie superalgebra $\mathfrak{d}(2, 1; \alpha)$, see refs. [175, 176]. Here $\frac{1}{2} \leq \alpha < 1$ is precisely the curvature ratio of eq. (8.11), and the case $\alpha = \frac{1}{2}$ gives $\mathfrak{d}(2, 1; \frac{1}{2}) \cong \mathfrak{osp}(4|2)$. The superalgebra $\mathfrak{d}(2, 1; \alpha)$ can be written in two inequivalent gradings. In figure 8.3 we illustrate the relative Dynkin diagrams.

Classical and perturbative aspects

Clearly the coset (8.12) can be equipped with a \mathbb{Z}_4 automorphisms, whence classical integrability follows, in the sense that the equations of motion admit the Lax representation that we briefly discussed in section 3.1. However, this as usual requires some care in dealing with the massless modes, that are not automatically present in the coset construction

³To preserve as much supersymmetry as possible, the light-cone geodesic runs through the time coordinate in AdS_3 and both of the spheres. The massless excitation comes from a coordinate on the spheres that is orthogonal to such geodesic.

Mass	$m = \alpha$	$m = 1$	$m = 1 - \alpha$	$m = 0$
Bosons	ϕ^{1L}, ϕ^{1R}	Φ^{2L}, Φ^{2R}	ϕ^{3L}, ϕ^{3R}	φ^0, φ^4

Table 8.1: Fundamental bosonic excitations of the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ in light-cone gauge. Fermions follow by supersymmetry. The massive modes are labelled (left-right) by the sign of a suitable target-space angular momentum, which can be interpreted as the chirality in the dual CFT_2 —much like in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$. The massless excitation φ^0 is shared between the two spheres, while φ^4 lives on the circle S^1 .

and must be added by hand. This was shown to be possible by Babichenko, Stefański and Zarembo in ref. [50] in a specific κ -gauge fixing, and then extended by Sundin and Wulff [51] to an arbitrary κ gauge.

A perturbative investigation of the spectrum in light-cone gauge yields again 8 + 8 bosonic and fermionic fundamental excitations. They come in eight supersymmetric doublets, with four different masses. We collect them in table 8.1. There are 4 + 4 “light” excitations ($\phi^j | \theta^j$) of mass $0 < m < 1$ and 2 + 2 “heavy” ones ($\Phi^j | \Theta^j$) of mass $m = 1$, plus the aforementioned massless modes ($\varphi^j | \vartheta^j$).

In ref. [50], besides establishing classical integrability for this background, the finite-gap equations for massive excitations were written down. In that context, it naturally appears that the heavy modes are actually composite, consisting of two light excitations of mass α and $1 - \alpha$. From the point of view of the S matrix, this is reflected in the presence of a tree-level light-light-heavy vertex.

In order to understand whether the heavy mode is truly composite one should analyse its (renormalised) two-point function. If this displays a pole at mass $m = 1$, then the heavy mode should be treated as a fundamental particle, and included in the asymptotic states. If the pole is replaced by a branch cut, then the mode should be treated as composite. This issue was investigated at one loop in ref. [51], where at least in the simpler case when $\alpha = \frac{1}{2}$ it was established that the latter scenario occurs.

Quantum integrability

The investigation of integrability beyond the classical limit was initiated by Ohlsson Sax and Stefański in ref. [112], by constructing a weakly-coupled spin chain with $\mathfrak{d}(2, 1; \alpha)^2$ symmetry. This corresponds to the strongly-coupled regime in the NLSM, *i.e.* to the opposite of what was considered in ref. [50]. Even at weak coupling, the spin chain appears substantially more complicated than the $\mathfrak{psu}(1, 1|2)^2$ one, which we discussed in chapter 5. The $\mathfrak{d}(2, 1; \alpha)^2$ spin chain is given by the direct product of two *alternating* spin chains. Each alternating chain has its odd and even sites in the short (infinite-dimensional) $\mathfrak{d}(2, 1; \alpha)$ representations $(\frac{\alpha}{2}; \frac{1}{2}; 0)$ and $(\frac{1-\alpha}{2}; 0; \frac{1}{2})$, respectively. At weak




Figure 8.4: Pictorial representation of the $\mathfrak{d}(2,1;\alpha)^2$ alternating spin chains. We can think of it as composed of two (left and right) alternating chain. For each of these, odd (cyan) sites are in the $(\frac{\alpha}{2}; \frac{1}{2}; 0)$ representation of $\mathfrak{d}(2,1;\alpha)$, while even (pink) sites are in the $(\frac{1-\alpha}{2}; \frac{1}{2}; 0)$ representation. The dotted squares represent the combination of sites which we identify as fundamental excitations of the whole $\mathfrak{d}(2,1;\alpha)^2$ chain.

coupling, the alternating spin chains are decoupled, and correspond to left- and right-moving excitations, as it happened for $\mathfrak{psu}(1,1|2)^2$. The setup is illustrated in figure 8.4.

In ref. [S5] this picture was used to study the all-loop integrability properties of the chain. As we did in chapter 5, the first step of this analysis was to pick a vacuum for the chain that preserved as much supersymmetry as possible, and study fundamental excitations on top of it. This resulted in $4 + 4$ excitations, the bosonic ones being

$$|\phi^{1L}\rangle, \quad |\phi^{1R}\rangle, \quad |\phi^{3L}\rangle, \quad |\phi^{3R}\rangle. \quad (8.13)$$

These are precisely the light modes of table 8.1. Each of these excitations forms a doublet with its fermionic partner, and transforms under $\mathfrak{su}(1|1)_{\text{c.e.}}^2$, that is, under the very same algebra that we have analysed at length in the previous chapters. In fact, should we consider only $|\phi^{1L}\rangle$ and $|\phi^{1R}\rangle$, we would have the same algebra and representation as in section 3.2, up to rescaling the mass. This rescaling can be easily done in terms of modified Zhukovski variables

$$\frac{x_{j,p}^+}{x_{j,p}^-} = e^{ip}, \quad x_{j,p}^+ + \frac{1}{x_{j,p}^+} - x_{j,p}^- - \frac{1}{x_{j,p}^-} = \frac{2im_j}{h}, \quad (8.14)$$

with $m_1 = \alpha$ for $|\phi^{1L,R}\rangle$ and $m_3 = 1 - \alpha$ for $|\phi^{3L,R}\rangle$. Then the dispersion relation reads

$$E = \frac{h}{2i} \left(x_{j,p}^+ - \frac{1}{x_{j,p}^+} - x_{j,p}^- + \frac{1}{x_{j,p}^-} \right) = \sqrt{m_j^2 + 4h^2 \sin^2 \left(\frac{p}{2} \right)}, \quad (8.15)$$

so that a large- h expansion⁴ reproduces the spectrum of table 8.1. Furthermore, for the same reasons as in the previous chapters, it is natural to take the spin chain to have a discrete left-right symmetry.

Despite the many similarities, let us point out two important differences between this spin chain and the $\mathfrak{psu}(1,1|2)^2$ one. Firstly, the present symmetry algebra $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ is *half* of the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ algebra of chapter 5. What earlier was only one tensor product

⁴Scaling as usual the momentum as $1/h$.

factor, which we introduced for convenience, is now the full, physical symmetry algebra of the spin-chain excitations. Secondly, even if when we restrict to a subset of particles of a given mass we just need to rescale the Zhukovski variables (8.14), the presence of different masses introduces novel features. In particular, scattering processes involving particles of different masses are new to this spin chain.

Using this spin-chain picture, an all-loop S matrix was worked out in ref. [S5]. We can summarise it as a block matrix

$$\mathbb{S} = \left(\begin{array}{c|c} \mathbb{S}_{11} & \mathbb{S}_{31} \\ \hline \mathbb{S}_{13} & \mathbb{S}_{33} \end{array} \right), \quad (8.16)$$

where \mathbb{S}_{jj} is given by rewriting the $\mathfrak{su}(1|1)^2$ S matrix of chapter 3 using the Zhukovski variables (8.14), and \mathbb{S}_{jk} with $j \neq k$ scatters particle of different masses. One finds that the mass quantum number is always transmitted, *e.g.*

$$|\mathcal{X}_{j,p} \mathcal{Y}_{k,q}\rangle \longrightarrow \# |\tilde{\mathcal{Y}}_{k,q} \tilde{\mathcal{X}}_{j,p}\rangle, \quad j \neq k, \quad (8.17)$$

where $\mathcal{X}_{j,p}, \mathcal{Y}_{k,q}$ are magnon excitations of definite mass and momentum and $\tilde{\mathcal{X}}_{j,p}, \tilde{\mathcal{Y}}_{k,q}$ are their scattering products. We can further subdivide the mixed-mass scattering by chirality *e.g.*

$$\mathbb{S}_{13} = \left(\begin{array}{c|c} \mathbb{S}_{13}^{\text{LL}} & \mathbb{S}_{13}^{\text{RL}} \\ \hline \mathbb{S}_{13}^{\text{LR}} & \mathbb{S}_{13}^{\text{RR}} \end{array} \right). \quad (8.18)$$

We then find that each block is given by the corresponding one from chapter 3, up to rescaling each of the two Zhukovski variables independently.

The resulting S matrix satisfies the Yang-Baxter equation and, after imposing unitarity and left-right symmetry, depends on *four* dressing factors

$$\mathcal{S}_{jj}, \quad \tilde{\mathcal{F}}_{jj}, \quad \mathcal{S}_{jk}, \quad \tilde{\mathcal{F}}_{jk}. \quad (8.19)$$

In reducing the number of the dressing factors to four, we assumed that the mass-dependence in *e.g.* \mathcal{S}_{11} and \mathcal{S}_{33} comes from the same functional dependence on m . Crossing symmetry relates the first two of dressing factors among themselves, and the last two among themselves. So far, there is no all-loop proposal for these dressing factors. Even if there have been some tree-level and one-loop worldsheet perturbative calculations that validate⁵ the proposed all-loop S matrix [50, 100, 101, 150], the understanding of this model is quite limited and calculations are more involved due to the presence of the additional parameter α .

⁵There may arise some confusion concerning ref. [101], where the one-loop matrix elements found for the limiting case $\alpha = 1$ are compared with the all-loop S matrix at $\frac{1}{2} \leq \alpha < 1$ and a mismatch is found. This is because, as we have seen, the S matrix radically changes in that limit, and the notion of fundamental particles is different at $\alpha < 1$ and $\alpha = 1$, see also the next subsection. Taking this into account, we can match the calculations of ref. [101] with *both* the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ and $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ S matrices, for appropriate values of α .




Figure 8.5: The Dynkin diagram for $\mathfrak{d}(2,1;\alpha)^2$ in mixed grading (see figure 8.3), with the various interaction terms of the Bethe ansatz indicated. Note how now, besides dressing phases that couple nodes of the same mass (in blue), there appear phases coupling nodes of mass α with ones of mass $1 - \alpha$ (in red).

From the proposed S matrix, the all-loop Bethe ansatz equations have been written in [S4], in a way much similar to what we did in section 6.2. Once again, the equations are written in a mixed grading for the two copies of $\mathfrak{d}(2,1;\alpha)$, and feature slightly different phases with respect to the early proposals that were reverse-engineered from the finite-gap equations, see refs. [50,112]. We schematically represent them in figure 8.5.

Massless modes and $\alpha \rightarrow 1$ limit

Once again, incorporating the massless excitations in the integrability description does not appear to be straightforward. Very recently, some progress has been made at the level of the finite-gap equations [111]. The original set of finite-gap equations did not include any of the massless modes of the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background, not even the one shared by the two spheres, which should be captured by the supercoset (8.12). In ref. [111] Lloyd and Stefański showed that such a mode can be accounted for by implementing the Virasoro constraints as suitable conditions on the residues at the poles of the quasimomenta—in particular, as weaker conditions than the ones assumed in the earlier literature. Therefore, accounting for the massless mode does not result in additional cuts in the algebraic curve description, but rather in a modification of the residues of the quasimomenta, which contribute to the determination of the energy of the state. Such a prescription can be also employed to accommodate the massless mode of S^1 , or the four modes of T^4 . This is a promising advance, which hopefully will help including massless modes in general all-loop integrability scenarios.

Another interesting direction is to exploit this background to *generate* massless modes. This can be done by taking the limit in which one of the spheres becomes flat, $\alpha \rightarrow 1$. At the level of $\mathfrak{d}(2,1;\alpha)$ this corresponds to a contraction of one of the $\mathfrak{su}(2)$ bosonic

subalgebras. The limit is quite subtle, because this eliminates one of the simple roots, see figure 8.3a, and introduces a novel fermionic one. At the level of the S matrix, the limit is non-trivial too. We argued that the mass-one excitations $\Phi^{2L,R}$ are composite in the case of $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$. In $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, there are four *fundamental* bosonic modes of mass one, two of which can be obtained from the one of mass α . The remaining two must be the once-composite modes $\Phi^{2L,R}$.

All this makes the comparison not straightforward. However, in the limit of the weakly-coupled spin chain some progress could be made [110]. There it was shown that the \mathbb{R} matrix is regular in the limit $\alpha \rightarrow 1$, where its representations can be studied. The $\mathfrak{d}(2,1;\alpha)$ alternating spin chain does not quite go to the $\mathfrak{psu}(1,1|2)$ homogeneous one. While the even sites are given by the $(\frac{1}{2}, \frac{1}{2})$ $\mathfrak{psu}(1,1|2)$ module (see section 5.1), the odd ones are given by $0 \oplus 0 \oplus (\frac{1}{2}, \frac{1}{2})$, *i.e.* also feature two singlets. These are related to the massless modes, and yield a degenerate vacuum. By writing down the corresponding Bethe ansatz equations and carefully studying their degeneracies it is possible to show that the degeneration is compatible with the chiral ring of the dual CFT—this is however quite subtle and requires considerations that go beyond the weakly-coupled regime.

8.3 Mixed-flux backgrounds

Up to this point, our focus has been on integrable non-linear σ models corresponding to superstring backgrounds supported by a pure Ramond-Ramond flux. This is in close analogy with the prototypical integrability example of the $\text{AdS}_5 \times \text{S}^5$ superstring. As we briefly mentioned in the introduction, however, AdS_3 backgrounds are special in that they can also be supported by a Neveu-Schwarz-Neveu-Schwarz flux, or by a combination of NSNS and RR fluxes. The pure NSNS theory is in a way simpler to quantise and, at least in the case of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$, is well-understood [41–44]⁶. Such a case can in fact be understood as a supersymmetric extension of the $SL(2) \times SU(2)$ Wess-Zumino-Witten (WZW) model, whence the free-string spectrum can be found by CFT techniques [42]. The mixed-flux case has also been considered [177, 178], but until recently the solution of the relative spectral problem appeared to be out of reach.

Classical integrability

In ref. [52], Cagnazzo and Zarembo investigated such mixed flux backgrounds. In general, their Lagrangian takes the form

$$\mathcal{L} = -\frac{\hbar}{2} \left(\gamma^{\alpha\beta} G_{\mu\nu}(X) + q \varepsilon^{\alpha\beta} B_{\mu\nu}(X) \right) \partial_\alpha X^\mu \partial_\beta X^\nu + \text{fermions}. \quad (8.20)$$

⁶The $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background has also been studied, but remains more obscure [48, 49].

In a coset formulation such as the one of chapter 2, this can be written as

$$S = -\frac{\hbar}{2} \left(\int d^2\sigma \gamma^{\alpha\beta} \text{str}(A_\alpha^{(2)} A_\beta^{(2)}) + \frac{q}{3} \int_{\mathcal{B}} d^3\sigma \varepsilon^{\alpha\beta\gamma} \text{str}(A_\alpha^{(2)} A_\beta^{(2)} A_\gamma^{(2)}) \right) + \text{fermions}. \quad (8.21)$$

Note that the last term is non-local, as it is a topological integral over a three-dimensional manifold \mathcal{B} such that its boundary $\partial\mathcal{B}$ is the string worldsheet. Locally its integrand is a total derivative, so that the equations of motion can be consistently written in terms of worldsheet fields only. The choice $q = 0$ gives back the pure RR case, while $q = 1$ corresponds to the pure NSNS one. Interestingly, the string-theoretical S duality exchanges the RR and NSNS fluxes, which corresponds to sending $q \rightarrow \sqrt{1 - q^2}$ in this description. However, S duality is not realized perturbatively, so that the relative invariance will not necessarily be manifest in the string spectrum. Using an appropriate extension of (8.21) to include fermions, it was shown in ref. [52] that the corresponding classical theory is integrable for a suitable choice of the couplings. Such a choice is precisely the same that guarantees κ symmetry as well as conformal invariance. In fact, the same picture holds at the classical level also when including the massless modes and considering the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background.

Towards quantum integrability

Following this realisation, Hoare and Tseytlin started investigating the integrability of the quantum theory in the S-matrix approach that we described in chapter 3. In ref. [102], they computed the tree-level S matrix for $q \neq 0$, focusing on the simplest case of massive excitations on $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$. In that case it turned out that the q -dependence was quite mild, and amounted to a relatively simple modification of some of the S-matrix elements and in particular of the tree-level dispersion relation

$$\sqrt{p^2 + 1} \quad \longrightarrow \quad \sqrt{(p \pm q)^2 + 1 - q^2}, \quad (8.22)$$

where the sign \pm should be chosen suitably for each the S-matrix element. Building on that, in ref. [66] they proposed an all-loop, mixed-flux S matrix in terms of a modification of the one proposed in ref. [S3]. The “matrix part” of S should then be given by replacing the usual⁷ definition of the Zukhovski variables

$$e^{ip} = \frac{x^+(p)}{x^-(p)}, \quad E(p) + 1 = i h (x^-(p) + x^+(p)), \quad (8.23)$$

with the q -dependent ones

$$e^{ip} = \frac{x_\pm^+(p)}{x_\pm^-(p)}, \quad E_\pm(p) + 1 \pm 2hq \sin \frac{p}{2} = i h \sqrt{1 - q^2} (x_\mp^-(p) + x_\pm^+(p)), \quad (8.24)$$

⁷The form in which we write the definition of the Zhukovski parameters here is equivalent to (3.47), as it can be seen by using the definition of $E(p)$ in terms of x^\pm .

where we in fact introduced two pairs of Zhukovski parameters x_{\pm}^{\pm} , each pair depending on one modified dispersion relation E_{\pm} , with⁸

$$E_{\pm}(p)^2 = (1 \pm 2hq \sin \frac{p}{2})^2 + 4h^2(1 - q^2) \sin^2 \left(\frac{p}{2}\right). \quad (8.25)$$

This conjecture for the S matrix is compatible with the tree-level one, and with the off-shell symmetry algebra of the mixed flux theory, which however does not fix the dispersion relation entirely. The dispersion relation can essentially be narrowed down to (8.25) by requiring periodicity in p , which is what happens in the pure-RR case. As we mentioned in the introduction, one can think of this periodicity as coming from a discretisation of the worldsheet whence the spin-chain picture emerges—which is well established in several instances of integrable (pure-RR) string backgrounds.

To further validate this proposal, Hoare, Stepanchuk and Tseytlin [67] have investigated solitonic solutions for the mixed flux theory. Studying the dyonic giant magnon solution, they were able to determine that the dispersion relation in that case takes the form

$$E_{\pm}(p)^2 = (1 \pm hqp)^2 + 4h^2(1 - q^2) \sin^2 \left(\frac{p}{2}\right), \quad (8.26)$$

that in fact is incompatible with the earlier conjecture (8.25), despite going to the same tree-level expression (8.22). Now the q -linear term in brackets spoils periodicity in the momentum p —a completely new and somewhat unexpected feature. Note that this also contradicts the naïve expectation of periodicity from a discretisation of the worldsheet that we discussed in the introduction and is at odds with a spin-chain description. This last picture is also validated from the study of the bound-state dispersion relation [67], but in light of its unexpected features some further investigation may be desirable, which may come through a two-loop calculation⁹ for the dispersion relation, in the spirit of [59, 179], that for these backgrounds has not been performed yet. In any case, the implications of such dispersion relation may be radically new and are yet to be understood.

We must conclude that so far integrability for mixed-flux backgrounds remains a bit of a puzzle even in the simplest case of $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ restricted to massive modes, but we have high hopes of witnessing significant progress in the subject in the coming years. We find this particularly exciting because it may offer a way to understand the relation between the notion of solvability in terms of the worldsheet S matrix (the pure-RR case) and by representation theory of chiral algebras (pure-NSNS). Such a structure has been partially uncovered in the case of relativistic massless integrable theories that we mentioned in section 8.1. In the case of the massless sine-Gordon model, it was possible to see how the higher charges from integrability fit into the Virasoro algebra,

⁸Special care is needed to consider the limit $q \rightarrow 1$.

⁹In the present case, a simpler near-flat-space calculation may not be sufficient to unveil the form of the dispersion relation, and a subtler near-BMN calculation would be necessary, perhaps taking advantage of the unitarity techniques that we mentioned in chapter 7.

thus relating the two descriptions, see ref. [171]. Whether something similar will happen in the $\text{AdS}_3/\text{CFT}_2$ setup and what role is played by the dual CFT_2 is a very interesting question.

8.4 Conclusions and outlook

In this chapter we have reviewed the most promising and relevant lines of research in $\text{AdS}_3/\text{CFT}_2$ integrability. As we have seen, rapid progress is being made in including massless modes into the integrability machinery, see refs. [110,111] and [S1]. More general backgrounds such as $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3$ [S4,S5] or the ones supported by mixed fluxes [66,67, 102] are also being actively investigated, and show indications of being integrable. All this offers a unique playground to gain a new understanding of integrability for massless (non-relativistic) S matrices, of the relation between the charges of the Zamolodchikov-Fadeev algebra with the Virasoro ones appearing in the CFT_2 as well as in the worldsheet CFT , and of string dualities.

Once the massless modes issue has been finally cleared, it would also be very interesting to further investigate the relations between the integrability construction and the symmetric-product orbifold CFT which is the dual to $\text{AdS}^3 \times \text{S}^3 \times \text{T}^4$ strings [10,41–46]. Quite likely, such a study would be most fruitful in a spin-chain framework such as the one of ref. [180]. Repeating that study for the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background could then shed some new light on the quite obscure CFT_2 dual arising in that case [48,49].¹⁰

We should also mention a related but somewhat more distant field of investigation, that is the study of *higher-spin theories* in AdS_3 backgrounds. Non-trivial theories involving fields of spin higher than two can be formulated on three-dimensional Minkowski space and on AdS_d spaces. Such fields may be thought as arising from the tensionless limit of string theory [182–184] whereby the tower of string states becomes massless. A special feature of AdS_3 is that there (like it happens for gravity [185, 186]) higher-spin theories can be described as Chern-Simons theories [187]. In this way, it was possible to show that asymptotically they enjoy a $\mathcal{W}_\infty(\lambda)$ infinite-dimensional symmetry [188,189], which also characterises minimal models in CFT_2 [190], see ref. [191] for a review. This provided a new approach to holographic dualities, and lead to rapid developments [192–195]—see also ref. [196] for a review—including a recent study of higher spins in the $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ background [197], whereby the same $\mathfrak{d}(2,1;\alpha)$ representations constructed in [112] and used in [S4,S5] emerge [198]. While a comparison of higher-spin theory results with the S-matrix integrability ones has not been performed yet, we expect these two approaches to make contact in the future. This may lead to a more quantitative understanding of the tensionless limit—at least at the level of the free string spectrum—adding to the recent progresses in flat space [199] and in the context of $\text{AdS}_4/\text{CFT}_3$ [200].

¹⁰A recent investigation on this topic by different techniques appeared in ref. [181].

Another interesting direction which should be explored in due course is to consider *deformations* of the AdS_3 backgrounds. This road has been taken in the case of $\text{AdS}_5/\text{CFT}_4$, where a plethora of deformations has been shown to preserve integrability, such as orbifolds [201, 202] and T-duality-shift-T-duality (TsT) deformations [203, 204] of the background, and quantum deformations (in the sense of quantum groups) of the underlying symmetry algebra [205–208], which also result in a string NLSM [209, 210], see also ref. [211] for a review of the spectral problem in these contexts. The rationale in these approaches is to get rid of as much manifest symmetry as possible in an effort to get closer to more realistic theories. This amounts to breaking R-symmetry and supersymmetry in the geometric deformations of the background, and even conformal (and Lorentz) symmetry in the case of quantum deformations. What could be new and very interesting in the case of AdS_3 is that integrable deformations of that sort may include BTZ-like black hole geometries, which as we remarked in the introduction are in fact locally isometric to AdS_3 . Putting an integrability handle on such geometries would be extremely interesting.

All of these avenues appear very exciting, and unique to three-dimensional gravity theories and its dual two-dimensional CFTs. We are confident that in the near future we will witness substantial progress along many of them.

In summary

In this thesis we have reviewed the S-matrix integrability approach to the $\text{AdS}_3/\text{CFT}_2$ holographic duality. Our detailed analysis and thorough comparisons with independent results leave no room for doubt on the possibility to understand the quantum spectral problem of these theories. Not only this is possible, but we believe that the research contained here takes us extremely close to this objective. From there, we envision the possibility of using integrability to explore several unique features of $\text{AdS}_3/\text{CFT}_2$, such as integrable deformations that may include black hole solutions. This could very well open a new chapter in our understanding of holography, of black hole physics, and of gravity.

A Appendices

A.1 Generalities of the $\mathfrak{psu}(1, 1|2)$ superalgebra

We have given the (anti)commutation relations of the superalgebra $\mathfrak{psu}(1, 1|2)$ in section 2.2. In this appendix we will collect some useful additional notions about it, see also ref. [175].

Serre-Chevalley bases

Superalgebras have in general several inequivalent Dynkin diagrams, corresponding to different choices of simple roots. Each such choice gives a set of Cartan generators \mathfrak{h}_i , and corresponding raising and lowering operators \mathfrak{e}_i and \mathfrak{f}_i , where the index i takes values from 1 to the rank of the algebra, which is 3 for $\mathfrak{psu}(1, 1|2)$. These generators satisfy an algebra of the form

$$[\mathfrak{h}_i, \mathfrak{h}_j] = 0, \quad [\mathfrak{e}_i, \mathfrak{f}_j] = \delta_{ij} \mathfrak{h}_j, \quad [\mathfrak{h}_i, \mathfrak{e}_j] = +A_{ij} \mathfrak{e}_j, \quad [\mathfrak{h}_i, \mathfrak{f}_j] = -A_{ij} \mathfrak{f}_j, \quad (\text{A.1})$$

where A_{ij} is the Cartan matrix.

In this thesis we will mainly consider two gradings of $\mathfrak{psu}(1, 1|2)$. In the $\mathfrak{su}(2)$ grading the simple roots are given by

$$\begin{aligned} \mathfrak{h}_1 &= +\mathbb{L}_3 - \mathbb{J}_3, & \mathfrak{e}_1 &= +\mathbb{Q}_{+--}, & \mathfrak{f}_1 &= +\mathbb{Q}_{-++}, \\ \mathfrak{h}_2 &= +2\mathbb{J}_3, & \mathfrak{e}_2 &= +\mathbb{J}_+, & \mathfrak{f}_2 &= +\mathbb{J}_-, \\ \mathfrak{h}_3 &= +\mathbb{L}_3 - \mathbb{J}_3, & \mathfrak{e}_3 &= +\mathbb{Q}_{+-+}, & \mathfrak{f}_3 &= -\mathbb{Q}_{-+-}. \end{aligned} \quad (\text{A.2})$$

This leads to the Cartan matrix

$$\begin{pmatrix} 0 & -1 & 0 \\ -1 & +2 & -1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (\text{A.3})$$

The corresponding Dynkin diagram is shown in figure A.1 (a). In the construction of the coset of chapter 2 and in the spin-chain one of chapter 5, we generally pick $\mathfrak{psu}(1, 1|2)_L$ to be in this grading.




Figure A.1: Three Dynkin diagrams for $\mathfrak{psu}(1, 1|2)$.

In the $\mathfrak{sl}(2)$ grading we have

$$\begin{aligned}
 \tilde{\mathfrak{h}}_1 &= -\mathbb{L}_3 + \mathbb{J}_3, & \tilde{\mathfrak{e}}_1 &= -\mathbb{Q}_{-++}, & \tilde{\mathfrak{f}}_1 &= +\mathbb{Q}_{+--}, \\
 \tilde{\mathfrak{h}}_2 &= +2\mathbb{L}_3, & \tilde{\mathfrak{e}}_2 &= +\mathbb{L}_-, & \tilde{\mathfrak{f}}_2 &= -\mathbb{L}_+, \\
 \tilde{\mathfrak{h}}_3 &= -\mathbb{L}_3 + \mathbb{J}_3, & \tilde{\mathfrak{e}}_3 &= -\mathbb{Q}_{-+-}, & \tilde{\mathfrak{f}}_3 &= -\mathbb{Q}_{+++},
 \end{aligned}
 \tag{A.4}$$

with the Cartan matrix

$$\begin{pmatrix} 0 & +1 & 0 \\ +1 & -2 & +1 \\ 0 & +1 & 0 \end{pmatrix}.
 \tag{A.5}$$

The Dynkin diagram for the $\mathfrak{sl}(2)$ grading is shown in figure A.1 (c). This is the grading that we generally use for $\mathfrak{psu}(1, 1|2)_R$.

There are also fermionic gradings of $\mathfrak{psu}(1, 1|2)$, in which all three raising operators \mathfrak{e}_i are odd. In particular we can choose them to be either

$$\mathbb{Q}_{+-+}, \quad \mathbb{Q}_{+--}, \quad \mathbb{Q}_{-++}, \quad \text{or} \quad \mathbb{Q}_{-+-}, \quad \mathbb{Q}_{--+}, \quad \mathbb{Q}_{+--}.$$

This leads to the Cartan matrices

$$\begin{pmatrix} 0 & +1 & 0 \\ +1 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \quad \text{and} \quad \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & +1 \\ 0 & +1 & 0 \end{pmatrix},
 \tag{A.6}$$

respectively, corresponding to the Dynkin diagram in figure A.1 (b).

A continuous automorphism

It is useful to note that $\mathfrak{psu}(1, 1|2)$ admits an $\mathfrak{u}(1)$ -automorphism \mathbb{U} , acting on the supercharges as

$$[\mathbb{U}, \mathbb{Q}_{a\kappa\pm}] = \pm \frac{1}{2} \mathbb{Q}_{a\kappa\pm},
 \tag{A.7}$$

and commuting with the bosonic charges. If one thinks of $\mathfrak{psu}(1, 1|2)$ as a contraction of the exceptional superalgebra $\mathfrak{d}(2, 1; \alpha)$ with $\alpha \rightarrow 0$, the generator \mathbb{U} can be identified with with one of the generators of $\mathfrak{su}(2) \subset \mathfrak{d}(2, 1; \alpha)$ that is contracted. This is described in more detail in ref. [112].

Supermatrix realisation

Once the reality condition (2.36) is imposed, we are left with eight even independent supermatrices. Eliminating trace and supertrace, we can identify the six even generators as

$$\begin{aligned}
 \mathbb{L}_1 &= \frac{1}{2} \begin{pmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \mathbb{J}_1 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{pmatrix}, \\
 \mathbb{L}_2 &= \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \mathbb{J}_2 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \\
 \mathbb{L}_3 &= \frac{1}{2} \begin{pmatrix} i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \mathbb{J}_3 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix},
 \end{aligned} \tag{A.8}$$

while the eight supercharges are

$$\begin{aligned}
 \mathbb{Q}_{11}^1 &= \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \mathbb{Q}_{11}^2 &= \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 \mathbb{Q}_{21}^1 &= \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}, & \mathbb{Q}_{21}^2 &= \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\
 \mathbb{Q}_{12}^1 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \mathbb{Q}_{12}^2 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 \mathbb{Q}_{22}^1 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, & \mathbb{Q}_{22}^2 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.
 \end{aligned} \tag{A.9}$$

While this choice of the generators satisfies (2.36), it can be useful to consider complex combinations such as $\mathbb{L}_\pm, \mathbb{J}_\pm$ and $\mathbb{Q}_{\pm\pm\pm}$. In particular we have

$$\mathbb{L}_\pm = \mathbb{L}_1 \pm i \mathbb{L}_2, \quad \mathbb{J}_\pm = \mathbb{J}_1 \pm i \mathbb{J}_2. \tag{A.10}$$

Two copies of $\mathfrak{psu}(1, 1|2)$

We can take the representation for \mathcal{M} , *i.e.* the one defining the left copy of $\mathfrak{psu}(1, 1|2)$, to be given by the previous expressions. The matrix in the representation $\widetilde{\mathcal{M}}$, instead, is in a representation that has opposite weights for the bosonic subalgebra. This can be realised by sending

$$\mathbb{L}_2 \rightarrow -\mathbb{L}_2, \quad \mathbb{L}_3 \rightarrow -\mathbb{L}_3, \quad \mathbb{J}_2 \rightarrow -\mathbb{J}_2, \quad \mathbb{J}_3 \rightarrow -\mathbb{J}_3. \tag{A.11}$$

In fact, this corresponds to exchanging raising with lowering operators \mathbb{L}_\pm and \mathbb{J}_\pm without affecting the commutation relations. The transformation straightforwardly extends to the whole superalgebra.

The transformation (A.11) implies that the bosonic subalgebras in each (left and right) copy of $\mathfrak{psu}(1, 1|2)$ have opposite notions of what are the positive bosonic roots. If we take this notion to be induced from the choice of simple roots, it follows that the two copies cannot be in the same grading. More specifically, if one copy is in the grading of fig. A.1a with Cartan matrix (A.3), then the other is in the grading of fig. A.1c with Cartan matrix (A.5). If one of them has the fully-fermion grading of fig. A.1b, then the other does too, and they have Cartan matrices of opposite signs, see (A.6).

It is also interesting to notice that then the \mathbb{Z}_4 -automorphism Ω of eq. (2.41) acts on a bosonic matrix as

$$\Omega(\mathbb{L}_j \oplus \mathbb{O}) = \mathbb{O} \oplus \tilde{\mathbb{L}}_j, \quad \Omega(\mathbb{J}_j \oplus \mathbb{O}) = \mathbb{O} \oplus \tilde{\mathbb{J}}_j, \quad (\text{A.12})$$

i.e. exchanges the left and right bosonic subalgebras.

A.2 Explicit expressions for the coset action

Here we will collect some explicit expressions that appear when evaluating the coset action of chapter 2 and expanding it.

Parametrisation of group elements

We parametrise a group element by (2.47). The light-cone coordinate $\Lambda(t, \phi)$ has been defined in eq. (2.48). The transverse coordinates $\{x_k\} = (z_1, z_2, y_1, y_2)$ are given by

$$g(x) = i \frac{\mathbb{I}_z + z_1 \Sigma_1 + z_2 \Sigma_2}{\sqrt{1 - \frac{1}{4}|z|^2}} + i \frac{\mathbb{I}_y + y_1 \Sigma_3 + y_2 \Sigma_4}{\sqrt{1 + \frac{1}{4}|y|^2}}, \quad (\text{A.13})$$

with

$$\begin{aligned} \mathbb{I}_z &= \frac{1}{2}(\mathbb{I} + \Sigma_+ \Sigma_-), & \Sigma_1 &= \mathbb{L}_1^L - \mathbb{L}_1^R, & \Sigma_3 &= \mathbb{J}_1^L - \mathbb{J}_1^R, \\ \mathbb{I}_y &= \frac{1}{2}(\mathbb{I} - \Sigma_+ \Sigma_-), & \Sigma_2 &= \mathbb{L}_2^L - \mathbb{L}_2^R, & \Sigma_4 &= \mathbb{J}_2^L - \mathbb{J}_2^R. \end{aligned} \quad (\text{A.14})$$

We have split the fermions in massive and massless ones. The former are θ_L^1, θ_L^2 and θ_R^1, θ_R^2 that in terms of a $\mathfrak{psu}(1, 1|2)^2$ matrix we write as

$$\Psi_m = \left(\begin{array}{cccc} 0 & 0 & 0 & -\theta_L^1 \\ 0 & 0 & \bar{\theta}_{2L} & 0 \\ 0 & \theta_L^2 & 0 & 0 \\ \bar{\theta}_{1L} & 0 & 0 & 0 \end{array} \right) \oplus \left(\begin{array}{cccc} 0 & 0 & 0 & -\theta_R^2 \\ 0 & 0 & -\bar{\theta}_{1R} & 0 \\ 0 & -\theta_R^1 & 0 & 0 \\ \bar{\theta}_{2R} & 0 & 0 & 0 \end{array} \right). \quad (\text{A.15})$$

The four massless fermions are instead

$$\Psi_l = \begin{pmatrix} 0 & 0 & \eta_L^1 & 0 \\ 0 & 0 & 0 & \eta_L^2 \\ -\bar{\eta}_{1L} & 0 & 0 & 0 \\ 0 & \bar{\eta}_{2L} & 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 0 & \bar{\eta}_R^2 & 0 \\ 0 & 0 & 0 & -\eta_R^1 \\ -\bar{\eta}_{2R} & 0 & 0 & 0 \\ 0 & -\bar{\eta}_{1R} & 0 & 0 \end{pmatrix}. \quad (\text{A.16})$$

In both cases, the bar denotes complex conjugation, *e.g.*

$$(\theta_L^1)^\dagger = \bar{\theta}_{1L}, \quad (\text{A.17})$$

and so on. We can now parametrise the corresponding group element as

$$g(\Psi_{m,l}) = 1 + \Psi_{m,l} + \frac{1}{2} \Psi_{m,l}^2. \quad (\text{A.18})$$

Note that one can think that such a parametrisation emerges equivalently from $g(\Psi) = \exp(\Psi)$ or $g(\Psi) = \Psi + \sqrt{1 + \Psi^2}$, since the expansions terminate at $O(\Psi^2)$.

Finally, the T^4 elements can be found from exponentiating the corresponding generators,

$$g(X) = e^{i \sum_{k=1}^4 X_k T^k} = \prod_{k=1}^4 e^{i X_k T^k}, \quad (\text{A.19})$$

where we made it explicit that they all commute.

Explicit expressions for perturbative evaluation

To define quantities related to the transverse bosonic fields x_k let us introduce indices a, s so that $x_a = (z_1, z_2)$ and $x_s = (y_1, y_2)$. Then the metric elements are

$$g_\pm = \left(\frac{1}{\sqrt{4 + |y|^2}} \pm \frac{1}{\sqrt{4 - |z|^2}} \right), \quad g_a = \frac{2i z_a}{\sqrt{4 - |z|^2}}, \quad g_s = \frac{2i y_s}{\sqrt{4 + |y|^2}}. \quad (\text{A.20})$$

It is useful to rewrite the parametrisation (A.13) of $g(x)$ in light-cone coordinates

$$g(x) = i g_+ \mathbb{I} + i g_- \Sigma_8 + g_k \Sigma_k, \quad (\text{A.21})$$

where \mathbb{I} is the identity, $\Sigma_8 = -\Sigma_+ \Sigma_-$ and Σ_k are given by (A.14). It is then easy to compute

$$g(x)^2 = G_+ \mathcal{I} + G_- \Sigma_8 + G_M \Sigma_M, \quad (\text{A.22})$$

where now

$$G_\pm = - \left(\frac{4 - |y|^2}{4 + |y|^2} \pm \frac{4 + |z|^2}{4 - |z|^2} \right), \quad G_a = - \frac{8 z_a}{4 - |z|^2}, \quad G_s = - \frac{8 y_s}{4 + |y|^2}. \quad (\text{A.23})$$

We can work out the expression of the momenta $\{p_k\} = (p_1^z, p_2^z, p_1^y, p_2^y)$ canonically conjugated to $\{x_k\} = (z_1, z_2, y_1, y_2)$, in terms of the auxiliary fields ϖ_k . They are

$$p_a = \frac{4\varpi_a}{4 - |z|^2}, \quad p_s = \frac{4\varpi_s}{4 + |y|^2}. \quad (\text{A.24})$$

Let us now look at the Lagrangian more in detail. The even and odd currents are given by

$$\begin{aligned} A_{\text{even}} &= -\frac{i}{2}g_x^{-1} \left(dx_+ \Sigma_+ (1 + 2\Psi_m^2) + \frac{1}{2}dx_- \Sigma_- - i[d\Psi_m, \Psi_m] \right) g_x - g_x^{-1} dg_x, \\ A_{\text{odd}} &= -i g_x^{-1} dx_+ \Sigma_+ \Psi_m g_x - g_x^{-1} d\Psi_m g_x, \end{aligned} \quad (\text{A.25})$$

where we used that the fermion parametrisation (A.15) is quadratic at most. The part of the even current that depends solely on the transverse bosonic coordinates is

$$A_{\text{even}}^\perp = -\frac{1}{2}g_x^{-1} [\dot{\Psi}_m, \Psi_m] g_x - g_x^{-1} \dot{g}_x. \quad (\text{A.26})$$

Using these expressions it is easy to get to eq. (2.57). Moreover we have

$$\text{str}(\varpi g_x^{-1} \dot{g}_x) = p_k \dot{x}_k. \quad (\text{A.27})$$

We can also further simplify

$$\begin{aligned} \text{str}(\varpi g_x^{-1} [\dot{\Psi}_m, \Psi_m] g_x) &= \frac{i}{4} (\varpi_+ G_+ - \varpi_- G_-) \text{str}(\Sigma_+ [\dot{\Psi}_m, \Psi_m]) \\ &\quad + \frac{1}{4} \varpi_k \text{str}(\Sigma_k g_x^{-1} [\dot{\Psi}_m, \Psi_m] g_x). \end{aligned} \quad (\text{A.28})$$

Solving the constraints

We have from $\mathcal{C}_1 = 0$

$$-x'_- = p_k x'_k + \frac{1}{4} \varpi_k \text{str}(\Sigma_k g_x^{-1} [\Psi'_m, \Psi_m] g_x) + \frac{i}{8} \text{str}(\Sigma_+ [\Psi'_m, \Psi_m]), \quad (\text{A.29})$$

which we can expand as

$$-x'_- = p_k x'_k - \frac{i}{4} \text{str}(\Sigma_+ \Psi_m \Psi'_m) + \dots, \quad (\text{A.30})$$

where the ellipsis indicate higher order terms. The quadratic constraint is

$$\mathcal{C}_2 = \frac{1}{G_+} (\varpi_- G_- + 1) \varpi_- + |\varpi_k|^2 + h \text{str}((A_1^{(2)})^2), \quad (\text{A.31})$$

where

$$\begin{aligned} A_1^{(2)} &= -\frac{i}{4} x'_- (g_x^{-1} \Sigma_- g_x - \Omega(g_x^{-1} \Sigma_- g_x)) - \frac{1}{2} (g_x^{-1} g'_x + g'_x g_x^{-1}) \\ &\quad - \frac{1}{4} (g_x^{-1} [\Psi'_m, \Psi_m] g_x - \Omega(g_x^{-1} [\Psi'_m, \Psi_m] g_x)). \end{aligned} \quad (\text{A.32})$$

By dropping higher order terms, from $\mathcal{C}_2 = 0$ we have

$$\varpi_- = -\frac{1}{2}|\varpi_k|^2 - \frac{1}{2}|x'_k|^2 + \dots \quad (\text{A.33})$$

Dropping the total τ -derivative \dot{x}_- , the Lagrangian reads

$$\mathcal{L}_2 = \mathbf{p}_- + p_k \dot{x}_k - \frac{i}{4} \text{str}(\Sigma_+ \Psi_m \dot{\Psi}_m) - \frac{i}{4} \varepsilon^{\alpha\beta} \text{str}(A_\alpha^{\text{odd}} \Omega(A_\beta^{\text{odd}})) + \dots, \quad (\text{A.34})$$

with

$$\mathbf{p}_- = \varpi_- - \frac{1}{2}(|x_k|^2 + \text{str}\Psi_m^2) + \dots \quad (\text{A.35})$$

For the purpose of expanding the Lagrangian and the Noether charge (2.67) in powers of the fields it is useful to note that at leading order it is

$$A_0^{\text{odd}} = -i\Psi_m - \dot{\Psi}_m + \dots, \quad A_1^{\text{odd}} = -\Psi'_m + \dots \quad (\text{A.36})$$

When plugged into the Lagrangian (A.34), and of using the explicit form of Ψ_m , this gives contributions of the form

$$-\frac{i}{4} \varepsilon^{\alpha\beta} \text{str}(A_\alpha^{\text{odd}} \Omega(A_\beta^{\text{odd}})) = \frac{1}{2} \text{str}(\Sigma_+ \Psi_m \Omega(\Psi'_m)) - \frac{i}{2} (\dot{\Psi}_m \Omega(\Psi'_m)) + \dots, \quad (\text{A.37})$$

where the double-derivative terms all vanish after two integrations by parts. By these expressions it is easy to derive the Lagrangian and Hamiltonian of the main text (2.71).

To derive some of these results is useful to use the identities

$$\Sigma_\pm g_x^{-1} = g_x \Sigma_\pm, \quad \Sigma_\pm \Psi_m = \mp \Psi_m \Sigma_\pm. \quad (\text{A.38})$$

A.3 Quadratic charges in terms of the fields

In order to compactly write down that form of the symmetry generators, it is useful to introduce complex bosonic coordinates

$$X_L^1 = z_1 + i z_2, \quad X_L^2 = y_1 + i y_2, \quad X_R^1 = z_1 - i z_2, \quad X_R^2 = y_1 - i y_2, \quad (\text{A.39})$$

and conjugate momenta

$$P_{1L} = p_1^z - i p_2^z, \quad P_{2L} = p_1^y - i p_2^y, \quad P_{1R} = p_1^z + i p_2^z, \quad P_{2R} = p_1^y + i p_2^y. \quad (\text{A.40})$$

It is clear that $(X_L^j)^\dagger = X_R^j$ and $(P_{jL})^\dagger = P_{jR}$. We will raise and lower indices by

$$X_{jL} = \delta_{jk} X_L^k, \quad X_{jR} = \delta_{jk} X_R^k, \quad P_L^j = \delta^{jk} P_{kL}, \quad P_R^j = \delta^{jk} P_{kR}. \quad (\text{A.41})$$

Similarly, if necessary, we will raise and lower the fermion indices by

$$\theta_{jL,R} = \delta_{jk} \theta_{L,R}^j, \quad \bar{\theta}_{L,R}^j = \delta^{jk} \bar{\theta}_{jL,R}. \quad (\text{A.42})$$

The quadratic Lagrangian becomes

$$\mathcal{L}_2 = P_{j_L} \dot{X}_L^j + P_{j_R} \dot{X}_R^j + i \bar{\theta}_{j_L} \dot{\theta}^{j_L} + i \bar{\theta}_{j_R} \dot{\theta}^{j_R} - \mathcal{H}_2. \quad (\text{A.43})$$

In particular, this implies that the non-vanishing Poisson brackets for the bosons are

$$\{X_L^j(\sigma), P_{k_L}(\tilde{\sigma})\} = \delta_k^j \delta(\sigma - \tilde{\sigma}), \quad \{X_R^j(\sigma), P_{k_R}(\tilde{\sigma})\} = \delta_k^j \delta(\sigma - \tilde{\sigma}), \quad (\text{A.44})$$

compatibly with the ones we originally found (2.74–2.75).

The form of the bosonic charges is then

$$\begin{aligned} \mathbb{H} &= \int d\sigma \left(\frac{1}{2} P_L^j P_{j_R} + 2X_L^{j'} X'_{j_R} + 2X_L^j X_{j_R} + \theta_L^j \theta'_{j_R} - \bar{\theta}_L^j \bar{\theta}'_{j_R} + \bar{\theta}_{j_L} \theta_L^j + \bar{\theta}_{j_R} \theta_R^j \right), \\ \mathbb{M} &= \int d\sigma \left(\bar{\theta}_{j_L} \theta_L^j - \bar{\theta}_{j_R} \theta_R^j + iP_{j_L} X_L^j - iP_{j_R} X_R^j \right), \\ \mathbb{N} &= \int d\sigma i \left(P_{1_L} X_L^1 - P_{2_L} X_L^2 - P_{1_R} X_R^1 + P_{2_R} X_R^2 \right). \end{aligned} \quad (\text{A.45})$$

The four supercharges $\mathbb{Q}^{j_L, R}$ read

$$\begin{aligned} \mathbb{Q}^{1L} &= e^{-\frac{i}{4}\pi} \int d\sigma e^{\frac{i}{2}x} \left(\frac{i}{2} P_{2R} \bar{\theta}_{1L} - X_L^2 \theta_R^{1'} + X_L^2 \bar{\theta}_{1L} \right. \\ &\quad \left. - \frac{i}{2} P_{1L} \theta_{2L} + X_R^1 \bar{\theta}'_{2R} + X_R^1 \theta_L^2 \right), \\ \mathbb{Q}^{1R} &= e^{-\frac{i}{4}\pi} \int d\sigma e^{\frac{i}{2}x} \left(\frac{i}{2} P_{2L} \bar{\theta}_{1R} - X_R^2 \theta_L^{1'} + X_R^2 \bar{\theta}_{1R} \right. \\ &\quad \left. - \frac{i}{2} P_{1R} \theta_R^2 + X_L^1 \bar{\theta}'_{2L} + X_L^1 \theta_R^2 \right), \\ \mathbb{Q}^{2L} &= e^{-\frac{i}{4}\pi} \int d\sigma e^{\frac{i}{2}x} \left(\frac{i}{2} P_{1L} \theta_L^1 + X_R^1 \bar{\theta}'_{1R} - X_R^1 \theta_L^1 \right. \\ &\quad \left. + \frac{i}{2} P_{2R} \bar{\theta}_{2L} + X_L^2 \theta_R^{2'} + X_L^2 \bar{\theta}_{2L} \right), \\ \mathbb{Q}^{2R} &= e^{-\frac{i}{4}\pi} \int d\sigma e^{\frac{i}{2}x} \left(\frac{i}{2} P_{1R} \theta_R^1 + X_L^1 \bar{\theta}'_{1L} - X_L^1 \theta_R^1 \right. \\ &\quad \left. + \frac{i}{2} P_{2L} \bar{\theta}_{2R} + X_R^2 \theta_L^{2'} + X_R^2 \bar{\theta}_{2R} \right), \end{aligned} \quad (\text{A.46})$$

and the remaining four can be found by taking the complex conjugate of these expressions. Notice how left and right supercharges differ only by exchanging the labels $L \leftrightarrow R$. The factor $e^{-\frac{i}{4}\pi}$ is inserted so that the charges take a simpler expression in terms of oscillators.

From field to oscillators

We can define bosonic lowering operators

$$\begin{aligned}
 a_{\text{L}}^{-}(p) &= \frac{1}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} \left(\omega_p X_{\text{L}}^1(\sigma) + \frac{i}{2} P_{1\text{R}}(\sigma) \right) e^{-ip\sigma}, \\
 a_{\text{R}}^{-}(p) &= \frac{1}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} \left(\omega_p X_{\text{R}}^1(\sigma) + \frac{i}{2} P_{1\text{L}}(\sigma) \right) e^{-ip\sigma}, \\
 a_{\text{L}}^{++}(p) &= \frac{1}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} \left(\omega_p X_{\text{L}}^2(\sigma) + \frac{i}{2} P_{2\text{R}}(\sigma) \right) e^{-ip\sigma}, \\
 a_{\text{R}}^{++}(p) &= \frac{1}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} \left(\omega_p X_{\text{R}}^2(\sigma) + \frac{i}{2} P_{2\text{L}}(\sigma) \right) e^{-ip\sigma},
 \end{aligned} \tag{A.47}$$

with the raising operator being the conjugate of these. Similarly, for fermions we have

$$\begin{aligned}
 a_{\text{L}}^{-+}(p) &= \frac{e^{+\frac{i}{4}\pi}}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} (f_p \theta_{\text{L}}^1(\sigma) - i g_p \bar{\theta}_{1\text{R}}(\sigma)) e^{-ip\sigma}, \\
 a_{\text{R}}^{-+}(p) &= \frac{e^{+\frac{i}{4}\pi}}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} (f_p \theta_{\text{R}}^1(\sigma) - i g_p \bar{\theta}_{1\text{L}}(\sigma)) e^{-ip\sigma}, \\
 a_{\text{L}}^{+-}(p) &= \frac{e^{-\frac{i}{4}\pi}}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} (f_p \bar{\theta}_{2\text{L}}(\sigma) + i g_p \theta_{\text{R}}^2(\sigma)) e^{-ip\sigma}, \\
 a_{\text{R}}^{+-}(p) &= \frac{e^{-\frac{i}{4}\pi}}{\sqrt{2\pi}} \int \frac{d\sigma}{\sqrt{\omega_p}} (f_p \bar{\theta}_{2\text{R}}(\sigma) + i g_p \theta_{\text{L}}^2(\sigma)) e^{-ip\sigma}.
 \end{aligned} \tag{A.48}$$

Once we drop the x_- -dependence from (A.46), and by making use of the explicit form of the wave-function parameters (2.81–2.84) with mass $m = 1$, the charges can be recast in the form (2.86–2.87).

A.4 Analytic properties of the dressing factors

In this appendix we present the proofs of some identities we used in chapter 4. In particular, we provide the proof that the phase $\chi^-(x, y)$ solves the crossing equation (4.44). This can be easily adapted to check that the Hernández-López phase as defined in (4.37) solves the “odd” crossing equation (4.38) with the choice of path depicted in figure 4.3.

Solving the “difference” crossing equation (4.44)

Let us define the following integral

$$\begin{aligned}
 \Phi^-(x, y) &= \oint \frac{dw}{8\pi} \frac{1}{x-w} \log \left[(y-w) \left(1 - \frac{1}{yw} \right) \right] \text{sign}((w-1/w)/i) \quad -x \leftrightarrow y \\
 &= \left(\oint - \oint \right) \frac{dw}{8\pi} \frac{1}{x-w} \log \left[(y-w) \left(1 - \frac{1}{yw} \right) \right] \quad -x \leftrightarrow y,
 \end{aligned} \tag{A.49}$$

which is reminiscent of (4.37). This function satisfies a property which will be crucial in what follows, that is

$$\Phi^-(x, y) - \Phi^-(1/x, y) = 0, \quad (\text{A.50})$$

which we will prove in the next subsection. Furthermore, when $|y| > 1$ is fixed and $|x| \geq 1$ approaches the unit circle, $\Phi^-(x, y)$ has a jump discontinuity. As we will discuss in a later subsection, the value of the discontinuity depends on whether x approaches the unit circle from below the real line, in which case

$$\Phi^-(e^{i\varphi+\epsilon}, y) = \Phi^-(e^{i\varphi-\epsilon}, y) + \delta_\uparrow(e^{i\varphi}, y) + O(\epsilon), \quad \epsilon > 0, \quad -\pi < \varphi < 0, \quad (\text{A.51})$$

with¹

$$\delta_\uparrow(x, y) = -\frac{i}{2} \log \left[(y-x) \left(1 - \frac{1}{xy} \right) \right], \quad (\text{A.52})$$

or from above, where

$$\Phi^-(e^{i\varphi+\epsilon}, y) = \Phi^-(e^{i\varphi-\epsilon}, y) + \delta_\downarrow(e^{i\varphi}, y) + O(\epsilon), \quad \epsilon > 0, \quad 0 < \varphi < \pi, \quad (\text{A.53})$$

with $\delta_\downarrow(x, y) = -\delta_\uparrow(x, y)$.

These ingredients are all we need to construct a solution of (4.44). In the physical region we define

$$\chi^-(x, y) \equiv \Phi^-(x, y) \quad |x|, |y| > 1. \quad (\text{A.54})$$

In order to continue this function to the crossed region, it is important to recall our choice of cuts of figure 4.3: both $x^+(z)$ and $x^-(z)$ will cross the unit circle *below the real line*. Therefore, we define

$$\chi^-(x, y) \equiv \Phi^-(x, y) + \delta_\uparrow(x, y) \quad |x| < 1, \quad |y| > 1, \quad (\text{A.55})$$

which is continuous across the lower half circle by construction. Using (A.50) we have

$$\chi^-(x, y) - \chi^-(1/x, y) = -\delta_\uparrow(1/x, y) = -\delta_\uparrow(x, y), \quad |x|, |y| > 1. \quad (\text{A.56})$$

Rewriting the left-hand-side of (4.44) in terms of $\chi(x, y)$ gives finally

$$\frac{\sigma^-(z_1, z_2)^2}{\sigma^-(z_1 + \omega_2, z_2)^2} = \frac{e^{-2i(\delta_\uparrow(x^+, y^+) + \delta_\uparrow(x^-, y^-))}}{e^{-2i(\delta_\uparrow(x^+, y^-) + \delta_\uparrow(x^-, y^+))}}, \quad (\text{A.57})$$

which coincides with (4.45).

¹More precisely, the following relation holds up to an arbitrary function of y only, and in an appropriate branch of the logarithm, see the next subsection. Such a function plays no role in the crossing equation.

Identity for $\Phi^-(x, y) - \Phi^-(1/x, y)$

In this subsection we prove equation (A.50). Define

$$F(x, y) = F^{\curvearrowright}(x, y) - F^{\curvearrowleft}(x, y) = \int\limits_{\curvearrowright} f(w, x, y)dw - \int\limits_{\curvearrowleft} f(w, x, y)dw, \quad (\text{A.58})$$

where $F^{\curvearrowright}, F^{\curvearrowleft}$ corresponds to the first and second integral, respectively, and

$$f(w, x, y) = \frac{1}{8\pi} \frac{1}{x-w} \log \left[(y-w) \left(1 - \frac{1}{yw} \right) \right], \quad (\text{A.59})$$

so that $\Phi^-(x, y) = F(x, y) - F(y, x)$. Since $f(w, x, y) - f(w, x, 1/y) = 0$, we see that

$$F^{\curvearrowright}(x, y) - F^{\curvearrowright}(x, 1/y) = 0, \quad \text{and} \quad F^{\curvearrowleft}(x, y) - F^{\curvearrowleft}(x, 1/y) = 0. \quad (\text{A.60})$$

A change of integration variable, $u = 1/w$, can be used to derive the following identity

$$\begin{aligned} F^{\curvearrowleft}(1/x, y) &= \int\limits_{\curvearrowleft} \frac{dw}{8\pi} \frac{1}{1/x-w} \log \left[(y-w) \left(1 - \frac{1}{yw} \right) \right] \\ &= \int\limits_{\curvearrowleft} \frac{du}{8\pi u^2} \frac{1}{1/x-1/u} \log \left[(y-u) \left(1 - \frac{1}{yu} \right) \right] \\ &= - \int\limits_{\curvearrowleft} \frac{du}{8\pi} \frac{1}{x-u} \log \left[(y-u) \left(1 - \frac{1}{yu} \right) \right] - \int\limits_{\curvearrowleft} \frac{du}{8\pi u} \log \left[(y-u) \left(1 - \frac{1}{yu} \right) \right] \\ &= -F^{\curvearrowright}(x, y) - \phi^-(y), \end{aligned} \quad (\text{A.61})$$

for arbitrary x with $|x| \neq 1$. Sending $x \rightarrow 1/x$ in the above equation gives

$$F^{\curvearrowright}(1/x, y) = -F^{\curvearrowleft}(x, y) - \phi^-(y). \quad (\text{A.62})$$

Combining equations (A.60), (A.61) and (A.62) and writing out Φ in terms of F^{\curvearrowright} and F^{\curvearrowleft} one may check that (A.50) holds.

Discontinuities of $\Phi^-(x, y)$ at $|x| = 1$

Let us split $\Phi^-(x, y)$ in terms of $F^{\curvearrowright}, F^{\curvearrowleft}$ as in (A.58), and focus on the discontinuities of $F^{\curvearrowleft}(x, y)$. The discontinuity in x follows immediately from Cauchy's theorem, and is given by

$$F^{\curvearrowleft}(e^{i\varphi+\epsilon}, y) = F^{\curvearrowleft}(e^{i\varphi-\epsilon}, y) + d_{\uparrow}^{(x)}(e^{i\varphi}, y) + O(\epsilon), \quad \epsilon > 0, \quad -\pi < \varphi < 0, \quad (\text{A.63})$$

crossing the unit circle from below, with

$$d_{\uparrow}^{(x)}(x, y) = \frac{i}{4} \log \left[(y-x) \left(1 - \frac{1}{xy} \right) \right]. \quad (\text{A.64})$$

Note that $F^{\curvearrowleft}(x, y)$ is continuous in x across the upper half-circle.

To find the discontinuity in y , we can consider $\partial_y F^\smile(x, y)$ for $|x|, |y| > 1$; bringing the derivative under the integral gets rid of the logarithm. The resulting function has a discontinuity on the lower half circle:

$$\partial_y F^\smile(x, e^{i\varphi+\epsilon}) = \partial_y F^\smile(x, e^{i\varphi-\epsilon}) + d_\uparrow^{(y')} (x, e^{i\varphi}) + O(\epsilon), \quad \epsilon > 0, \quad -\pi < \varphi < 0. \quad (\text{A.65})$$

The appropriate primitive of $d_\uparrow^{(y')} (x, y)$ in y gives the discontinuity of $F^\smile(x, y)$ from below. Such a primitive is

$$d_\uparrow^{(y)} (x, y) = -\frac{i}{4} \log(x - y) + \phi_\uparrow(x), \quad (\text{A.66})$$

where $\phi_\uparrow(x)$ is arbitrary function. Furthermore, there is also a discontinuity on the upper half circle:

$$\partial_y F^\smile(x, e^{i\varphi+\epsilon}) = \partial_y F^\smile(x, e^{i\varphi-\epsilon}) + d_\downarrow^{(y')} (x, e^{i\varphi}) + O(\epsilon), \quad \epsilon > 0, \quad 0 < \varphi < \pi, \quad (\text{A.67})$$

whose primitive is

$$d_\downarrow^{(y)} (x, y) = \frac{i}{4} (\log(1 - xy) - \log x - \log y) + \phi_\downarrow(x). \quad (\text{A.68})$$

It is easy to repeat this analysis for $F^\frown(x, y)$, where we find essentially the same results up to exchanging the upper and lower circles. Putting everything together proves (A.52) up to such an arbitrary function of x , which however drops out of the crossing equation (A.57), canceling among the contributions of the four χ 's. Since the crossing relations (A.57) are written in exponential form, the logarithmic branch cuts of the discontinuities play no role.

If we instead had considered the discontinuities of $\Phi^-(x, y)$ when crossing the unit circle in the *upper* half-plane we would have found an extra minus sign upon crossing.

Singularities of the dressing phases

Let us investigate the singularities of the dressing phases $\theta(x, y)$ and $\tilde{\theta}(x, y)$. They are defined in terms of $\chi(x, y)$, $\tilde{\chi}(x, y)$ by (4.47). Since the analytic properties of the BES phase are well known [75, 142], we will focus on the semisum and semidifference of the HL phase with $\chi^-(x, y)$. We are interested in logarithmic singularities as x and y take particular positions with respect to each other. To find them, let us consider the integrals

$$\Theta^\pm(x, y) = \frac{1}{2} (-\Phi^{\text{HL}}(x, y) \pm \Phi^-(x, y)), \quad (\text{A.69})$$

where $\Phi^{\text{HL}}(x, y)$ is the integral defining the HL phase in the physical region,

$$\Phi^{\text{HL}}(x, y) = \left(\oint - \oint \right) \frac{dw}{4\pi} \frac{1}{x-w} (\log(y-w) - \log(y-1/w)), \quad (\text{A.70})$$

and $\Phi^-(x, y)$ is defined in (A.49). Singularities may arise only at $y = x$ or $y = 1/x$. However, we can evaluate explicitly

$$\Theta^\pm(x, y)|_{y=x} = 0, \quad \Theta^\pm(x, y)|_{y=1/x} = \frac{1}{4\pi} (4\text{Li}_2(x) - \text{Li}_2(x^2)), \quad (\text{A.71})$$

with $|y| > 1$, so that no singularity arises from the integral representation. Since this coincides with the phase in the physical region, we can conclude that there is no discontinuity for the phases at $x = y$ when both variables are in the physical region.

When $y = 1/x$ one of the variables, *e.g.* x , must be inside the unit circle. Therefore, as explained earlier in this appendix, we must continue $\Theta^\pm(x, y)$ in x through the lower half circle. In order to do this we need to find the discontinuity of $\Theta^\pm(x, y)$ there. In appendix A.4 we worked out the discontinuity of $\Phi^-(x, y)$ to be as in (A.52), *i.e.*,

$$\delta_{\uparrow}^-(x, y) = -\frac{i}{2} \log \left[(y-x) \left(1 - \frac{1}{xy} \right) \right]. \quad (\text{A.72})$$

Using Cauchy's theorem we find that $\Phi^{\text{HL}}(x, y)$ satisfies

$$\Phi^{\text{HL}}(e^{i\varphi+\epsilon}, y) = \Phi^{\text{HL}}(e^{i\varphi-\epsilon}, y) + \delta_{\uparrow}^{\text{HL}}(e^{i\varphi}, y) + O(\epsilon), \quad \epsilon > 0, \quad -\pi < \varphi < 0, \quad (\text{A.73})$$

with

$$\delta_{\uparrow}^{\text{HL}}(x, y) = -\frac{i}{2} \log \left[\frac{y-x}{y-1/x} \right]. \quad (\text{A.74})$$

Using this to analytically continue (A.69) we have that when $|x| < 1$ and $|y| > 1$, there is no singularity in $\tilde{\chi}(x, y)$ at $y = 1/x$. However $\chi(x, y)$ has a logarithmic singularity such that

$$e^{2i\chi(x, y)} \approx \left(y - \frac{1}{x} \right), \quad \text{when } y \approx 1/x. \quad (\text{A.75})$$

A.5 Dualisation of the Bethe ansatz equations

While performing the nesting procedure in chapter 6, we had to chose a set of level-I excitations. Here we will show that the different choices we could make lead there, to equivalent equations (*i.e.* that describe the same spectrum), by performing a fermionic duality [27, 156] of the Bethe equations.

Duality for the $\mathfrak{su}(1|1)_{\text{c.e.}}^2$ Bethe equations

Let us start from the simpler case we treated in section 6.2. We will work in the spin-chain frame for definiteness.

The idea is that one can write a new set of equations equivalent to the previous one, in which all the auxiliary roots y, \tilde{y} are replaced by a dual set of auxiliary roots $\hat{y}, \hat{\tilde{y}}$. We will see that the resulting equations have the same form as the original ones, up to

exchanging the left and right sectors, which makes it clear that they correspond to the choice V_B^I in (6.17).

Let us define the polynomial $\mathcal{P}(\xi)$ as

$$\mathcal{P}(\xi) = \prod_{j=1}^{M_L^I} (\xi - x_j^+) \prod_{j=1}^{M_R^I} (\xi - \frac{1}{\tilde{x}_j}) - \prod_{j=1}^{M_L^I} (\xi - x_j^-) \prod_{j=1}^{M_R^I} (\xi - \frac{1}{\tilde{x}_j^+}). \quad (\text{A.76})$$

This is a polynomial of degree $n = M_L^I + M_R^I - 1$. Then the Bethe equations for the left and right auxiliary excitations can be rewritten respectively as

$$\mathcal{P}(y) = 0, \quad \mathcal{P}(1/\tilde{y}) = 0. \quad (\text{A.77})$$

Another zero of the polynomial is at $\xi = 0$. The equation $\mathcal{P}(0) = 0$ is equivalent to the level-matching condition. We denote the remaining zeros by \hat{y} and $\hat{\tilde{y}}$, because they correspond to the dualisation of respectively left and right excitations. We can thus rewrite $\mathcal{P}(\xi)$ as

$$\mathcal{P}(\xi) = \xi \prod_{j=1}^{M_L^{II}} (\xi - y_j) \prod_{j=1}^{\widehat{M}_L^{II}} (\xi - \hat{y}_j) \prod_{j=1}^{M_R^{II}} (\xi - \frac{1}{\tilde{y}_j}) \prod_{j=1}^{\widehat{M}_R^{II}} (\xi - \frac{1}{\tilde{\hat{y}}_j}) \quad (\text{A.78})$$

for $\widehat{M}_L^{II} = M_L^I - K_L^{II} - 1$ and $\widehat{M}_R^{II} = M_R^I - K_R^{II} - 1$ roots. Let us now dualise the Bethe equations for level-I excitations. We write the expression $\mathcal{P}(x_k^+)/\mathcal{P}(x_k^-)$ using the two possible representations for the polynomial, getting the equation

$$\begin{aligned} \frac{x_k^+}{x_k^-} \prod_{j=1}^{M_L^{II}} \frac{x_k^+ - y_j}{x_k^- - y_j} \prod_{j=1}^{\widehat{M}_L^{II}} \frac{x_k^+ - \hat{y}_j}{x_k^- - \hat{y}_j} \prod_{j=1}^{M_R^{II}} \frac{x_k^+ - 1/\tilde{y}_j}{x_k^- - 1/\tilde{y}_j} \prod_{j=1}^{\widehat{M}_R^{II}} \frac{x_k^+ - 1/\tilde{\hat{y}}_j}{x_k^- - 1/\tilde{\hat{y}}_j} = \\ - \prod_{j=1}^{M_L^I} \frac{x_k^+ - x_j^-}{x_k^- - x_j^+} \prod_{j=1}^{M_R^I} \frac{x_k^+ - 1/\tilde{x}_j^+}{x_k^- - 1/\tilde{x}_j^-}, \end{aligned} \quad (\text{A.79})$$

that becomes

$$\begin{aligned} \left(\frac{x_k^+}{x_k^-} \right)^{-1 - M_R^{II} - \widehat{M}_R^{II} + M_R^I} \prod_{\substack{j=1 \\ j \neq k}}^{M_L^I} \frac{x_k^+ - x_j^-}{x_k^- - x_j^+} \prod_{j=1}^{M_L^{II}} \frac{x_k^- - y_j}{x_k^+ - y_j} \prod_{j=1}^{M_R^{II}} \frac{1 - \frac{1}{x_k^- \tilde{y}_j}}{1 - \frac{1}{x_k^+ \tilde{y}_j}} = \\ \prod_{j=1}^{\widehat{M}_L^{II}} \frac{x_k^+ - \hat{y}_j}{x_k^- - \hat{y}_j} \prod_{j=1}^{\widehat{M}_R^{II}} \frac{1 - \frac{1}{x_k^+ \tilde{\hat{y}}_j}}{1 - \frac{1}{x_k^- \tilde{\hat{y}}_j}} \prod_{j=1}^{M_R^I} \frac{1 - \frac{1}{x_k^- \tilde{x}_j^-}}{1 - \frac{1}{x_k^+ \tilde{x}_j^+}}, \end{aligned} \quad (\text{A.80})$$

where the exponent of x_k^+/x_k^- is in fact 0. With the help of this substitution, the Bethe equation for left excitations can thus be rewritten as

$$\left(\frac{x_k^+}{x_k^-} \right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{M_L^I} \mathcal{S}_{kj}^{\text{LL}} \prod_{j=1}^{\widehat{M}_L^{II}} \frac{x_k^+ - \hat{y}_j}{x_k^- - \hat{y}_j} \prod_{j=1}^{M_R^I} \sqrt{\frac{1 - \frac{1}{x_k^- \tilde{x}_j^-}}{1 - \frac{1}{x_k^+ \tilde{x}_j^+}}} \mathcal{S}_{kj}^{\text{LR}} \prod_{j=1}^{\widehat{M}_R^{II}} \frac{1 - \frac{1}{x_k^- \tilde{\hat{y}}_j}}{1 - \frac{1}{x_k^+ \tilde{\hat{y}}_j}}. \quad (\text{A.81})$$

This equation has precisely the same form as the original equation for the right excitations (6.48).

Similarly, using $\mathcal{P}(1/\tilde{x}_k^+)/\mathcal{P}(1/\tilde{x}_k^-)$, one can obtain the dualised Bethe equation for type $\tilde{1}$ excitations

$$\left(\frac{\tilde{x}_k^+}{\tilde{x}_k^-}\right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{M_R^1} \frac{\tilde{x}_k^+ - \tilde{x}_j^-}{\tilde{x}_k^- - \tilde{x}_j^+} \mathcal{S}_{kj}^{\text{RR}} \prod_{j=1}^{\widehat{M}_R^1} \frac{\tilde{x}_k^- - \widehat{y}_j}{\tilde{x}_k^+ - \widehat{y}_j} \prod_{j=1}^{M_L^1} \sqrt{\frac{1 - \frac{1}{\tilde{x}_k^+ x_j^+}}{1 - \frac{1}{\tilde{x}_k^- x_j^-}}} \mathcal{S}_{kj}^{\text{RL}} \prod_{j=1}^{\widehat{M}_L^1} \frac{1 - \frac{1}{\tilde{x}_k^- y_j}}{1 - \frac{1}{\tilde{x}_k^+ y_j}}, \quad (\text{A.82})$$

which has the same form as the original Bethe equations for type 1 excitations.

We have found a new set of Bethe equations, with the same form of the original ones up to exchanging left with right excitations and substituting (y, K_2) and $(\tilde{y}, K_{\tilde{2}})$ by $(\widehat{y}, \widehat{K}_{\tilde{2}})$ and $(\widehat{\tilde{y}}, \widehat{K}_2)$, respectively.

Duality for the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$. Bethe equations

As argued in section 6.3, there are four different possible gradings in which we can write the all-loop Bethe equations for the $\mathfrak{psu}(1|1)_{\text{c.e.}}^4$ S matrix. Once again, these are related by dualities. Labelling the excitation numbers K_j by the corresponding Cartan elements as in eqs. (6.76–6.77), let us define again the polynomial

$$\mathcal{P}(\xi) = \prod_{j=1}^{K_2} (\xi - x_j^+) \prod_{j=1}^{K_{\tilde{2}}} (\xi - \frac{1}{\tilde{x}_j^-}) - \prod_{j=1}^{K_2} (\xi - x_j^-) \prod_{j=1}^{K_{\tilde{2}}} (\xi - \frac{1}{\tilde{x}_j^+}), \quad (\text{A.83})$$

of degree $n = K_2 + K_{\tilde{2}} - 1$. The Bethe equations for auxiliary roots $y_1, y_3, y_{\tilde{1}}, y_{\tilde{3}}$ can be written respectively as

$$\mathcal{P}(y_1) = 0, \quad \mathcal{P}(y_3) = 0, \quad \mathcal{P}(1/y_{\tilde{1}}) = 0, \quad \mathcal{P}(1/y_{\tilde{3}}) = 0. \quad (\text{A.84})$$

We can choose to dualise either the auxiliary roots $y_1, y_{\tilde{1}}$ or $y_3, y_{\tilde{3}}$. In the first case we consider a set of dual $\widehat{K}_1, \widehat{K}_{\tilde{1}}$ roots such that $K_1 + \widehat{K}_1 = K_2 - 1$ and $K_{\tilde{1}} + \widehat{K}_{\tilde{1}} = K_{\tilde{2}} - 1$. The polynomial can thus be rewritten as

$$\mathcal{P}(\xi) = \xi \prod_{j=1}^{K_1} (\xi - y_{1,j}) \prod_{j=1}^{\widehat{K}_1} (\xi - \widehat{y}_{1,j}) \prod_{j=1}^{K_{\tilde{1}}} \left(\xi - \frac{1}{y_{\tilde{1},j}}\right) \prod_{j=1}^{\widehat{K}_{\tilde{1}}} \left(\xi - \frac{1}{\widehat{y}_{\tilde{1},j}}\right) \quad (\text{A.85})$$

As we did in the previous subsection, we can evaluate the quantities $\frac{\mathcal{P}(x_k^+)}{\mathcal{P}(x_k^-)}$ and $\frac{\mathcal{P}(1/\tilde{x}_k^-)}{\mathcal{P}(1/\tilde{x}_k^+)}$. The resulting identities, plugged into the Bethe ansatz equations, allow us to rewrite them

as

$$1 = \prod_{j=1}^{K_2} \frac{\widehat{y}_{1,k} - x_j^-}{\widehat{y}_{1,k} - x_j^+} \prod_{j=1}^{K_2} \frac{1 - \frac{1}{\widehat{y}_{1,k} x_j^+}}{1 - \frac{1}{\widehat{y}_{1,k} x_j^-}}, \quad (\text{A.86})$$

$$\left(\frac{x_k^+}{x_k^-} \right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{K_2} \frac{1 - \frac{1}{x_k^+ x_j^-}}{1 - \frac{1}{x_k^- x_j^+}} \sigma^2(x_k, x_j) \prod_{j=1}^{\widehat{K}_1} \frac{x_k^+ - \widehat{y}_{1,j}}{x_k^- - \widehat{y}_{1,j}} \prod_{j=1}^{K_3} \frac{x_k^- - y_{3,j}}{x_k^+ - y_{3,j}} \quad (\text{A.87})$$

$$\begin{aligned} & \times \prod_{j=1}^{K_2} \frac{1 - \frac{1}{x_k^+ \tilde{x}_j^-}}{1 - \frac{1}{x_k^- \tilde{x}_j^+}} \widetilde{\sigma}^2(x_k, \tilde{x}_j) \prod_{j=1}^{\widehat{K}_1} \frac{1 - \frac{1}{x_k^+ \widehat{y}_{1,j}}}{1 - \frac{1}{x_k^- \widehat{y}_{1,j}}} \prod_{j=1}^{K_3} \frac{1 - \frac{1}{x_k^- y_{3,j}}}{1 - \frac{1}{x_k^+ y_{3,j}}}, \\ 1 &= \prod_{j=1}^{K_2} \frac{y_{3,k} - x_j^+}{y_{3,k} - x_j^-} \prod_{j=1}^{K_2} \frac{1 - \frac{1}{y_{3,k} x_j^-}}{1 - \frac{1}{y_{3,k} x_j^+}}, \end{aligned} \quad (\text{A.88})$$

$$1 = \prod_{j=1}^{K_3} \frac{\widehat{y}_{1,k} - \tilde{x}_j^+}{\widehat{y}_{1,k} - \tilde{x}_j^-} \prod_{j=1}^{K_2} \frac{1 - \frac{1}{\widehat{y}_{1,k} x_j^-}}{1 - \frac{1}{\widehat{y}_{1,k} x_j^+}}, \quad (\text{A.89})$$

$$\left(\frac{\tilde{x}_k^+}{\tilde{x}_k^-} \right)^\ell = \prod_{\substack{j=1 \\ j \neq k}}^{K_2} \frac{1 - \frac{1}{\tilde{x}_k^+ x_j^-}}{1 - \frac{1}{\tilde{x}_k^- x_j^+}} \sigma^2(\tilde{x}_k, \tilde{x}_j) \prod_{j=1}^{\widehat{K}_1} \frac{\tilde{x}_k^- - \widehat{y}_{1,j}}{\tilde{x}_k^+ - \widehat{y}_{1,j}} \prod_{j=1}^{K_3} \frac{\tilde{x}_k^+ - y_{3,j}}{\tilde{x}_k^- - y_{3,j}} \quad (\text{A.90})$$

$$\begin{aligned} & \times \prod_{j=1}^{K_2} \frac{1 - \frac{1}{\tilde{x}_k^+ x_j^-}}{1 - \frac{1}{\tilde{x}_k^- x_j^+}} \widetilde{\sigma}^2(\tilde{x}_k, x_j) \prod_{j=1}^{\widehat{K}_1} \frac{1 - \frac{1}{\tilde{x}_k^- \widehat{y}_{1,j}}}{1 - \frac{1}{\tilde{x}_k^+ \widehat{y}_{1,j}}} \prod_{j=1}^{K_3} \frac{1 - \frac{1}{\tilde{x}_k^- y_{3,j}}}{1 - \frac{1}{\tilde{x}_k^+ y_{3,j}}}, \\ 1 &= \prod_{j=1}^{K_3} \frac{y_{3,k} - \tilde{x}_j^-}{y_{3,k} - \tilde{x}_j^+} \prod_{j=1}^{K_2} \frac{1 - \frac{1}{y_{3,k} x_j^-}}{1 - \frac{1}{y_{3,k} x_j^+}}. \end{aligned} \quad (\text{A.91})$$

The above equations are the ones that we would have found if we had chosen Φ_{+-}^L, Φ_{+-}^R to be the fields composing the level-II vacuum. Similarly, one could have dualised the auxiliary roots $y_3, y_{\bar{3}}$ and obtained the Bethe equations corresponding to the choice of Φ_{+-}^L, Φ_{-+}^R in the level-II vacuum. We do not write them, since they are equal to the ones written above after exchanging 1 and 3, as expected. Two consecutive dualisations, *i.e.* dualising $y_1, y_{\bar{1}}$ and then $y_3, y_{\bar{3}}$ (in any order) give Bethe equations corresponding to the choice of Φ_{--}^L, Φ_{++}^R in the level-II vacuum. They are equal to the Bethe equations derived in section 6.3 after exchanging $L \leftrightarrow R$.

Samenvatting


In dit proefschrift hebben we gekeken naar bepaalde supersymmetrische snaartheorieën in gekromde ruimtes die de drie-dimensionale anti-De Sitter ruimte, AdS_3 , bevatten. Aan de ene kant zijn deze theorieën van zwaartekracht in drie dimensies, hoewel ze eenvoudiger zijn dan de zwaartekracht in onze wereld, toch rijk aan vele interessante fenomenen. Aan de andere kant horen deze zwaartekrachtstheorieën volgens het holografische principe equivalent te zijn aan een passende kwantumveldentheorie (QFT) met conforme symmetrie in twee dimensies, een CFT_2 . In dit lage aantal dimensies zijn dergelijke conforme veldentheorieën grotendeels vastgelegd door een oneindig-dimensionale symmetrie-algebra, maar blijven toch niet triviaal.

Ons onderzoek beschouwt een bepaald regime, de 't Hooft limiet, waar de snaren zich vrij bewegen. De natuurlijke observabele in dit regime is het energiespectrum van de snaar, dat via holografie correspondeert met het spectrum van de voortbrenger van dilataties in een bepaalde sector van de CFT. Perturbatieve berekeningen van het energiespectrum van de snaar zijn mogelijk, in wezen door de vrije snaartheorie te beschouwen als een niet-lineair σ -model (NLSM) van het “wereldvlak” van de snaar (meetkundig een tweedimensionale cylinder) naar een “doelruimte” (achtergrond) van de vorm $\text{AdS}_3 \times \mathcal{M}$, waar \mathcal{M} een variëteit zonder rand is. Hier zal \mathcal{M} een product van cirkels en drie-dimensionale bollen zijn, namelijk $\mathcal{M} = \text{S}^3 \times \text{T}^4$ of $\mathcal{M} = \text{S}^3 \times \text{S}^3 \times \text{S}^1$.

Wat deze theorieën erg interessant maakt is onze overtuiging dat alle observabelen op efficiënte wijze kunnen worden berekend voor elke waarde van de koppelingsconstante, zonder terug te vallen op perturbatietheorie—dit is mogelijk dankzij verborgen symmetrieën van de theorie die resulteren in haar zogenoemde *integreerbaarheid*. Het belangrijkste resultaat van dit proefschrift is een overweldigende hoeveelheid bewijs dat de beschouwde $\text{AdS}_3/\text{CFT}_2$ -duale modellen integreerbaar zijn op kwantumniveau.

Het concept integreerbaarheid

In een klassieke theorie betekent integreerbaarheid dat er oneindig veel bewegingsconstanten zijn, die de bewegingsvergelijkingen oplosbaar maken als ware het lineaire differentiaal- of integraalvergelijkingen. Voor onze NLSMs is dit eind 2009 bewezen door Babichenko,



Figuur 1: Een verstrooiingsproces met drie inkomende deeltjes resulteert in drie uitgaande deeltjes (middelste diagram). In een integreerbare theorie kan dit proces worden beschreven als twee verschillende opeenvolgingen van verstrooiingen van twee deeltjes, die equivalent horen te zijn. Dit is een niet-triviale conditie op de S-matrix van twee deeltjes.

Stefański en Zarembo. Dit is vergelijkbaar met de situatie in de $\text{AdS}_5/\text{CFT}_4$ dualiteit—daar is gevonden dat klassieke vrije snaren in $\text{AdS}_5 \times S^5$ integreerbaar zijn en bovendien dat de kern van het kwantum-spectrale probleem kan worden opgelost via integreerbaarheid. Toch bleven er veel vragen over: de $\text{AdS}_3/\text{CFT}_2$ dualiteit bood nieuwe uitdagingen die niet kunnen worden herleid tot de beter begrepen $\text{AdS}_5/\text{CFT}_4$ versie. Om deze uitdagingen goed te kunnen beschrijven zullen we nu het idee van integreerbaarheid in een kwantumveldentheorie kort samenvatten.

In QFT kunnen we integreerbaarheid zien als factorisatie van het verstrooiingsproces en het ontbreken van de vorming van nieuwe deeltjes. In een integreerbare theorie verstrooien M deeltjes naar M deeltjes en kan een $M \rightarrow M$ verstrooiingsproces precies gezien worden als een opeenvolging van $2 \rightarrow 2$ verstrooiingen. Dit is een zeer niet-triviaal feit, zoals aangegeven in figuur 1. In een dergelijke theorie volstaat het dus om de $2 \rightarrow 2$ verstrooiings-matrix (S-matrix) \mathbb{S} uit te rekenen, om elk verstrooiingsproces te beschrijven. Bovendien is het zo dat \mathbb{S} vaak (bijna) volledig is vastgelegd door symmetrieën. In de praktijk draait men deze redenering om: onder de aanname dat de theorie integreerbaar is rekent men \mathbb{S} uit en laat men zien dat het resultaat voldoet aan een aantal consistentie-checks en overeenkomt met perturbatieve berekeningen.

In ons geval is het NLSM gedefinieerd op een cylinder met (eindige) ruimtelijke omtrek ℓ , waar het concept van asymptotische toestanden niet bestaat. Om deze reden moeten we eerst de “decompactificatielimiet” $\ell \rightarrow \infty$ beschouwen, waarin de cylinder een vlak wordt. In dit domein kan de S-matrix gedefinieerd worden. Met deze S-matrix is het mogelijk het energiespectrum te vinden door te eisen dat de verstrooiings-golffuncties ℓ -periodiek zijn. Hieruit volgen kwantisatiecondities voor de impulsen van de (quasi)deeltjes en vervolgens het energiespectrum via de dispersierelatie. Wikkelingseffecten in de vorm

van deeltjes die meerdere keren om de cylinder heen bewegen zijn een subtiel punt, maar ook deze passen in het raamwerk van integreerbaarheid.

Ons geval onderscheidt zich van $\text{AdS}_5/\text{CFT}_4$ in hoofdzakelijk drie opzichten. De duidelijkste is dat onze theorie minder manifeste symmetrieën heeft dan haar hoger dimensionale tegenhanger, waarbij een deel van de continue (Lie) symmetrieën vervangen wordt door een discrete symmetrie. Vervolgens is er weinig bekend over de duale CFT; in het bijzonder is het daar niet mogelijk perturbatieve berekeningen uit te voeren. Ten slotte, als gevolg van de vlakke richtingen in de meetkunde van de doelruimte (de cirkels) bevat het excitatiespectrum van het NLSM massaloze deeltjes die een belangrijk technisch obstakel vormden in de integreerbare aanpak. Desondanks beargumenteren we in dit proefschrift dat deze technische complicaties kunnen worden opgelost en dat de constructie gebaseerd op integreerbaarheid nog steeds kan worden toegepast.

Onze resultaten binnen $\text{AdS}_3/\text{CFT}_2$

Onze resultaten bestaan voornamelijk uit het opzetten en implementeren van de integreerbaarheidsaanpak zoals hierboven beschreven, in de context van $\text{AdS}_3/\text{CFT}_2$. We beschrijven het simpelste geval in deze context in detail, namelijk de massieve sector van $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$.


We vinden de precieze vorm van de $2 \rightarrow 2$ S-matrix door het bestuderen van de symmetrieën van de wereldvlakexcitaties en bekijken enkele van haar analytische eigenschappen. Ook vinden we hetzelfde resultaat vanuit een duaal oogpunt, waar het wereldvlak vervangen is door een spinketen en men spingolven verstrooit. We geven verder de kwantisatiecondities voor de impulsen waarmee het energiespectrum gevonden kan worden. We controleren dat onze procedure consistent is met de aangenomen integreerbaarheid en dat het de perturbatieve berekeningen die in deze context zijn gedaan reproduceert, welke tot de tweede lus orde in de QFT gaan.

In het laatste hoofdstuk van dit proefschrift beschrijven we hoe een dergelijke aanpak kan worden uitgebreid naar de ingewikkeldere $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ achtergrond en hoe massaloze deeltjes hier binnen passen, met referenties naar de originele artikelen van de auteur voor meer details.

Figuur 2 geeft een leuke weergave van een aantal van onze belangrijkste resultaten. Dit figuur illustreert de structuur van de vergelijkingen die het energiespectrum van de massieve deeltjes in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ en $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ beschrijven, op een wijze waarbij de symmetrieën van deze theorieën zich manifesteren.

Vooruitzicht

Het integreerbaarheidsplaatje voor $\text{AdS}_3/\text{CFT}_2$ is nog niet af. Wij zijn desondanks van mening dat de resultaten die we hier hebben gepresenteerd een volledig beeld dichtbij



Figuur 2: Een grafische weergave van de kwantisatiecondities op de impulsen van deeltjes (Bethe-Yang vergelijkingen). De verschillende soorten deeltjes worden weergegeven door de knopen in het Dynkin diagram van de supersymmetrie algebra van beide theorieën ($\mathfrak{psu}(1, 1|2)^2$ en $\mathfrak{d}(2, 1; \alpha)^2$ respectievelijk). De lijnen geven verscheidene interactietermen in de Bethe-Yang vergelijkingen weer.

brengen en dat we in de nabije toekomst een complete beschrijving van het spectrum van snaren in $\text{AdS}_3 \times \text{S}^3 \times \text{T}^4$ en $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ zullen hebben, inclusief de wikkeleffecten.

In onze optiek is dit geen einddoel, maar juist het beginpunt van nieuwe en spannende ontwikkelingen—op dit moment lijken er drie het meest uitnodigend. Ten eerste is het mogelijk de achtergronden die wij hier hebben beschouwd te realiseren via een één-parameter familie van snaartheorieën—de technische benaming is dat ze worden ondersteund door een mix van verschillende fluxen. Hier hebben we ons geconcentreerd op een specifiek punt in deze familie, maar er zijn aanwijzingen dat het betreffende NLSM integreerbaar zou kunnen zijn binnen de hele familie. De oplossing van dit probleem zou nieuwe inzichten geven in de eigenschappen van snaartheorie en in het bijzonder in dualiteiten. Een tweede, gerelateerde onderzoeksrichting is het begrijpen van de relatie tussen de behouden grootheden die in de integreerbaarheidsaanpak naar boven komen en de behouden grootheden die de oneindig-dimensionale symmetrie-algebra van de CFT_2 vormen. Opmerkelijk genoeg kan men hier aan twee conforme veldentheorieën denken: één die dual is aan de snaartheorie op AdS_3 en één op het wereldvlak van de snaar zelf. Tot slot is het mogelijk dat integreerbaarheid behouden blijft onder andere vervormingen van de achtergrond-geometrie (zoals bijvoorbeeld de ruimte uitdelen door de actie van een discrete groep), net als in $\text{AdS}_5/\text{CFT}_4$. Dit is met name interessant omdat er zwarte gaten in AdS_3 bestaan die min of meer van deze vorm zijn en ondanks hun eenvoud interessante fysische eigenschappen hebben. Het toepassen van integreerbaarheid in deze gevallen biedt wellicht een belangrijke stap voorwaarts in ons begrip van zwarte gaten en dus in de zwaartekracht.

Curriculum vitae

I was born on December the Second, 1987, in Como, Italy. There, I attended the *Liceo Scientifico Paolo Giovio* where I graduated in July 2005 with highest marks, one year in advance for outstanding merits.

I have attended the *Università degli Studi di Padova* in Padova, Italy from 2005 to 2010, where I obtained my Bachelor in Physics in July 2008 and my Master in Physics in July 2010, both with highest marks. I also attended the *Scuola Galileiana di Studi Superiori* in Padova, from 2005 to 2010, where I graduated in October 2010 with highest marks. During those years, I have been a visiting student at the *Max-Planck-Institut für Quantenoptik* in Garching bei München, Germany in the group of Prof. Dr. Juan Ignacio Cirac, and at the Perimeter Institut for Theoretical Physics in Waterloo, Canada, in the group of Prof. Dr. Lee Smolin.

From October 2010 to April 2014, I have been a doctoral student at the *Universiteit Utrecht*, under the supervision of Prof. Dr. Gleb Arutyunov. I have have been awarded the *Dutch Research School in Theoretical Physics* prize for best poster presentation at the national “Trends in Theory” symposium in Dalfsen in May 2013, and the *Italian Physical Society* “Vito Volterra” young researcher prize in September 2013.

Since January 2014, I hold a three-year early stage researcher position at the *Humboldt-Universität zu Berlin* in Berlin, Germany, in the group of Prof. Dr. Matthias Staudacher.

Acknowledgements

I would like to thank Gleb Arutyunov for his supervision and encouragement throughout my doctoral studies. To me he has always been a guide to and an example of how to make good Physics, besides being an extraordinary person.

This thesis represents the completion of a journey that started because of a passion for knowledge. I would like to thank those that helped me to keep that passion alive.

Firstly, thanks to all the people I had to pleasure to collaborate with: Gleb, Riccardo Borsato, Franco Cardin, Javier Cerrillo, Juan Ignacio Cirac, Sergey Frolov, Razvan Gurau, Tim Koslowski, Giuseppe de Marco, Olof Ohlsson Sax, Vincent Rivasseau, Norbert Schuch, Bogdan Stefański, Stijn van Tongeren and Alessandro Torrielli. Thanks in particular to Riccardo, Olof, Bogdan and Alessandro for the long and fruitful collaboration that lead to the results described here.

Thanks to many other friends and colleagues with whom I also greatly enjoyed discussing. In particular, my understanding of integrability benefited from interacting with Zoltan Bajnok, Lorenzo Bianchi, Diego Bombardelli, Alessandra Cagnazzo, Ben Hoare, Valentina Forini, Wellington Galleas, Grisha Korchemsky, Laura Koster, Fedor Levkovich-Maslyuk, Tom Lloyd, Takuya Matsumoto, Andrej Stepanchuk, Per Sundin, Ryo Suzuki, Arkady Tseytlin and Kostya Zarembo. More generally, it was a pleasure to talk about Physics (and Mathematics) with Gian Maria dall'Ara, Jogundas Armaitis, Marco Baggio, Dario Benedetti, Antonio Bassetto, Paolo Ciatti, Franz Ciceri, Marcos Crichigno, Thessa Fokkema, Nava Gaddam, Dražen Glavan, Alessandra Gnecci, Vivian Jacobs, Stefanos Katmadas, Kurt Lechner, Marco di Liberto, Gustavo Lucena-Gómez, Vito Mandorino, Pieralberto Marchetti, Enrico Massa, Liuba Mazzanti, Daniele Oriti, Paolo Pasti, Valentina Pozzoli, Andrea Prinsloo, Diego Redigolo, Wilke van der Schee, Simone Speziale, Chiara Toldo, Mario Tonin, Thomas Troppenz, Stefan Vandoren, Oscar Varela, Bram Wouters and Tobias Zingg.

Special thanks to Alessandro, Bogdan, Olof, Riccardo and Stijn for their comments on some of the topics presented here, and in particular to Bogdan for initial collaboration on the topics that evolved into part of chapter 2. Thanks to Gustavo for educating me on the topic of higher-spin theories, to Jules, Laura and Stijn for help with the Dutch translations and to Riccardo for helping me with uncountable practicalities.

Thanks also to the people at ITF who made my stay in Utrecht a very pleasant one: to the students, post-docs and professors, many of whom have already been mentioned above, and to the secretaries and to our manager Joost. Thanks to the many friends I have met there, inside and outside of the University, to the friends met in Perugia, in Padova and to the ones in Como: it is always good to spend time with you. Particular thanks to Wilke and Oscar: it was a pleasure to share an office with you.

Thanks to Maria.

Thanks to Fiorella.

Thanks to Almut.

Thanks to my family.

Thanks and apologies to whom this thesis is dedicated to, to my brother



MATA
SONDINI

References

- [S1] R. Borsato, O. Ohlsson Sax, A. Sfondrini and B. Stefański, Jr., “All-loop worldsheet S matrix for $AdS_3 \times S^3 \times T^4$ ”, [arxiv:1403.4543](#).
- [S2] R. Borsato, O. Ohlsson Sax, A. Sfondrini, B. Stefański, Jr. and A. Torrielli, “Dressing phases of AdS_3/CFT_2 ”, *Phys.Rev. D* **88**, 066004 (2013), [arxiv:1306.2512](#).
- [S3] R. Borsato, O. Ohlsson Sax, A. Sfondrini, B. Stefański, Jr. and A. Torrielli, “The all-loop integrable spin-chain for strings on $AdS_3 \times S^3 \times T^4$: the massive sector”, *JHEP* **1308**, 043 (2013), [arxiv:1303.5995](#).
- [S4] R. Borsato, O. Ohlsson Sax and A. Sfondrini, “All-loop Bethe ansatz equations for AdS_3/CFT_2 ”, *JHEP* **1304**, 116 (2013), [arxiv:1212.0505](#).
- [S5] R. Borsato, O. Ohlsson Sax and A. Sfondrini, “A dynamic $\mathfrak{su}(1|1)^2$ S -matrix for AdS_3/CFT_2 ”, *JHEP* **1304**, 113 (2013), [arxiv:1211.5119](#).
- [S6] R. Gurau, V. Rivasseau and A. Sfondrini, “Renormalization: an advanced overview”, [arxiv:1401.5003](#).
- [S7] A. Sfondrini, “An introduction to universality and renormalization group techniques”, *PoS ModaveVIII*, 005 (2012), [arxiv:1210.2262](#).
- [S8] G. Arutyunov, S. Frolov and A. Sfondrini, “Exceptional operators in $\mathcal{N} = 4$ super Yang-Mills”, *JHEP* **1209**, 006 (2012), [arxiv:1205.6660](#).
- [S9] A. Sfondrini and S. J. van Tongeren, “Lifting asymptotic degeneracies with the Mirror TBA”, *JHEP* **1109**, 050 (2011), [arxiv:1106.3909](#).
- [S10] A. Sfondrini and T. A. Koslowski, “Functional renormalization of noncommutative scalar field theory”, *Int.J.Mod.Phys. A* **26**, 4009 (2011), [arxiv:1006.5145](#).
- [S11] F. Cardin, G. De Marco and A. Sfondrini, “Finite reduction and Morse index estimates for mechanical systems”, *Nonlinear Differential Equations and Applications* **18**, 557 (2011), [arxiv:1004.1133](#).
- [S12] A. Sfondrini, J. Cerrillo, N. Schuch and J. I. Cirac, “Simulating two- and three-dimensional frustrated quantum systems with string-bond states”, *Phys. Rev. B* **81**, 214426 (2010), [arxiv:0908.4036](#).

References (cont.)

- [1] J. D. Bekenstein, “*Black holes and entropy*”, *Phys.Rev.* D7, 2333 (1973).
- [2] J. M. Bardeen, B. Carter and S. Hawking, “*The four laws of black hole mechanics*”, *Commun.Math.Phys.* 31, 161 (1973).
- [3] S. Hawking, “*Black hole explosions?*”, *Nature* 248, 30 (1974).
- [4] S. Hawking, “*Particle creation by black holes*”, *Commun.Math.Phys.* 43, 199 (1975).
- [5] J. D. Bekenstein, “*A universal upper Bound on the entropy to energy ratio for bounded systems*”, *Phys.Rev.* D23, 287 (1981).
- [6] G. 't Hooft, “*Dimensional reduction in quantum gravity*”, [gr-qc/9310026](#).
- [7] R. Bousso, “*The holographic principle*”, *Rev.Mod.Phys.* 74, 825 (2002), [hep-th/0203101](#).
- [8] J. D. Brown and M. Henneaux, “*Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity*”, *Commun.Math.Phys.* 104, 207 (1986).
- [9] L. Susskind, “*The world as a hologram*”, *J.Math.Phys.* 36, 6377 (1995), [hep-th/9409089](#).
- [10] J. M. Maldacena, “*The large- N limit of superconformal field theories and supergravity*”, *Adv.Theor.Math.Phys.* 2, 231 (1998), [hep-th/9711200](#).
- [11] E. Witten, “*Anti-de Sitter space and holography*”, *Adv.Theor.Math.Phys.* 2, 253 (1998), [hep-th/9802150](#).
- [12] S. Gubser, I. R. Klebanov and A. M. Polyakov, “*Gauge theory correlators from noncritical string theory*”, *Phys.Lett.* B428, 105 (1998), [hep-th/9802109](#).
- [13] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “*Large- N field theories, string theory and gravity*”, *Phys.Rept.* 323, 183 (2000), [hep-th/9905111](#).
- [14] G. 't Hooft, “*A planar diagram theory for strong interactions*”, *Nucl.Phys.* B72, 461 (1974).
- [15] J. Minahan and K. Zarembo, “*The Bethe ansatz for $\mathcal{N}=4$ superYang-Mills*”, *JHEP* 0303, 013 (2003), [hep-th/0212208](#).
- [16] N. Beisert, C. Kristjansen and M. Staudacher, “*The Dilatation operator of conformal $\mathcal{N}=4$ superYang-Mills theory*”, *Nucl.Phys.* B664, 131 (2003), [hep-th/0303060](#).
- [17] N. Beisert, “*The complete one loop dilatation operator of $\mathcal{N}=4$ superYang-Mills theory*”, *Nucl.Phys.* B676, 3 (2004), [hep-th/0307015](#).
- [18] N. Beisert, “*The Dilatation operator of $\mathcal{N}=4$ super Yang-Mills theory and integrability*”, *Phys.Rept.* 405, 1 (2004), [hep-th/0407277](#).
- [19] I. Bena, J. Polchinski and R. Roiban, “*Hidden symmetries of the $AdS_5 \times S^5$ superstring*”, *Phys.Rev.* D69, 046002 (2004), [hep-th/0305116](#).
- [20] D. M. Hofman and J. M. Maldacena, “*Giant magnons*”, *J.Phys.* A39, 13095 (2006), [hep-th/0604135](#).
- [21] G. Arutyunov, S. Frolov, J. Russo and A. A. Tseytlin, “*Spinning strings in $AdS_5 \times S^5$ and integrable systems*”, *Nucl.Phys.* B671, 3 (2003), [hep-th/0307191](#).

- [22] A. A. Tseytlin, “*Spinning strings and AdS/CFT duality*”, hep-th/0311139.
- [23] G. Arutyunov, J. Russo and A. A. Tseytlin, “*Spinning strings in $AdS_5 \times S^5$: new integrable system relations*”, Phys.Rev. D69, 086009 (2004), hep-th/0311004.
- [24] V. Kazakov, A. Marshakov, J. Minahan and K. Zarembo, “*Classical/quantum integrability in AdS/CFT*”, JHEP 0405, 024 (2004), hep-th/0402207.
- [25] G. Arutyunov, S. Frolov and M. Staudacher, “*Bethe ansatz for quantum strings*”, JHEP 0410, 016 (2004), hep-th/0406256.
- [26] N. Beisert, “*The $SU(2|2)$ dynamic S-matrix*”, Adv.Theor.Math.Phys. 12, 945 (2008), hep-th/0511082.
- [27] N. Beisert and M. Staudacher, “*Long-range $psu(2, 2|4)$ Bethe ansatze for gauge theory and strings*”, Nucl.Phys. B727, 1 (2005), hep-th/0504190.
- [28] N. Beisert, B. Eden and M. Staudacher, “*Transcendentality and crossing*”, J.Stat.Mech. 0701, P01021 (2007), hep-th/0610251.
- [29] G. Arutyunov and S. Frolov, “*Foundations of the $AdS_5 \times S^5$ superstring. Part I*”, J.Phys. A42, 254003 (2009), arxiv:0901.4937.
- [30] N. Beisert, C. Ahn, L. F. Alday, Z. Bajnok, J. M. Drummond et al., “*Review of AdS/CFT integrability: an overview*”, Lett.Math.Phys. 99, 3 (2012), arxiv:1012.3982.
- [31] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “ *$\mathcal{N}=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*”, JHEP 0810, 091 (2008), arxiv:0806.1218.
- [32] T. Klose, “*Review of AdS/CFT integrability, chapter IV.3: $\mathcal{N}=6$ Chern-Simons and strings on $AdS_4 \times CP^3$* ”, Lett.Math.Phys. 99, 401 (2012), arxiv:1012.3999.
- [33] M. Banados, C. Teitelboim and J. Zanelli, “*The black hole in three-dimensional space-time*”, Phys.Rev.Lett. 69, 1849 (1992), hep-th/9204099.
- [34] M. Banados, M. Henneaux, C. Teitelboim and J. Zanelli, “*Geometry of the (2+1) black hole*”, Phys.Rev. D48, 1506 (1993), gr-qc/9302012.
- [35] S. Carlip, “*The (2+1)-Dimensional black hole*”, Class.Quant.Grav. 12, 2853 (1995), gr-qc/9506079.
- [36] A. Strominger and C. Vafa, “*Microscopic origin of the Bekenstein-Hawking entropy*”, Phys.Lett. B379, 99 (1996), hep-th/9601029.
- [37] A. Strominger, “*Black hole entropy from near horizon microstates*”, JHEP 9802, 009 (1998), hep-th/9712251.
- [38] J. L. Cardy, “*Operator content of two-dimensional conformally invariant theories*”, Nucl.Phys. B270, 186 (1986).
- [39] S. Hyun, “*U duality between three-dimensional and higher-dimensional black holes*”, J.Korean Phys.Soc. 33, S532 (1998), hep-th/9704005.
- [40] K. Sfetsos and K. Skenderis, “*Microscopic derivation of the Bekenstein-Hawking entropy formula for nonextremal black holes*”, Nucl.Phys. B517, 179 (1998), hep-th/9711138.

- [41] A. Giveon, D. Kutasov and N. Seiberg, “Comments on string theory on AdS_3 ”, *Adv.Theor.Math.Phys.* 2, 733 (1998), [hep-th/9806194](#).
- [42] J. M. Maldacena and H. Ooguri, “Strings in AdS_3 and $SL(2, \mathbb{R})$ WZW model 1. The spectrum”, *J.Math.Phys.* 42, 2929 (2001), [hep-th/0001053](#).
- [43] J. M. Maldacena, H. Ooguri and J. Son, “Strings in AdS_3 and the $SL(2, \mathbb{R})$ WZW model. 2. Euclidean black hole”, *J.Math.Phys.* 42, 2961 (2001), [hep-th/0005183](#).
- [44] J. M. Maldacena and H. Ooguri, “Strings in AdS_3 and the $SL(2, \mathbb{R})$ WZW model. 3. Correlation functions”, *Phys.Rev.* D65, 106006 (2002), [hep-th/0111180](#).
- [45] F. Larsen and E. J. Martinec, “ $U(1)$ charges and moduli in the $D1$ - $D5$ system”, *JHEP* 9906, 019 (1999), [hep-th/9905064](#).
- [46] N. Seiberg and E. Witten, “The $D1 / D5$ system and singular CFT”, *JHEP* 9904, 017 (1999), [hep-th/9903224](#).
- [47] N. Berkovits, C. Vafa and E. Witten, “Conformal field theory of AdS background with Ramond-Ramond flux”, *JHEP* 9903, 018 (1999), [hep-th/9902098](#).
- [48] S. Elitzur, O. Feinerman, A. Giveon and D. Tsabar, “String theory on $AdS_3 \times S^3 \times S^3 \times S^1$ ”, *Phys.Lett.* B449, 180 (1999), [hep-th/9811245](#).
- [49] S. Gukov, E. Martinec, G. W. Moore and A. Strominger, “The search for a holographic dual to $AdS_3 \times S^3 \times S^3 \times S^1$ ”, *Adv.Theor.Math.Phys.* 9, 435 (2005), [hep-th/0403090](#).
- [50] A. Babichenko, B. Stefański, Jr. and K. Zarembo, “Integrability and the AdS_3/CFT_2 correspondence”, *JHEP* 1003, 058 (2010), [arxiv:0912.1723](#).
- [51] P. Sundin and L. Wulff, “Classical integrability and quantum aspects of the $AdS_3 \times S^3 \times S^3 \times S^1$ superstring”, *JHEP* 1210, 109 (2012), [arxiv:1207.5531](#).
- [52] A. Cagnazzo and K. Zarembo, “ B -field in AdS_3/CFT_2 Correspondence and Integrability”, *JHEP* 1211, 133 (2012), [arxiv:1209.4049](#).
- [53] R. Metsaev and A. A. Tseytlin, “Type IIB superstring action in $AdS_5 \times S^5$ background”, *Nucl.Phys.* B533, 109 (1998), [hep-th/9805028](#).
- [54] R. Metsaev, C. B. Thorn and A. A. Tseytlin, “Light cone superstring in AdS space-time”, *Nucl.Phys.* B596, 151 (2001), [hep-th/0009171](#).
- [55] G. Arutyunov and S. Frolov, “Integrable Hamiltonian for classical strings on $AdS_5 \times S^5$ ”, *JHEP* 0502, 059 (2005), [hep-th/0411089](#).
- [56] S. Frolov, J. Plefka and M. Zamaklar, “The $AdS_5 \times S^5$ superstring in light-cone gauge and its Bethe equations”, *J.Phys.* A39, 13037 (2006), [hep-th/0603008](#).
- [57] G. Arutyunov, S. Frolov, J. Plefka and M. Zamaklar, “The off-shell symmetry algebra of the light-cone $AdS_5 \times S^5$ superstring”, *J.Phys.* A40, 3583 (2007), [hep-th/0609157](#).
- [58] G. Arutyunov, S. Frolov and M. Zamaklar, “The Zamolodchikov-Faddeev algebra for $AdS_5 \times S^5$ superstring”, *JHEP* 0704, 002 (2007), [hep-th/0612229](#).
- [59] T. Klose, T. McLoughlin, J. Minahan and K. Zarembo, “World-sheet scattering in $AdS_5 \times S^5$ at two loops”, *JHEP* 0708, 051 (2007), [arxiv:0704.3891](#).

- [60] C.-N. Yang, “Some exact results for the many body problems in one dimension with repulsive delta function interaction”, *Phys.Rev.Lett.* 19, 1312 (1967).
- [61] H. Bethe, “On the theory of metals. 1. Eigenvalues and eigenfunctions for the linear atomic chain”, *Z.Phys.* 71, 205 (1931).
- [62] E. H. Fradkin, “*Field theories of condensed matter systems*”, Addison-Wesley (1991).
- [63] M. Kruczenski, “Spin chains and string theory”, *Phys.Rev.Lett.* 93, 161602 (2004), [hep-th/0311203](#).
- [64] M. Kruczenski, A. Ryzhov and A. A. Tseytlin, “Large-spin limit of $AdS_5 \times S^5$ string theory and low-energy expansion of ferromagnetic spin chains”, *Nucl.Phys.* B692, 3 (2004), [hep-th/0403120](#).
- [65] N. Beisert, R. Hernandez and E. Lopez, “A crossing-symmetric phase for $AdS_5 \times S^5$ strings”, *JHEP* 0611, 070 (2006), [hep-th/0609044](#).
- [66] B. Hoare and A. Tseytlin, “Massive S -matrix of $AdS_3 \times S^3 \times T^4$ superstring theory with mixed 3-form flux”, *Nucl.Phys.* B873, 395 (2013), [arxiv:1304.4099](#).
- [67] B. Hoare, A. Stepanchuk and A. Tseytlin, “Giant magnon solution and dispersion relation in string theory in $AdS_3 \times S^3 \times T^4$ with mixed flux”, [arxiv:1311.1794](#).
- [68] J. Ambjorn, R. A. Janik and C. Kristjansen, “Wrapping interactions and a new source of corrections to the spin-chain/string duality”, *Nucl.Phys.* B736, 288 (2006), [hep-th/0510171](#).
- [69] M. Lüscher, “Volume dependence of the energy spectrum in massive quantum field theories. 1. Stable particle states”, *Commun.Math.Phys.* 104, 177 (1986).
- [70] M. Lüscher, “Volume dependence of the energy spectrum in massive quantum field theories. 2. Scattering states”, *Commun.Math.Phys.* 105, 153 (1986).
- [71] Z. Bajnok and R. A. Janik, “Four-loop perturbative Konishi from strings and finite size effects for multiparticle states”, *Nucl.Phys.* B807, 625 (2009), [arxiv:0807.0399](#).
- [72] C.-N. Yang and C. Yang, “Thermodynamics of one-dimensional system of bosons with repulsive delta function interaction”, *J.Math.Phys.* 10, 1115 (1969).
- [73] A. Zamolodchikov, “Thermodynamic Bethe ansatz in relativistic models. Scaling three-state Potts and Lee-Yang models.”, *Nucl.Phys.* B342, 695 (1990).
- [74] G. Arutyunov and S. Frolov, “On string S -matrix, bound states and TBA”, *JHEP* 0712, 024 (2007), [arxiv:0710.1568](#).
- [75] G. Arutyunov and S. Frolov, “The dressing factor and crossing equations”, *J.Phys.* A42, 425401 (2009), [arxiv:0904.4575](#).
- [76] G. Arutyunov and S. Frolov, “String hypothesis for the $AdS_5 \times S^5$ mirror”, *JHEP* 0903, 152 (2009), [arxiv:0901.1417](#).
- [77] G. Arutyunov and S. Frolov, “Thermodynamic Bethe ansatz for the $AdS_5 \times S^5$ mirror model”, *JHEP* 0905, 068 (2009), [arxiv:0903.0141](#).
- [78] D. Bombardelli, D. Fioravanti and R. Tateo, “Thermodynamic Bethe ansatz for planar AdS/CFT : a proposal”, *J.Phys.* A42, 375401 (2009), [arxiv:0902.3930](#).

- [79] N. Gromov, V. Kazakov, A. Kozak and P. Vieira, “*Exact spectrum of anomalous dimensions of planar $\mathcal{N}=4$ supersymmetric Yang-Mills theory: TBA and excited states*”, *Lett.Math.Phys.* 91, 265 (2010), [arxiv:0902.4458](#).
- [80] N. Gromov, V. Kazakov and P. Vieira, “*Exact spectrum of anomalous dimensions of planar $\mathcal{N}=4$ supersymmetric Yang-Mills theory*”, *Phys.Rev.Lett.* 103, 131601 (2009), [arxiv:0901.3753](#).
- [81] A. Cavaglia, D. Fioravanti and R. Tateo, “*Extended Y-system for the AdS_5/CFT_4 correspondence*”, *Nucl.Phys.* B843, 302 (2011), [arxiv:1005.3016](#).
- [82] J. Balog and A. Hegedus, “ *$AdS_5 \times S^5$ mirror TBA equations from Y-system and discontinuity relations*”, *JHEP* 1108, 095 (2011), [arxiv:1104.4054](#).
- [83] R. Suzuki, “*Hybrid NLIE for the mirror $AdS_5 \times S^5$* ”, *J.Phys.* A44, 235401 (2011), [arxiv:1101.5165](#).
- [84] J. Balog and A. Hegedus, “*Hybrid-NLIE for the AdS/CFT spectral problem*”, *JHEP* 1208, 022 (2012), [arxiv:1202.3244](#).
- [85] N. Gromov, V. Kazakov, S. Leurent and D. Volin, “*Quantum spectral curve for AdS_5/CFT_4* ”, [arxiv:1305.1939](#).
- [86] G. Arutyunov and S. Frolov, “*Comments on the mirror TBA*”, *JHEP* 1105, 082 (2011), [arxiv:1103.2708](#).
- [87] G. Arutyunov, S. Frolov and R. Suzuki, “*Exploring the mirror TBA*”, *JHEP* 1005, 031 (2010), [arxiv:0911.2224](#).
- [88] G. Arutyunov, S. Frolov and R. Suzuki, “*Five-loop Konishi from the mirror TBA*”, *JHEP* 1004, 069 (2010), [arxiv:1002.1711](#).
- [89] J. Balog and A. Hegedus, “*Five-loop Konishi from linearized TBA and the XXX magnet*”, *JHEP* 1006, 080 (2010), [arxiv:1002.4142](#).
- [90] G. Arutyunov, S. Frolov and S. J. van Tongeren, “*Bound states in the mirror TBA*”, *JHEP* 1202, 014 (2012), [arxiv:1111.0564](#).
- [91] S. Leurent, D. Serban and D. Volin, “*Six-loop Konishi anomalous dimension from the Y-system*”, *Phys.Rev.Lett.* 109, 241601 (2012), [arxiv:1209.0749](#).
- [92] S. Leurent and D. Volin, “*Multiple zeta functions and double wrapping in planar $\mathcal{N}=4$ SYM*”, *Nucl.Phys.* B875, 757 (2013), [arxiv:1302.1135](#).
- [93] N. Gromov, V. Kazakov and P. Vieira, “*Exact spectrum of planar $\mathcal{N} = 4$ supersymmetric Yang-Mills theory: Konishi dimension at any coupling*”, *Phys.Rev.Lett.* 104, 211601 (2010), [arxiv:0906.4240](#).
- [94] S. Frolov, “*Konishi operator at intermediate coupling*”, *J.Phys.* A44, 065401 (2011), [arxiv:1006.5032](#).
- [95] S. Frolov, “*Scaling dimensions from the mirror TBA*”, *J.Phys.* A45, 305402 (2012), [arxiv:1201.2317](#).
- [96] N. Gromov and P. Vieira, “*The AdS_4/CFT_3 algebraic curve*”, *JHEP* 0902, 040 (2009), [arxiv:0807.0437](#).

- [97] R. A. Janik, “*The $AdS_5 \times S^5$ superstring worldsheet S -matrix and crossing symmetry*”, Phys.Rev. D73, 086006 (2006), [hep-th/0603038](#).
- [98] K. Zarembo, “*Strings on semisymmetric superspaces*”, JHEP 1005, 002 (2010), [arxiv:1003.0465](#).
- [99] K. Zarembo, “*Algebraic curves for integrable string backgrounds*”, [arxiv:1005.1342](#).
- [100] N. Rughoonauth, P. Sundin and L. Wulff, “*Near BMN dynamics of the $AdS_3 \times S^3 \times S^3 \times S^1$ superstring*”, JHEP 1207, 159 (2012), [arxiv:1204.4742](#).
- [101] P. Sundin and L. Wulff, “*Worldsheet scattering in AdS_3/CFT_2* ”, JHEP 1307, 007 (2013), [arxiv:1302.5349](#).
- [102] B. Hoare and A. Tseytlin, “*On string theory on $AdS_3 \times S^3 \times T^4$ with mixed 3-form flux: tree-level S -matrix*”, Nucl.Phys. B873, 682 (2013), [arxiv:1303.1037](#).
- [103] M. Beccaria, F. Levkovich-Maslyuk, G. Macorini and A. Tseytlin, “*Quantum corrections to spinning superstrings in $AdS_3 \times S^3 \times M^4$: determining the dressing phase*”, JHEP 1304, 006 (2013), [arxiv:1211.6090](#).
- [104] M. Beccaria and G. Macorini, “*Quantum corrections to short folded superstring in $AdS_3 \times S^3 \times M^4$* ”, JHEP 1303, 040 (2013), [arxiv:1212.5672](#).
- [105] M. C. Abbott, “*The $AdS_3 \times S^3 \times S^3 \times S^1$ Hernández-López phases: a semiclassical derivation*”, J. Phys. A46, 445401 (2013), [arxiv:1306.5106](#).
- [106] L. Bianchi, V. Forini and B. Hoare, “*Two-dimensional S -matrices from unitarity cuts*”, JHEP 1307, 088 (2013), [arxiv:1304.1798](#).
- [107] O. T. Engelund, R. W. McKeown and R. Roiban, “*Generalized unitarity and the worldsheet S matrix in $AdS_n \times S^n \times M^{10-2n}$* ”, JHEP 1308, 023 (2013), [arxiv:1304.4281](#).
- [108] P. Sundin, “*Worldsheet two- and four-point functions at one loop in $AdS(3) / CFT(2)$* ”, [arxiv:1403.1449](#).
- [109] A. Pakman, L. Rastelli and S. S. Razamat, “*Diagrams for symmetric product orbifolds*”, JHEP 0910, 034 (2009), [arxiv:0905.3448](#).
- [110] O. Ohlsson Sax, B. Stefański, Jr. and A. Torrielli, “*On the massless modes of the AdS_3/CFT_2 integrable systems*”, JHEP 1303, 109 (2013), [arxiv:1211.1952](#).
- [111] T. Lloyd and B. Stefański, Jr., “ *AdS_3/CFT_2 , finite-gap equations and massless modes*”, [arxiv:1312.3268](#).
- [112] O. Ohlsson Sax and B. Stefański, Jr., “*Integrability, spin-chains and the AdS_3/CFT_2 correspondence*”, JHEP 1108, 029 (2011), [arxiv:1106.2558](#).
- [113] A. A. Tseytlin, “*Superstrings in AdS in light cone gauge*”, Int.J.Mod.Phys. A16, 900 (2001), [hep-th/0009226](#).
- [114] M. B. Green and J. H. Schwarz, “*Covariant description of superstrings*”, Phys.Lett. B136, 367 (1984).
- [115] M. T. Grisaru, P. S. Howe, L. Mezincescu, B. Nilsson and P. Townsend, “ *$\mathcal{N} = 2$ superstrings in a supergravity background*”, Phys.Lett. B162, 116 (1985).

- [116] L. Wulff, “*The type II superstring to order θ^4* ”, JHEP 1307, 123 (2013), [arxiv:1304.6422](#).
- [117] G. Arutyunov and S. Frolov, “*Uniform light-cone gauge for strings in $AdS_5 \times S^5$: solving $SU(1|1)$ sector*”, JHEP 0601, 055 (2006), [hep-th/0510208](#).
- [118] G. Arutyunov, S. Frolov and M. Zamaklar, “*Finite-size effects from giant magnons*”, Nucl.Phys. B778, 1 (2007), [hep-th/0606126](#).
- [119] D. E. Berenstein, J. M. Maldacena and H. S. Nastase, “*Strings in flat space and pp waves from $\mathcal{N}=4$ superYang-Mills*”, JHEP 0204, 013 (2002), [hep-th/0202021](#).
- [120] J. Souriau, “*Structure des systèmes dynamiques.*”, Dunod (1970).
- [121] N. Beisert and B. I. Zwiebel, “*On symmetry enhancement in the $\mathfrak{psu}(1,1|2)$ sector of $\mathcal{N}=4$ SYM*”, JHEP 0710, 031 (2007), [arxiv:0707.1031](#).
- [122] J. R. David and B. Sahoo, “*Giant magnons in the $D1$ - $D5$ system*”, JHEP 0807, 033 (2008), [arxiv:0804.3267](#).
- [123] B. Hoare and A. Tseytlin, “*Towards the quantum S -matrix of the Pohlmeyer reduced version of $AdS_5 \times S^5$ superstring theory*”, Nucl.Phys. B851, 161 (2011), [arxiv:1104.2423](#).
- [124] P. Sundin and L. Wulff, “*The low energy limit of the $AdS_3 \times S^3 \times M_4$ spinning string*”, JHEP 1310, 111 (2013), [arxiv:1306.6918](#).
- [125] O. Babelon, D. Bernard and M. Talon, “*Introduction to classical integrable systems*”, Cambridge University Press (2007).
- [126] L. Faddeev, L. Takhtajan and A. Reyman, “*Hamiltonian methods in the theory of solitons*”, Springer (2007).
- [127] M. Gaudin, “*La fonction d’onde de Bethe*”, Masson (1983).
- [128] A. B. Zamolodchikov and A. B. Zamolodchikov, “*Factorized S matrices in two dimensions as the exact solutions of certain relativistic quantum field models*”, Annals Phys. 120, 253 (1979).
- [129] L. Faddeev, “*How algebraic Bethe ansatz works for integrable model*”, [hep-th/9605187](#).
- [130] C. S. Gardner, J. M. Greene, M. D. Kruskal and R. M. Miura, “*Method for solving the Korteweg-de Vries equation*”, Phys. Rev. Lett. 19, 1095 (1967).
- [131] P. D. Lax, “*Integrals of nonlinear equations of evolution and solitary waves*”, Communications on pure and applied mathematics 21, 467 (1968).
- [132] V. Zakharov and L. Faddeev, “*Korteweg-de Vries equation: a completely integrable Hamiltonian system*”, Functional Analysis and Its Applications 5, 280 (1971), <http://dx.doi.org/10.1007/BF01086739>.
- [133] A. Shabat and V. Zakharov, “*Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media*”, Soviet Physics JETP 34, 62 (1972).
- [134] I. Gel’fand and B. Levitan, “*On the determination of a differential equation from its spectral function*”, American Mathematical Society (1955).
- [135] V. Marchenko, “*Sturm-Liouville operators and applications*”, Birkäuser (1986).

- [136] L. Faddeev, “*Quantum completely integral models of field theory*”, *Sov.Sci.Rev.* C1, 107 (1980).
- [137] J. Plefka, F. Spill and A. Torrielli, “*On the Hopf algebra structure of the AdS/CFT S-matrix*”, *Phys.Rev.* D74, 066008 (2006), [hep-th/0608038](#).
- [138] A. Torrielli, “*Review of AdS/CFT integrability, chapter VI.2: Yangian algebra*”, *Lett.Math.Phys.* 99, 547 (2012), [arxiv:1012.4005](#).
- [139] A. Torrielli, “*Yangians, S-matrices and AdS/CFT*”, *J.Phys.* A44, 263001 (2011), [arxiv:1104.2474](#).
- [140] G. Delius, “*Exact S-matrices with affine quantum group symmetry*”, *Nucl.Phys.* B451, 445 (1995), [hep-th/9503079](#).
- [141] N. Beisert, V. Dippel and M. Staudacher, “*A Novel long range spin chain and planar $\mathcal{N}=4$ super Yang-Mills*”, *JHEP* 0407, 075 (2004), [hep-th/0405001](#).
- [142] N. Dorey, D. M. Hofman and J. M. Maldacena, “*On the singularities of the magnon S-matrix*”, *Phys.Rev.* D76, 025011 (2007), [hep-th/0703104](#).
- [143] P. Vieira and D. Volin, “*Review of AdS/CFT Integrability, chapter III.3: the dressing factor*”, *Lett.Math.Phys.* 99, 231 (2012), [arxiv:1012.3992](#).
- [144] R. Hernández and E. López, “*Quantum corrections to the string Bethe ansatz*”, *JHEP* 0607, 004 (2006), [hep-th/0603204](#).
- [145] L. Castillejo, R. Dalitz and F. Dyson, “*Low’s scattering equation for the charged and neutral scalar theories*”, *Phys.Rev.* 101, 453 (1956).
- [146] H.-Y. Chen, N. Dorey and K. Okamura, “*The Asymptotic spectrum of the $\mathcal{N}=4$ super Yang-Mills spin chain*”, *JHEP* 0703, 005 (2007), [hep-th/0610295](#).
- [147] G. Arutyunov and S. Frolov, “*The S-matrix of string bound states*”, *Nucl.Phys.* B804, 90 (2008), [arxiv:0803.4323](#).
- [148] H.-Y. Chen, N. Dorey and K. Okamura, “*Dyonic giant magnons*”, *JHEP* 0609, 024 (2006), [hep-th/0605155](#).
- [149] N. Dorey, “*Magnon bound states and the AdS/CFT correspondence*”, *J.Phys.* A39, 13119 (2006), [hep-th/0604175](#).
- [150] M. C. Abbott, “*Comment on strings in $AdS_3 \times S^3 \times S^3 \times S^1$ at one loop*”, *JHEP* 1302, 102 (2013), [arxiv:1211.5587](#).
- [151] R. Eden, P. Landshoff, D. Olive and J. Polkinghorne, “*The analytic S matrix*”, Cambridge University Press (2002).
- [152] N. Beisert, “*Review of AdS/CFT integrability, chapter VI.1: superconformal symmetry*”, *Lett.Math.Phys.* 99, 529 (2012), [arxiv:1012.4004](#).
- [153] N. Beisert and M. Staudacher, “*The $\mathcal{N}=4$ SYM integrable super spin chain*”, *Nucl.Phys.* B670, 439 (2003), [hep-th/0307042](#).
- [154] N. Beisert, “*The $su(2|3)$ dynamic spin chain*”, *Nucl.Phys.* B682, 487 (2004), [hep-th/0310252](#).

- [155] A. Rej, “*Review of AdS/CFT integrability, chapter I.3: long-range spin chains*”, *Lett.Math.Phys.* 99, 85 (2012), [arxiv:1012.3985](#).
- [156] F. H. Essler and V. E. Korepin, “*A new solution of the supersymmetric T-J model by means of the quantum inverse scattering method*”, [hep-th/9207007](#).
- [157] E. Ogievetsky and P. Wiegmann, “*Factorized S matrix and the Bethe ansatz for simple Lie groups*”, *Phys.Lett.* B168, 360 (1986).
- [158] T. Klose, T. McLoughlin, R. Roiban and K. Zarembo, “*Worldsheet scattering in $AdS_5 \times S^5$* ”, *JHEP* 0703, 094 (2007), [hep-th/0611169](#).
- [159] J. M. Maldacena and I. Swanson, “*Connecting giant magnons to the pp-wave: an interpolating limit of $AdS_5 \times S^5$* ”, *Phys.Rev.* D76, 026002 (2007), [hep-th/0612079](#).
- [160] N. Beisert and T. Klose, “*Long-range $gl(n)$ integrable spin chains and plane-wave matrix theory*”, *J.Stat.Mech.* 0607, P07006 (2006), [hep-th/0510124](#).
- [161] T. Bargheer, N. Beisert and F. Loebbert, “*Long-range deformations for integrable spin chains*”, *J.Phys.* A42, 285205 (2009), [arxiv:0902.0956](#).
- [162] N. Beisert, J. Minahan, M. Staudacher and K. Zarembo, “*Stringing spins and spinning strings*”, *JHEP* 0309, 010 (2003), [hep-th/0306139](#).
- [163] N. Beisert, S. Frolov, M. Staudacher and A. A. Tseytlin, “*Precision spectroscopy of AdS/CFT*”, *JHEP* 0310, 037 (2003), [hep-th/0308117](#).
- [164] N. Gromov and P. Vieira, “*Complete 1-loop test of AdS/CFT*”, *JHEP* 0804, 046 (2008), [arxiv:0709.3487](#).
- [165] N. Gromov, S. Schafer-Nameki and P. Vieira, “*Quantum wrapped giant magnon*”, *Phys.Rev.* D78, 026006 (2008), [arxiv:0801.3671](#).
- [166] J. R. David and B. Sahoo, “*S-matrix for magnons in the D1-D5 system*”, *JHEP* 1010, 112 (2010), [arxiv:1005.0501](#).
- [167] C. Ahn and D. Bombardelli, “*Exact S-matrices for AdS_3/CFT_2* ”, [arxiv:1211.4512](#).
- [168] A. B. Zamolodchikov and A. B. Zamolodchikov, “*Massless factorized scattering and sigma models with topological terms*”, *Nucl.Phys.* B379, 602 (1992).
- [169] P. Fendley, H. Saleur and A. B. Zamolodchikov, “*Massless flows. 1. The Sine-Gordon and $O(n)$ models*”, *Int.J.Mod.Phys.* A8, 5717 (1993), [hep-th/9304050](#).
- [170] P. Fendley, H. Saleur and A. B. Zamolodchikov, “*Massless flows, 2. The Exact S matrix approach*”, *Int.J.Mod.Phys.* A8, 5751 (1993), [hep-th/9304051](#).
- [171] P. Fendley and H. Saleur, “*Massless integrable quantum field theories and massless scattering in (1+1)-dimensions*”, [hep-th/9310058](#).
- [172] L. F. Alday, G. Arutyunov and S. Frolov, “*Green-Schwarz strings in TsT-transformed backgrounds*”, *JHEP* 0606, 018 (2006), [hep-th/0512253](#).
- [173] D. J. Gross and A. Neveu, “*Dynamical symmetry breaking in asymptotically free field theories*”, *Phys.Rev.* D10, 3235 (1974).
- [174] E. Witten, “*Chiral symmetry, the $1/n$ expansion, and the $SU(N)$ Thirring model*”,

- Nucl.Phys. B145, 110 (1978).
- [175] L. Frappat, P. Sorba and A. Sciarrino, “*Dictionary on Lie superalgebras*”, hep-th/9607161.
- [176] F. Gavarini, “*Chevalley Supergroups of type $D(2, 1; \alpha)$* ”, arxiv:1006.0464.
- [177] J. Rahmfeld and A. Rajaraman, “*The GS string action on $AdS_3 \times S^3$ with Ramond-Ramond charge*”, Phys.Rev. D60, 064014 (1999), hep-th/9809164.
- [178] I. Pesando, “*The GS type IIB superstring action on $AdS_3 \times S^3 \times T^4$* ”, JHEP 9902, 007 (1999), hep-th/9809145.
- [179] J. Murugan, P. Sundin and L. Wulff, “*Classical and quantum integrability in AdS_2/CFT_1* ”, JHEP 1301, 047 (2013), arxiv:1209.6062.
- [180] A. Pakman, L. Rastelli and S. S. Razamat, “*A spin chain for the symmetric product CFT_2* ”, JHEP 1005, 099 (2010), arxiv:0912.0959.
- [181] D. Tong, “*The holographic dual of $AdS_3 \times S^3 \times S^3 \times S^1$* ”, arxiv:1402.5135.
- [182] B. Sundborg, “*Stringy gravity, interacting tensionless strings and massless higher spins*”, Nucl.Phys.Proc.Suppl. 102, 113 (2001), hep-th/0103247.
- [183] E. Witten, “*Spacetime reconstruction*”, Talk at the John Schwarz 60th Birthday Symposium, (2001), <http://theory.caltech.edu/jhs60/witten/1.html>.
- [184] E. Sezgin and P. Sundell, “*Massless higher spins and holography*”, Nucl.Phys. B644, 303 (2002), hep-th/0205131.
- [185] A. Achucarro and P. Townsend, “*A Chern-Simons action for three-dimensional anti-de Sitter supergravity theories*”, Phys.Lett. B180, 89 (1986).
- [186] E. Witten, “*(2+1)-dimensional gravity as an exactly soluble system*”, Nucl.Phys. B311, 46 (1988).
- [187] M. Blencowe, “*A consistent interacting massless higher spin field theory in $D = (2 + 1)$* ”, Class.Quant.Grav. 6, 443 (1989).
- [188] A. Campoleoni, S. Fredenhagen, S. Pfenninger and S. Theisen, “*Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields*”, JHEP 1011, 007 (2010), arxiv:1008.4744.
- [189] M. Henneaux and S.-J. Rey, “*Nonlinear W_∞ as asymptotic symmetry of three-dimensional higher spin anti-de Sitter gravity*”, JHEP 1012, 007 (2010), arxiv:1008.4579.
- [190] A. Zamolodchikov, “*Infinite additional symmetries in two-dimensional conformal quantum field theory*”, Theor.Math.Phys. 65, 1205 (1985).
- [191] P. Bouwknegt and K. Schoutens, “*W symmetry in conformal field theory*”, Phys.Rept. 223, 183 (1993), hep-th/9210010.
- [192] M. R. Gaberdiel and R. Gopakumar, “*An AdS_3 dual for minimal model CFTs*”, Phys.Rev. D83, 066007 (2011), arxiv:1011.2986.
- [193] M. R. Gaberdiel and T. Hartman, “*Symmetries of holographic minimal models*”,

- JHEP 1105, 031 (2011), [arxiv:1101.2910](#).
- [194] C. Candu and M. R. Gaberdiel, “*Supersymmetric holography on AdS_3* ”, JHEP 1309, 071 (2013), [arxiv:1203.1939](#).
- [195] M. Henneaux, G. Lucena Gómez, J. Park and S.-J. Rey, “*Super- W_∞ asymptotic symmetry of higher-spin AdS_3 supergravity*”, JHEP 1206, 037 (2012), [arxiv:1203.5152](#).
- [196] M. R. Gaberdiel and R. Gopakumar, “*Minimal model holography*”, J.Phys. A46, 214002 (2013), [arxiv:1207.6697](#).
- [197] M. R. Gaberdiel and R. Gopakumar, “*Large $\mathcal{N} = 4$ holography*”, JHEP 1309, 036 (2013), [arxiv:1305.4181](#).
- [198] M. Gaberdiel, “*Large $\mathcal{N} = 4$ minimal model holography*”, Talk at the Integrability in Gauge and String Theory workshop, (2013), <http://web.science.uu.nl/IGST13/>.
- [199] A. Sagnotti and M. Taronna, “*String lessons for higher-spin interactions*”, Nucl.Phys. B842, 299 (2011), [arxiv:1006.5242](#).
- [200] C.-M. Chang, S. Minwalla, T. Sharma and X. Yin, “*ABJ Triality: from higher spin fields to strings*”, J.Phys. A46, 214009 (2013), [arxiv:1207.4485](#).
- [201] S. Kachru and E. Silverstein, “*4-D conformal theories and strings on orbifolds*”, Phys.Rev.Lett. 80, 4855 (1998), [hep-th/9802183](#).
- [202] A. E. Lawrence, N. Nekrasov and C. Vafa, “*On conformal field theories in four-dimensions*”, Nucl.Phys. B533, 199 (1998), [hep-th/9803015](#).
- [203] O. Lunin and J. M. Maldacena, “*Deforming field theories with $U(1) \times U(1)$ global symmetry and their gravity duals*”, JHEP 0505, 033 (2005), [hep-th/0502086](#).
- [204] S. Frolov, “*Lax pair for strings in Lunin-Maldacena background*”, JHEP 0505, 069 (2005), [hep-th/0503201](#).
- [205] N. Beisert and P. Koroteev, “*Quantum deformations of the one-dimensional Hubbard model*”, J.Phys. A41, 255204 (2008), [arxiv:0802.0777](#).
- [206] N. Beisert, W. Galleas and T. Matsumoto, “*A quantum affine algebra for the deformed Hubbard chain*”, J.Phys. A45, 365206 (2012), [arxiv:1102.5700](#).
- [207] B. Hoare, T. J. Hollowood and J. L. Miramontes, “*q-deformation of the $AdS_5 \times S^5$ superstring S-matrix and its relativistic limit*”, JHEP 1203, 015 (2012), [arxiv:1112.4485](#).
- [208] M. de Leeuw, T. Matsumoto and V. Regelskis, “*The bound state S-matrix of the deformed Hubbard chain*”, JHEP 1204, 021 (2012), [arxiv:1109.1410](#).
- [209] F. Delduc, M. Magro and B. Vicedo, “*An integrable deformation of the $AdS_5 \times S^5$ superstring action*”, [arxiv:1309.5850](#).
- [210] G. Arutyunov, R. Borsato and S. Frolov, “*S-matrix for strings on η -deformed $AdS_5 \times S^5$* ”, [arxiv:1312.3542](#).
- [211] S. J. van Tongeren, “*Integrability of the $AdS_5 \times S^5$ superstring and its deformations*”, [arxiv:1310.4854](#).