



# Fermi National Accelerator Laboratory

FERMILAB-Pub-82/74-EXP  
7180.557  
(Submitted to Phys. Rev. Lett.)

## PRODUCTION OF HIGH TRANSVERSE ENERGY EVENTS IN p-NUCLEUS COLLISIONS AT 400 GeV/c

B. Brown, P. Devenski,<sup>a</sup> S. Gronemeyer, H. Haggerty,  
E. Malamud, and P. Rapp<sup>b</sup>  
Fermi National Accelerator Laboratory, Batavia, Illinois 60510

and

R. Abrams, H. Goldberg, C. Halliwell, F. Lopez,  
S. Margulies, D. McLeod, and J. Solomon  
University of Illinois-Circle Campus, Chicago, Illinois 60680

and

A. Dzierba, J. Florian, R. Heinz, J. Krider, J. Martin,  
D. Petersen, P. Smith, and S. Teige  
Indiana University, Bloomington, Indiana 47405

and

R. Ellsworth,<sup>c</sup> R. Glasser, R. Holmes, L. Myrianthopoulos,<sup>d</sup>  
H. Strobele,<sup>e</sup> G. Yodh, and A. Zieminski<sup>f</sup>  
University of Maryland, College Park, Maryland 20742

and

S. Ahn and T. Watts  
Rutgers University, New Brunswick, New Jersey 08903

October 1982



## Abstract

High transverse energy events produced in proton-nucleus collisions at 400 GeV/c were studied using the Fermilab Multiparticle Spectrometer. The cross sections for such interactions increase more rapidly than the atomic mass number for both small and large acceptance data. The nuclear events are less planar than the corresponding high transverse energy pp events.

It is now widely accepted that the interactions of high-energy particles with nuclear targets provide a unique opportunity to study the space-time development of the hadron production process.<sup>(1)</sup> The hadronic matter formed as a consequence of a primary interaction may interact again with another nucleon long before it has time to convert into hadrons.

In this approach the nucleus is treated as a kind of detector which helps

observe phenomena not accessible to ordinary detectors used in high energy physics. It has also been argued that high energy nuclear experiments provide information on the structure of the incident hadrons.<sup>(1)</sup>

Cross sections for inclusive particle production in hadron-nucleus-reactions have been found<sup>(1-3)</sup> to grow as a power of  $A$ , the number of nucleons in the nucleus:  $d^3\sigma/d^3p \propto A^\alpha$ . The exponent  $\alpha$  depends on the flavor and momentum of the produced particle, in particular on the particle transverse momentum,  $p_t$ . For particles emitted around  $90^\circ$  in the hadron-nucleon center of mass frame, the values of  $\alpha$  increase from 0.8 observed in the low- $p_t$  region<sup>(2)</sup> to values larger than 1 for  $p_t \geq 2$  GeV/c.<sup>(3)</sup> The latter effect, called<sup>(4)</sup> the "anomalous nuclear enhancement", is also observed when cross sections for production of a group of particles (a 'jet') are studied.<sup>(5)</sup> Values of  $\alpha \geq 1$  are often considered as an indication that hard scattering of hadronic constituents plays a dominant role in the production process.<sup>(4,6,7)</sup>

In this paper we investigate the  $A$ -dependence of cross sections for the emission of a certain amount of transverse energy,  $E_t$  ( $E_t \approx \sum |p_t|$  for relativistic secondaries) into a given solid angle,  $\Delta\Omega^*$ , in the proton-nucleon c.m. system. The studied range of transverse energy extends up to 80% of the available proton-nucleon c.m. energy and results are presented for both small and large values of  $\Delta\Omega^*$ . This is the first time that such data have been

available and they should provide new types of sensitive tests for models of hadron-nucleus interactions.

These data came from experiment E557, performed in the Fermilab Multiparticle Spectrometer using 400 GeV protons.

The beam was incident on a 45 cm  $H_2$  target followed downstream by two interchangeable metal foils, of Al, Cu or Pb. The results from the hydrogen data as well as details of the apparatus have already been published.<sup>(8-9)</sup>

In this report we analyze nuclear target data taken with three different triggers. The low  $E_t$  'interacting beam' trigger required only the detection of an inelastic collision. The high  $E_t$  triggers made use of the highly segmented calorimeter<sup>(10)</sup> placed downstream of the magnetic spectrometer, 9.1 m from the nuclear targets; the triggers consisted of the 'interacting beam' trigger with an additional requirement that the  $E_t$  detected in the entire calorimeter (' $2\pi$ ' trigger) or in a smaller region of the calorimeter (' $\pi/3$ ' trigger), exceeded a threshold.

The geometrical acceptance of the apparatus for the ' $2\pi$ ' trigger was estimated to be  $2\pi$  in azimuth for the polar angle range  $47^\circ < \theta^* < 127^\circ$  as measured in the proton-nucleon center-of-mass frame. This corresponds to c.m. rapidity range  $-0.70 < y^* < 0.84$  and to  $\Delta\Omega^* \approx 8.0$  steradians. The c.m. solid angle covered by the ' $\pi/3$ ' trigger was  $\Delta\Omega^* \approx 1.35$  steradians. The acceptances for pA collisions given above were larger by 2% than those for pp interactions.<sup>(8,9)</sup> The difference in the acceptances for the two nuclear foils was negligible. Table I lists types and thicknesses of the nuclear targets as well as the number of events from each used in the analysis.

The magnetic spectrometer was used only to locate the position of the interaction vertex. The distributions of the vertex coordinate along the

incident beam direction,  $z_{\text{vertex}}$ , are shown in Figs. 1a and 1b for the 'interacting beam' data and for the high- $E_t$  ' $2\pi$ ' data, respectively. It is clear from these figures that the relative cross sections for different nuclei depend strongly on the observed  $E_t$ . To determine cross sections for the different targets a fit was performed to the  $z_{\text{vertex}}$  distributions for various regions of  $E_t$ . The fitted curve was a sum of two vertex resolution functions and a constant background. The background term accounted for  $\approx 10\%$  of all events found in the nuclear targets region. An example of such a fit is presented in Fig. 1c.

Cross sections,  $d\sigma/dE_t$ , are shown in Fig. 2 for the large acceptance data (including interacting beam data). The data are consistent with the parametrization  $d\sigma/dE_t = A^\alpha(E_t)$  when the hydrogen data are not included. The pp data lie below extrapolated straight lines drawn through the nuclear points. A similar deviation of the pp data has been also seen for single particle production.<sup>(3)</sup>

The values of the exponent  $\alpha(E_t)$  were determined separately for pairs of nuclear foils for which data were collected simultaneously. This way systematic errors due to threshold effects, calibration of the  $E_t$  scale, and absolute normalization have been minimized. The errors in the target thicknesses, estimates of the background and the acceptance differences (for  $H_2$  and Al) lead to an estimated 0.03-0.05 systematic error in the values of  $\alpha$ . There is very good agreement between values of  $\alpha$  obtained for the Al-Cu and Al-Pb pairs. However, the Al-Cu data are not available for the entire  $E_t$  range.

The variation of  $\alpha$  as a function of  $E_t$  for the ' $2\pi$ ' azimuthal acceptance trigger is shown in Fig. 3a. The exponent  $\alpha$  agrees with the expected value of  $\approx 0.7$  at low  $E_t$ <sup>(1)</sup>, becomes 1 at  $E_t \approx 10$  GeV, and, after rising to  $\approx 1.20$  for  $E_t \geq 15$  GeV, flattens off. A similar dependence of  $\alpha$  on  $E_t$  is

also observed for the smaller acceptance trigger (' $\pi/3$ ') (see Fig. 3b). This behavior of  $\alpha$ , as seen in Fig. 3a and 3b, resembles the dependence of  $\alpha$  on  $p_t$  found for single particle production.<sup>(3)</sup>

The trend of  $\alpha$  to exceed unity has also been reported for the production of "jets" from an experiment<sup>(5)</sup> using a small acceptance calorimeter. However,  $\alpha$  in ref. (5) was determined only by comparison of pA and pp data. The values of  $\alpha$  for the ' $\pi/3$ ' trigger obtained the same way (shown in Fig. 4) are in good agreement with those of ref. (5).

The ' $2\pi$ ' p-nucleus data exhibit event structure that is slightly more isotropic in azimuth than that observed for the corresponding pp data<sup>(8,9)</sup>. This is quantified in Table II using variable planarity as defined in refs. (8,9,11). One also observes an excess of transverse energy emitted backwards in the proton-nucleon center-of-mass as compared to the hydrogen data. This is in qualitative agreement with results reported from many low- $p_t$  nuclear experiments<sup>(1)</sup>. There is little variation in the event structure, as seen by the calorimeter, between Al, Cu and Pb targets (Table II).

Explanations for the observed behavior of  $\alpha$  for both single particle and small azimuthal acceptance data are numerous and varied.<sup>(1,4,5,6,7)</sup> Many authors propose to explain the 'anomalous nuclear enhancement' as due to multiple scattering of the incoming hadron constituents<sup>(6)</sup> and/or 'multijet' production.<sup>(7)</sup> These explanations exploit the standard four jet QCD model for pp scattering and therefore cannot be directly applied to the ' $2\pi$ ' data, for which the model was found inappropriate.<sup>(8,9,11)</sup> There are two recent approaches to description of ' $2\pi$ ' pp data: QCD model of parton showers<sup>(12)</sup> and a low- $p_t$  type multiple Pomeron exchange model.<sup>(13)</sup> The latter is an extension of a dual parton model which accurately explains many aspects of soft hadronic

collisions<sup>(14)</sup> and which has already been generalized successfully to describe soft hadron-nucleus collisions.<sup>(15)</sup> The new pA data present a new challenge for these models.

We conclude that cross sections for the production of high transverse energy events in proton-nucleus collisions increase more rapidly than the atomic mass number. This effect is observed for both small and large azimuthal acceptance data despite the substantial differences in the event structure of the two cases. The nuclear events are less planar than the corresponding pp events and exhibit more energy emitted backward in the proton-nucleon center of mass.

We are grateful for the excellent technical support given us by the Multiparticle Spectrometer facility group led by Dan Green.

This work was supported in part by the U. S. Department of Energy and the National Science Foundation.

- (a) Present address: Higher Institute for Chemistry and Technology,  
Sofia, Bulgaria.
- (b) Present address: University of Maryland, College Park, Maryland
- (c) Present address: George Mason University, Fairfax, Virginia.
- (d) Present address: Dept. of Radiology, University of Chicago, Illinois.
- (e) Present address: Institute for H.E.P. Heidelberg, Fed. Rep. of Germany.
- (f) On leave of absence from University of Warsaw, Poland.

1. For recent reviews on hadron-nucleus interactions, see e.g.:  
W. Busza, Acta. Phys. Polonica B8, 333 (1977);  
A. Bialas, Proceedings of the 9th Multiparticle Symposium, Tabor,  
1978, p. C1.  
Materials of the Fermilab Workshop on  $A^A$  physics, compiled by  
L. Voyvodic, Fermilab preprint Conf. 82-39 (1982).
2. D. Chaney et al., Phys. Rev. Lett. 40, 71 (1978).
3. J. Cronin et al., Phys. Rev. D11, 3105 (1975);  
L. Kluberg et al., Phys. Rev. Lett. 38, 670 (1977);  
D. Antreasyan et al., Phys. Rev. D19, 764 (1979).
4. A. Krzywicki, Phys. Rev. D14, 152 (1976).
5. C. Bromberg et al., Phys. Rev. Lett. 42, 1202 (1979); Nucl. Phys.  
B171, 38 (1980).
6. J. Pumplin and E. Yen, Phys. Rev. D11, 1812 (1975);  
P. M. Fishbane et al., Phys. Rev. D12, 2133 (1976); D16, 122 (1977);  
J. H. Kühn, Phys. Rev. D13, 2948 (1976).
7. F. Takagi, Phys. Rev. Lett. 43, 1296 (1979);  
V. V. Zmushko, Sov. J. Nucl. Phys. 32, 231 (1980);  
D. Treleani and G. Wilk, Nuovo Cimento 60A, 201 (1980);  
U. Sukhatme and G. Wilk, SLAC-PUB-2844 (1981); Phys. Rev. D - in press.



8. "Properties of high transverse energy hadronic events", B. Brown et al., Fermilab preprint CONF. 82/34-EXP (1982), to be published in Proceedings of the 17th Rencontre de Moriond, 1982.
9. B. Brown et al., Phys. Rev. Lett. 49, 711 (1982).
10. P. Rapp et al., Nucl. Inst. and Meth. 188, 285 (1981).
11. C. De Marzo et al., Phys. Letters 112B, 173 (1982).
12. G. C. Fox and R. L. Kelly, LBL-CALTECH preprint, LBL-13985, CALT-68-890 (1982).
13. F. W. Bopp and P. Aurenche, Z. Physics C13, 205 (1982).
14. A. Capella et al., Phys. Letters 81B, 68 (1979); Z. Physics C3, 329 (1980).
15. A. Capella and J. Tran Thanh Van, Phys. Letters 93B, 146 (1980); Z. Physics C10, 249 (1981).

## Figure Captions

Fig. 1. Distributions of the interaction vertex coordinate along the incident beam direction,  $z_{\text{vertex}}$ . The nuclear targets region is marked in black in Figs. a and b.

- (a) 'Interacting beam' data
  - (b) ' $2\pi$ ' acceptance data with  $E_t > 15$  GeV
  - (c) Enlargement of nuclear target region for the ' $2\pi$ ' data.
- The curve superimposed on the data represents a fit described in the text.

Fig. 2. Cross sections,  $d\sigma/dE_t$  for the ' $2\pi$ ' acceptance data as a function of atomic mass number,  $A$ . Different symbols represent various  $E_t$  regions:  $1 < E_t < 11$  GeV (full circles);  $15 < E_t < 16$  GeV (full squares);  $16 < E_t < 17$  GeV (upward triangles);  $17 < E_t < 18$  GeV (open circles);  $18 < E_t < 19$  GeV (downward triangles);  $19 < E_t < 21$  GeV (open squares). The lines represent  $A^\alpha$  fit to the complex nuclei data.

Fig. 3. Exponent  $\alpha$  as a function of  $E_t$  for:

- (a) The ' $2\pi$ ' data
- (b) The ' $\pi/3$ ' data.

The 0.05 systematic error in the value of  $\alpha$  is not included.

Fig. 4. Exponent ' $\alpha_{H_2-Al}$ ' =  $\ln(\sigma_{pAl}/\sigma_{pp})/\ln 27$  as a function of  $E_t$ .

- (a) ' $2\pi$ ' data.
- (b) ' $\pi/3$ ' data (open circles). Data of ref. (5) (triangles) are shown for comparison. They correspond to  $\Delta\Omega^* \approx 1$  sr and are plotted as a function of  $p_t$  of a 'jet'.

The 0.03 systematic error in the value of ' $\alpha_{H_2-Al}$ ' is not included.

Table I

Targets used in this experiment. Number of events taken with different triggers.

Target	A	Thickness (g.cm <sup>-2</sup> )	Number of events *10 <sup>-3</sup>		
			'Inter. beam'	'2 $\pi$ '	' $\pi/3$ '
H <sub>2</sub>	1.0	2.832	32	21.5	5.0
Al	27.0	0.214	0.8	7.5	1.3
Cu	63.5	0.356	0.25	6.5	2.2
Pb	207.2	0.174	0.4	3.7	0.2

Table II

Mean planarity,  $\langle P \rangle$ , and average ratio of transverse to total energy observed in the calorimeter,  $\langle E_t/E_{tot} \rangle$  for the '2 $\pi$ ' data ( $E_t > 15$  GeV).

The value of  $\langle E_t/E_{tot} \rangle$  for H<sub>2</sub> target has been increased by 0.0008 to account for the difference in the acceptance.

Target	$\langle P \rangle$	$\langle E_t/E_{tot} \rangle$
H <sub>2</sub>	0.406 $\pm$ 0.005	0.0628 $\pm$ 0.0005
Al	0.385 $\pm$ 0.004	0.0668 $\pm$ 0.0003
Cu	0.374 $\pm$ 0.005	0.0669 $\pm$ 0.0003
Pb	0.376 $\pm$ 0.004	0.0681 $\pm$ 0.0003




Figure 1




Figure 2




Figure 3




Figure 4